Diversity and specialization of mammalian SWI/SNF complexes

Weidong Wang,¹ Yutong Xue,¹ Sharleen Zhou,² Ann Kuo,¹ Bradley R. Cairns,³ and Gerald R. Crabtree^{1,4}

¹Howard Hughes Medical Institute, Department of Developmental Biology, Stanford University, Stanford, California 94305-5428 USA; ²Department of Molecular and Cell Biology, Howard Hughes Medical Institute, University of California, Berkeley, California 94720-3204 USA; ³Department of Cell Biology, Stanford University, Stanford, California 94305 USA

The SWI/SNF complex in yeast facilitates the function of transcriptional activators by opposing chromatin-dependent repression of transcription. We demonstrate that in mammals SWI/SNF complexes are present in multiple forms made up of 9–12 proteins that we refer to as BRG1-associated factors (BAFs) ranging from 47 to 250 kD. We have isolated cDNAs for human BAF155, BAF170, and BAF60. BAF155 and BAF170 are encoded by separate genes that are both homologs of yeast SWI3. Both contain a region of similarity to the DNA binding domain of myb, but lack the basic residues known to be necessary for interaction with DNA. The two SWI3 homologs copurify on antibody columns specific for either BAF155 or BAF170, indicating that they are in the same complex. BAF60 is encoded by a novel gene family. An open reading frame from yeast, which is highly homologous, encodes the previously uncharacterized 73-kD subunit of the yeast SWI/SNF complex required for transcriptional activation by the glucocorticoid receptor (Cairns et al., this issue). BAF60a is expressed in all tissues examined, whereas BAF60b and BAF60c are expressed preferentially in muscle and pancreas, respectively. BAF60a is present within the 2000-kD BRG1 complex, whereas BAF60b is in a distinct complex that shares some but not all subunits with the BRG1 complex. The observed similarity between mammalian BAF190, BAF170, BAF155, BAF60, and BAF47 and yeast SNF2/SWI2, SWI3, SWI3, SWP73, and SNF5, respectively, underscores the similarity of the mammalian and yeast complexes. However, the complexes in mammals are more diverse than the SWI/SNF complex in yeast and are likely dedicated to developmentally distinct functions.

[Key Words: SWI; SNF; BRG1; BAF; SWP; chromatin]

Received May 23, 1996; revised version accepted July 22, 1996.

Chromatin is actively remodeled during development, as indicated by the observations that the same genetic locus in different tissues varies dramatically in its sensitivity to DNase I (Weintraub and Groudine 1976; Wu and Gilbert 1981) and restriction enzymes (McGhee et al. 1981). During development of specific cell types, chromatin remodeling of certain genes appears to precede the transcriptional activation of the gene (Siebenlist et al. 1986), suggesting that chromatin remodeling may occur in anticipation of developmental transitions. The mechanisms underlying these developmental changes in chromatin structure are unclear. In *Saccharomyces cerevisiae*, the products of five genes, *SWI1*(*ADR6*), *SWI2*(*SNF2*), *SWI3*, *SNF5*, and *SNF6*, are thought to function by remodeling chromatin to aid transcription factors required for specific inducible genes such as *HO* and *SUC2* (Neugeborn and Carlson 1984; Stern et al. 1984; Breeden and Nasmyth 1987; for review, see Winston and Carlson 1992; Carlson and Laurent 1994; Peterson and Tamkun

1995). Recent evidence from both genetic and biochemical studies indicates that these five proteins function together as one multisubunit complex. For example, single and double SWI/SNF mutants were shown to have similar phenotypes (Peterson and Herskowitz 1992). Each of the five SWI/SNF gene products is involved in regulation of the same set of genes (for review, see Peterson and Herskowitz 1992; Winston and Carlson 1992). Immunoprecipitation using antibodies against different SWI/SNF proteins purified all five proteins, indicating they are subunits of one complex. Six additional polypeptides have been found in the complex and two of them were identified as TFG3 and SNF11 (Cairns et al. 1994; Peterson et al. 1994; Treich et al. 1995; Cairns et al. 1996a).

The SWI/SNF complex appears to remodel chromatin structures by an unknown mechanism (for review, see Winston and Carlson 1992; Carlson and Laurent 1994; Wolffe 1993; Peterson and Tamkun 1995). Several genetic suppressors of *swi/snf* mutants have been identified as components of chromatin, such as H2A, H2B, H3, H4 (Hirschhorn et al. 1992, 1995; Prelich and Winston 1993; Kruger et al. 1995). The chromatin structure at the

⁴Corresponding author.

SUC2 promoter is altered in *swi1/snf* mutants, a change that is independent of transcription (Hirschhorn et al. 1992). Moreover, the purified yeast SWI/SNF complex has an ATP-dependent activity that enables it to disrupt mononucleosomes and facilitate binding of GAL4 derivatives to the nucleosome core in vitro (Côté et al. 1994). Thus, one model for the function of the SWI/SNF complex is that it facilitates the ability of transcriptional activators to overcome the repressive effects of chromatin (Travers 1992).

Several lines of evidence suggest that higher eukaryotes contain homologs of the yeast SWI/SNF complex. First, transcriptional activation by the rat glucocorticoid receptor (GR), the *Drosophila bicoid* factor, and *ftz* were shown to be dependent on SWI/SNF function when ectopically expressed in yeast (Laurent and Carlson 1992; Peterson and Herskowitz 1992; Yoshinaga et al. 1992). In addition, two *Drosophila* homologs of the SWI/SNF proteins, *brm* (a homolog of SWI2) and *inr1* (a homolog of SNF5), have been identified and are present in a high-molecular-weight complex similar in size to the yeast complex (Tamkun et al. 1992; Dingwall et al. 1995). *brm* was originally identified as a suppressor of mutations in *polycomb* (Kennison and Tamkun 1988), a repressor of several homeotic genes that is thought to act by regionally compacting chromatin (for review, see Moehrl and Paro 1994). Thus, *brm* itself may be involved in modification of chromatin structure. *brm* and *INR1* are required for proper expression of several homeotic genes (Tamkun et al. 1992; Brizuela et al. 1994; Dingwall et al. 1995). Flies lacking either one die at embryonic stages.

In humans, the two closest relatives of SWI2, BRG1 and hbrm, have been implicated to be functional homologs of SWI2 (Khavari et al. 1993; Muchardt and Yaniv 1993; Chiba et al. 1994). Another protein, INI1/hSNF5, has been reported to be an SNF5 homolog (Kalpana et al. 1994; Muchardt et al. 1995). BRG1 is present in a high-molecular-mass complex of 2 MD, similar in size to the yeast complex (Khavari et al. 1993). Kwon et al. (1994) have reported partial purification of two BRG1-containing complexes from HeLa nuclear extracts by following BRG1 immunoblot reactivity. They showed that their partially purified fractions contain an activity similar to the yeast SWI/SNF complex that can disrupt nucleosomes and facilitate the binding of GAL4 derivatives and TBP to a nucleosome core (Kwon et al. 1994; Imbalzano et al. 1994). We have recently reported complete purification of distinct complexes containing either BRG1 or hbrm using a combination of conventional chromatography and immunoaffinity methods (Wang et al. 1996). The purified BRG1 or hbrm complexes contain 9–12 polypeptides (depending on the cell lines used for the purification). They were referred to as BAFs (BRG1 associated factors). These polypeptides are different from the partially purified fractions reported by Kwon et al. (1994). The discrepancy most likely relates to the fact that the affinity columns allow a single-step purification of 500- to 1000-fold. By microsequencing, immunoblotting, and immunoprecipitation, we showed that the BAF47 subunit is INI1/hSNF5, a mammalian homolog

of SNF5. Like the yeast SWI/SNF complex, the complexes we purified possess an ATP-dependent nucleosome disruption activity and are able to facilitate binding of GAL4-VP16 to mononucleosomes. These data strongly suggest that BRG1- and hbrm-containing complexes are mammalian homologs of the yeast SWI/SNF complex. Purification of SWI/SNF complexes from different mammalian cell lines revealed the presence of multiple forms of SWI/SNF complexes within the same cell or in different cell lines.

As a first step toward understanding the role of the SWI/SNF complex in the developmental remodeling of chromatin, we report the isolation of genes encoding subunits BAF170, BAF155, and BAF60. Our results underscore the similarity of the yeast and mammalian complexes, because BAF155 and BAF170 are homologs of SWI3. Furthermore, the sequence of BAF60 is highly similar to an open reading frame in *Saccharomyces cerevisiae*, that encodes the 73-kD component of the yeast SWI/SNF complex (SWP73) (Cairns et al., this issue), strongly indicating that BAF60 is a functional component of the mammalian complex. BAF60 is encoded by a multigene family that is expressed selectively and participates in distinct SWI/SNF-like complexes. These results imply that although the SWI/SNF complex is likely to have a universal function common to all eukaryotes, this function has been adapted to specialized cellular and developmental needs in multicellular eukaryotes.

Results

Purification of mammalian SWI/SNF complexes

We have purified the SWI/SNF complexes from the human YT cell line using a combination of conventional chromatography and immunoaffinity purification with the BRG1 antibody (Fig. 1; Wang et al. 1996). Two forms of BRG1 complexes, as judged by their different subunit composition, have been purified from this cell line. Complex A, which is the more abundant form in the cell, contains nine subunits with apparent molecular masses of 250, 190, 170, 150, 110, 60, 57, 53, and 47 kD (Fig. 1b, lane 1). The BAF47 subunit of the complex has been identified as INI1/hSNF5 by microsequencing, immunoblotting, and immunoprecipitation (Wang et al. 1996). Purification with an antibody against INI1/hSNF5 yields an almost identical pattern as with the BRG1 antibody (Fig. 1b, lane 2). The data strongly indicate that these nine polypeptides are components of one complex.

Microsequencing and cloning of BAF155 and BAF170 reveals that they belong to the same gene family as yeast SWI3

To define additional subunits of the BRG1 complexes we separated individual subunits of purified complex A by preparative gel electrophoresis and determined the sequence of peptides from BAF155, BAF170, and BAF60 (see Materials and Methods). Four peptide sequences were obtained for human BAF155 after Lys-C endopro-

Figure 1. Purification of mammalian SWI/SNF complex. (a) Purification scheme for mammalian SWI/SNF complex A. Two different forms of mammalian SWI/SNF complexes (A and B) were present in most cell lines and tissues and both BRG1 and hbrm complexes exist in two forms (Wang et al. 1996). The purification scheme is for complex A from cultured cells. Purification using animal tissues requires an extra S300 column. (b) The silver stain gel of the purified human SWI/SNF complex A. The antibodies against BRG1 or hSNF5/INI1/BAF47 were used in immunoaffinity purification and were indicated on top of each lane. The eluted polypeptides were named as BAFxx (see text; xx refers the apparent molecular weight). The band between BAF53 and BAF57 is the immunoglobulin heavy chain.

teinase digestion (Fig. 2). cDNA clones were isolated from a human Jurkat T cell cDNA library (see Materials and Methods). Sequencing and mapping of selected clones revealed a single long open reading frame of 1104 amino acids. All peptide sequences obtained from microsequencing were found in this open reading frame. Immunoprecipitation and immunoblotting using an antibody against a region of the clone confirms that it is indeed the gene encoding human BAF155 (see Fig. 8 and below).

Three high-quality peptide sequences were obtained for human BAF170 after Lys-C endoprotease digestion (Fig. 2a; see Materials and Methods). Degenerate oligos were designed based on the longest peptide sequence (amino acids 733–752 of the cloned BAF170; Fig. 2a) and were used to isolate cDNA clones from a Jurkat T-cell cDNA library under low-stringency washing conditions. Sequencing of the clones obtained revealed a single open reading frame of 1213 amino acids containing all three peptide sequences from microsequencing. Immunoprecipitation and immunoblotting with an antibody against part of the sequence showed that it is indeed the gene encoding human BAF170 (see Fig. 8 and below).

Interestingly, BAF170 and BAF155 are highly similar to each other; they are 62% identical and 77% similar at the amino acid level (Fig. 3a). A search of the NR database at the NCBI with BAF170 and BAF155 protein sequences using the BLAST program revealed two signifi-

cant matches, yeast SWI3 protein (retrospective probability 10^{-73}) and a second open reading frame from *Saccharomyces cerevisiae* identified by the genome sequencing project, YFK7 (SwissProtein bank Accession no. P43609, retrospective probability 10^{-66}). Sequence comparison between SWI3 and YFK7 showed that they are also similar to each other. Overall, they are 30% identical and 52% similar. We will refer to YFK7 as SWI3b (Fig. 3). The similarity between the two yeast SWI3 genes is much lower than the similarity between the two human genes (Table 1).

The conserved regions of the SWI3 gene family include a myb-like tryptophan repeat and a coiled-coil region

The alignment of the two human and the two yeast genes revealed three conserved regions in all four proteins (Fig. 3a,b). Region I is the longest and has the greatest similarity of all three regions. Most of the conserved residues are prolines, hydrophobic, and aromatic amino acids, suggesting that this domain is hidden inside of the complex.

Region II of these proteins is a tryptophan-repeat domain, identified originally in the myb family of proteins (Fig. 3c). This domain is present in two or three tandem repeats in the myb family of proteins from all species and is involved in sequence-specific DNA binding (for review, see Luscher and Eisenman 1990). Recently, this domain was termed SANT domain and was thought to bind DNA as it does in the myb family of proteins (Aasland 1996). However, a bacterial fusion protein containing all three conserved regions of BAF170 has no detectable DNA-binding activity, as determined by gel-shift assays (data not shown).

The conserved region III of the SWI3 gene family contains a predicted leucine-zipper region first recognized in the yeast SWI3 gene (Peterson and Herskowitz 1992). The leucine-zipper or coiled-coil was identified as a dimerization domain for a variety of DNA-binding transcription factors, such as C/EBP, Fos-Jun-GCN4, ATF and Myc-Mad-Max families of proteins (for review, see Johnson and McKnight 1989). Because the mammalian BRG1 complex contains both BAF155 and BAF170 (see below), it is possible that this coiled-coil region serves as the dimerization domain for the complex.

Table 1. The homology between the human and yeast SWI3 gene homologs

Identity/similarity	BAF155	BAF170	ySWI3
BAF155			
BAF177	62/77%		
ySWI3	30/50%	30/49%	
ySWI3b	29/51%	31/51%	30/52%

Each of the two human SWI3 homologs (BAF155 and BAF170) or yeast proteins (SWI3 and SWI3b) were aligned against each other by using the Bestfit program of UWGCG. The calculated percentage of identical or similar amino acids is shown. SWI3b refers to a yeast open reading frame, YFK7 (see text).

Wang et al.

BAF155	1	MAAAAGGGPGTAVGATGFGDSAAAAG	27	618	AGRGWTEQETLLLEALEMYKDDWNVKSEHVGSRTQDECILHFLRLPIED	667
BAF170	1		1	597	ATREWTEQETLLLEALEMYKDDWNVKSEHVGSRTQDECILHFLRLPIED	646
	28	LAVYRRKDGGPATKFWESPETVSQLDSEVRLGKHYKYYHADAPTNKTL	77	668	PYLENSDASLGPLAYQPVFPFSQGNPVMSTVAPLASVVDPRVASAAAKAA	717
	1	MAV.RKKDGGPVNYKYEAADVTQFDNVRLWLKKNYKRYQAEPPTNKSL	49	647	PYLEDSASLGPLAYQPIPPFSQGNPVMSTVAPLASVVDPRVASAAAKSA	696
	78	AGLVVQLLQPOEDAFGKHVTNPAPTKLPACFPMDFKAGGALCHILGAAAYK	127	718	LEEFSSRVREVPLELVEAHVKVQEAARAAGKVDPTYGLESSCIAGTGPD	767
	50	SSLVVQLLQPOEEVFGKHVSNAPLTKLPICPLDFKAGGSLCHILAAAYK	99	697	LEEFSSMKKEVPPTALVEAHVKVQEAARAAGKVDPTYGLESSCIAGTGPD	746
	128	YKNEQGWRRPDLQNPFRMDRNVEMPMNIKTTLVQNNCLTRPNYILIPDID	177	768	EPEKLE..GAEEKMEADPDGQOPEKAENK....VENETDE.....	802
	100	FKSDQGWRRYDFQNPFRMDRNVEMPMNIKTTLVQNNCLSRPNIFLCPEIE	149	747	EPERIEESGNDSEARVEGQATDEKKEPKPEPREGGGAIEEAKKETSAPKK	796
	178	LKLANLKLDIKRHQGTPTDEKSKASHIYPYSSQDDEWLRLPVMRKEK	227	803	.GDKAQDGENEKNSEKQDSEVSEDTKSEKETEENKELSSCTCKERESDT	851
	150	PKLLGKLDIKRHQGTPTDEKSKASHIYPYSSQDDEWLRLPVMRKEK	199	797	DEEKKGEGDSEKSEKSDGDPVDPEK..EKEPKEGQEVVLKEVVESEGER	845
	228	QVLVHWGYPDSYDVTWVSHNDVDAEIEDPPIPEKPKVHVHWKILDTDIFN	277	852	GKKKVEHEISEGNVATAAALASAAKAKHLAAVEERIKISLVALLVET	901
	200	QVLLHWGYYPDSYDVTWVSHNDVDAEIEDPPIPEKPKVHVHWKILDTDIFN	249	846	.KTKVERDIEGNLSTAAAAALAAAVKAKHLAAVEERIKISLVALLVET	894
	278	EMNEEDYEVDEMRKPVSPQRISTKN.EEPVRSP..ERRDRKASANARK	324	902	QMKKLEIKLRHFEGLETIMDRREKALEQQRQQLITERQNFHMEQLKYAEL	951
	250	EMNEEDYEVNDKPNVSRKKISAKTLTDEVNSPDSDRDKG.GNYKK	298	895	QMKKLEIKLRHFEGLETIMDRREKALEYQRRQLADRAQFHMELQKYPEN	944
	325	RKHSPPPPPTPTTSERKKSGKKQASLYGKRRSQKEDEQEDLTDKMDP	374	952	RARQQMEQQHQQNPQQAHQ..HSGGPGLAGLGAAGHPGMMHPQPPPYPL	1000
	299	RKRSPPSP..TPEVKEEKKKGSTPYTKSKRGHREEQEDLTDKMDP	346	945	RARQQHQQHQQHQQPPPPALPPGSGQPIPTGAAGPPAVHGLAVAPASVV	994
	375	TPVPNIIEVVLPKNVNLKDDSENPVKGQTVADLDEQDEETVTAGGKEDE	424	1001	M.....HHQMPPPHPPQGGQIP.....	1017
	347	SPVPNVREVTLPTVNTKDDSESAPVKGQTMIDLDEQDESMETTGKDED	396	995	PAPAGSGAPPGSLGPSEQIGAGSTRGPQQQPPAGAPQPGAVPPGVPPPG	1044
	425	D...PAKGDQSRSDVLGDNVTEQTNHIIIPSYASWFDYNCIHVIERRAL	471	1018	.GPGSMMPGQHMPGRMIPPTVAANIHPGSGGPTPPGMP.....	1053
	397	ENSTGNKGEQTKNPDHEDNVTEQTNHIIIPSYAAWFDYNSVHAIERRAL	446	1045	PHGPSPPFNQQTTPPMMPGAVPGSGHPGAGNAPLGLPFGMPPPPPPPAP	1094
	472	PEFFNGKNSKTPETIYLAYRNFMIDSYRLNPQVEYLTSTACRRNLGTDVCA	521	1054	PMPGNILG.....PRVPLTAPNGMYP	1074
	447	PEFFNGKNSKTPETIYLAYRNFMIDSYRLNPQVEYLTSTACRRNLGTDVCA	496	1095	SIIPFGSLADISINLPAFPNLMGSPSPVRPGTLPPPNLPSMANLHP	1144
	522	VMRVHAGGEQWGLVNYQVDPESRPMAMGPPPTPHFNVLDATP..LACASD	569	1075	..PPPPQQPPPPPPADGVPPPPAPGPPASAA*.....	1104
	497	ISRVHAFLEQWGLINQVDAESRPTPMGPPPTSHFVLADTFSGLVPLQP	546	1145	NLPATTMTSPSLPLGPGGLGSAQAQSPAIVAAVQGNLLPSASPLPDPTPL	1194
	570	LRSPQVPAQQLNFPPEKNKEKPVLDQNPGLRDTIYSKKTLA..KSK.GAS	617			
	547	KTPQQTSAQQMLNFPDKGKEKPTDMQNPGLRDTMYTKKNAPSKSKAAS	596	1195	PPDPTAPSPGTPTVPVPPQ*.....	1213

Figure 2. BAF155 and BAF170 are highly similar to each other. The predicted amino acid sequences for human BAF155 (*top*) and BAF170 (*bottom*). The underlined sequences indicate the peptide sequences obtained from microsequencing after Lys-C digestion. The proteins used for microsequencing are from human KB cells. The sequences are aligned using the Bestfit program of UWGCG. The sequence of BAF155 was obtained from three overlapping cDNA clones. The total length of DNA is 5188 bp, very close to the size of its mRNA on Northern blot (~5.5 kb). The sequence of BAF170 was obtained from a contiguous cDNA clone of 4022 bp. It is shorter than its mRNA determined by Northern blot (~5 kb; see Fig. 4 for Northern blots).

In addition to these conserved regions, the amino termini of BAF155 and BAF170 are highly similar to each other but not to the yeast genes (Fig. 3a). Their carboxyl termini are highly proline-rich and somewhat glutamine-rich. Pro-rich and Glu-rich domains are common in a variety of transcription activators (Courey et al. 1989; Mermod et al. 1989).

BAF155 and BAF170 are expressed in a wide range of tissues

BAF155 and BAF170 mRNA were found in all tissues that have been examined and were expressed at particularly high levels in muscle (Fig. 4). In addition, Western blots of nuclear extracts from cultured cell lines derived from different tissues show that they are expressed in all

cell lines examined (data not shown). Finally, the cDNAs of BAF155 and BAF170 match ESTs obtained from melanocytes, testis, breast, spleen, liver, placenta, infant brain, and fetal brain for BAF155, and ESTs obtained from liver and spleen, ovary, breast, placenta, fetal brain, and testis for BAF170 (Fig. 4).

BAF60 is a homolog of SWP73, a novel SWI/SNF protein

Six peptides obtained from microsequencing of BAF60 from calf thymus matched to an open reading frame [musd15kzl, protein information resource (PIR) accession no. A30222; Fig. 5] adjacent to the mouse glycerol-3-phosphate dehydrogenase gene (Johnston et al. 1989). musd15kzl has no known function, but is expressed ubiquitously in mice. To verify that this gene encodes

Figure 3. Human *BAF155* and *BAF170* belong to the yeast *SWI3* gene family. (a) The schematic presentation of conserved regions between two human genes, *BAF155* and *BAF170* and two yeast genes, *SWI3* and *SWI3b* (*SWI3b* refers to an open reading frame, YFK7, SW accession no. p43609). The shaded area indicates the conserved regions. The numbers above each shaded box indicate the percentage of identical and similar amino acid residues of each region using the corresponding region of *BAF155* as the standard. They were calculated using the Bestfit program of UWGCG. (b) The three highly conserved regions among the *SWI3* gene family of proteins. The myb-like tryptophan repeat and predicted coiled-coil region are indicated by a bracket. The asterisks mark the hydrophobic residues in the coiled-coil that are separated by seven amino acids. The alignment was performed using the Pileup and Pretty program of UWGCG. (c) The alignment of the myb-like tryptophan repeat of the *SWI3* gene family with myb and other transcription factors (see text). We found that the domain is also present as a single repeat in the nuclear receptor corepressor (N-CoR; Horlein et al. 1995) and transcription factor TFC5 (a subunit of TFIIB; Kassavetis et al. 1995). The helix-turn-helix structure was based on the solution structure of c-myb DNA binding domain (Ogata et al. 1992). The asterisks indicate the conserved tryptophans. The arrows indicate the amino acid residues that presumably interact with DNA if the domain functions similarly as the domain in c-myb. The computer search of the NR data bank at the NCBI was done using the BLAST search program.

BAF60, we made rabbit polyclonal antibodies against two different regions. We find that the human version of d15kzl cofractionates with BRG1 through several ion exchange columns (data not shown). In particular, they cofractionate as a 2-MD complex on a Superose 6 sizing column (Fig. 6). Immunoblotting of the complexes purified with antibodies against BRG1, hbrm, BAF155, and BAF170 showed that they all contain a homolog of d15kzl (see Fig. 8, below). Thus, we renamed the

musd15kzl open reading frame mBAF60a. The published musd15kzl sequence has several errors in the carboxy-terminal part of the coding region and has no starting methionine. We corrected those errors and defined its starting methionine (Fig. 5; below).

Comparison of the mBAF60a sequence with known yeast SWI/SNF proteins revealed no significant similarities. However, searching the data bank with the BAF60a protein sequence identified three genes with high BLAST

Figure 4. BAF155 and BAF170 are expressed in a wide range of tissues. (a) Northern blot analysis of mRNA from different human tissues as indicated on the top of each lane. The arrows indicate the presence of the mRNA for either *BAF155*, *BAF170*, or β -actin. (*) The muscle actin that was also detected by the β -actin probe. *BAF155* and *BAF170* cDNAs match many EST sequences in the dBEST data bank at NCBI. These EST sequences are derived from several different tissues at different developmental stages, suggesting *BAF155* and *BAF170* were expressed in these tissues. For *BAF155*, these tissues include: fetal brain (T06718, R95776), infant brain (z43024), testis (T19008, t25695), breast (R50245, t25696, H21971, R55216), placenta (R67673, R24632), melanocyte (N44961, N36960), liver and spleen (R95825, H54140, H53515, R95825). For *BAF170*, the tissues include: fetal brain (M61970), ovary (T50228), testis (T19008), breast (R73206, H24688, R73142), placenta (R24526, R24632), spleen and liver (T64879, T91030, T50242).

scores: one *C. elegans* open reading frame CEZK1128-5 (PIR accession no. z47357, BLAST score 10^{-165}), and two *Saccharomyces cerevisiae* sequences, SCPCPETIT (PIR accession no. s19063, BLAST score 10^{-11}) and YCU2 [SWISS-PROT (SW) accession no. P25632, BLAST score 10^{-10}]. The gene product of SCPCPETIT encodes the SWP73 subunit of the yeast SWI/SNF complex (Cairns et al., this issue). This gene appears to be essential for the function of the SWI/SNF complex, because null mutants have *swi⁻* and *snf⁻* phenotypes. This gene was referred to as SWP73. The sequence YCU2 is 32% identical and 54% similar to SWP73. The alignment of *mBAF60a* with the two yeast genes and the one *C. elegans* gene showed that regions of similarity are present throughout their entire sequences, with many conserved amino acid residues being the same (Fig. 5). *BAF60a* is therefore a mammalian homolog of yeast SWP73 protein and is likely to play roles similar to those of SWP73 in the yeast complex.

Cloning of a family of human genes that are related to *mBAF60a* and SWP73

Based on the sequence of *mBAF60a* we isolated a family of human genes very similar to *mBAF60a* by screening cDNA libraries (see Materials and Methods). We referred to its members as human *BAF60a*, *b*, and *c*. The human *BAF60a* gene is likely the gene corresponding to

mBAF60a in humans, because it has the highest homology to the mouse *mBAF60a* gene at both the DNA (92% identical) and protein level (98% identical and 99% similar). *hBAF60a* is likely a splicing variant because it lacks a region of the mouse gene *mBAF60a* (amino acids 384–423; Fig. 5) that is present in all other homologs.

hBAF60b and *hBAF60c* are highly similar to *BAF60a* and to each other (~70% identical and ~80% similar among all three human genes, Table 2; Fig. 5). *hBAF60b* encodes a predicted protein of 475 amino acids. Immunoblotting with an antibody against *hBAF60b* showed that it is not detectable in the complex purified with the BRG1 antibody (data not shown). Its immunoreactivity was found to fractionate as a 500-kD complex on a Superose 6 sizing column, in contrast to *BAF60a*, which cofractionates with BRG1 as a complex of ~2 MD (Fig. 6; Fig. 9, below). The sequence of *hBAF60c* predicts a protein of 469 amino acids. *BAF60c* is highly expressed in muscle cells (Fig. 7, top), in contrast to *BAF60b* which is expressed in many tissues with the highest level in pancreas (Fig. 7, middle). *BAF60a* was reported to be ubiquitously expressed (Johnston et al. 1989; Fig. 7, bottom).

All three human genes are roughly equal in similarity to yeast SWP73 and SWP73b (Table 2). The alignment of the three human genes with the *BAF60/SWP73* genes from other species showed that they have a divergent amino terminus, but most of the carboxy-terminal part of the protein is conserved. The data strongly suggest that *BAF60* and SWP73 belong to a large gene family whose products could be involved in nucleosome remodeling. Searching of the PIR data bank or using the University of Wisconsin Genetics Computer Group (UWGCG) program to analyze the *BAF60/SWP73* gene family did not reveal any obvious structures or motifs. However, we noticed that the amino-terminal nonconserved region of *BAF60a* contains a proline-glycine rich region that is similar to a region in BRG1 and the *brm* family of proteins.

The *hbrm* complex has a subunit composition similar to the BRG1 complex

We have shown previously that the two mammalian SWI2 homologs, BRG1 and *hbrm*, are present in separate complexes (Wang et al. 1996). As demonstrated above, the BRG1 complex purified with BRG1 antibody contains BAF170, BAF155, and BAF60a. All three subunits are homologs of yeast SWI/SNF proteins. Western blots using antibodies against these proteins confirmed their presence in the BRG1 complex (Fig. 8a). The complex purified with *hbrm* antibody contains INI1/hSNF5 as determined by Western blot, and has a pattern of polypeptides on a silver-stained gel similar to that of the BRG1 complex (Wang et al. 1996; see Fig. 8b, cf. lanes 1,2). We immunoblotted the *hbrm*-associated polypeptides with antibodies against BAF170, BAF155, and BAF60a and found that all three are present in the *hbrm* complex (Fig. 8c, lane 1). Thus, *hbrm* and BRG1 complexes have similar subunit compositions, suggesting that they have similar functions in vivo.

The mammalian SWI/SNF complex contains two SWI3-like subunits

We have shown above that both BAF170 and BAF155 are homologs of the yeast SWI3 gene. Silver stain gels of the complex (Fig. 1b, lanes 1,2; also see Fig. 8b, lanes 1,2) suggest that these two subunits are present in equal molecular ratio in the complex (staining with Coomassie Blue or Ponceau S shows similar results; data not shown). To determine whether BAF170 and BAF155 are in the same complex or separate complexes, we purified BAF170- or BAF155-associated polypeptides using immobilized antibody against BAF170 or BAF155 and visualized the purified proteins by silver-staining of the SDS-PAGE (Fig. 8b, lanes 3,4) and by Western blotting of the eluted polypeptides (Fig. 8c, lanes 2,3). The data demonstrate that the mammalian SWI/SNF complexes containing BAF170 also contain BAF155 and that those containing BAF155 also contain BAF170.

The diversity of mammalian SWI/SNF complexes as evidenced by the presence of a distinct BAF60b complex

Based on the different tissue distributions of the members of the BAF60 gene family, we explored the possibility that they may define biochemically distinct complexes. Using an antibody against the nonconserved amino terminus of BAF60b, we affinity-purified the BAF60b-associated polypeptides (see Materials and Methods). The silver-staining pattern of these polypeptides looks remarkably similar to that of the BRG1 complex purified with the BRG1 antibody (Fig. 9a, lanes 2,3), suggesting that they could be similar or identical. Western blotting revealed that the BAF60b complex does not contain any BAF60a immunoreactivity, indicating that BAF60b and BAF60a are indeed subunits of distinct complexes (Fig. 9b). Conversely, the BAF60b complex contains abundant immunoreactivity for BAF170, BAF155, and BAF47, suggesting that these polypeptides are shared between these complexes (Fig. 9c). In addition, the BAF60b complex contains a subunit (referred to as BAF190x; Fig. 9a, lane 3) that has a similar but slightly slower mobility than the BRG1 subunit. Western blotting showed that it has very low BRG1 immunoreactivity and somewhat higher hbrm reactivity (Fig. 9d). Because several subunits of the mammalian SWI/SNF complex are encoded by gene families, microsequencing will be needed to determine the identities of these BAF60b-associated polypeptides. Preliminary data indicate that in addition to BAF60b, at least one other subunit is different from the BRG1 complex (W. Wang and G.R. Crabtree, unpubl.).

We noted that the molecular mass of all subunits of the BAF60b complex is ~1 MD, which is bigger than the size determined using a Superose 6 sizing column (500 kD). It is possible that the complex was partially dissociated during the fractionation or that the complex exists in a nonglobular shape. We also noted that the *Drosophila* NURF complex (Tsukiyama et al. 1995) fractionates

at a size similar to the BAF60b complex and contains a subunit of 55 kD. We therefore investigated whether BAF60b is a subunit of the human NURF complex. Using an antibody against the human homolog of ISWI, hSNF2L (Okabe et al. 1992; the antibody is a generous gift of Robert Roeder, Rockefeller University, New York), we found that BAF60b and hSNF2L do not have identical chromatographic properties on a Superose 6 sizing column (Fig. 6) or ion-exchange columns (P11 and DE52; data not shown). The complex purified with BAF60b antibody does not contain immunoreactivity for hSNF2L (Fig. 9d).

Discussion

Structural and functional similarities between the mammalian and yeast SWI/SNF complexes

We previously described the complete purification of mammalian SWI/SNF complexes and demonstrated that the complexes are heterogeneous and that they contain homologs of yeast SWI2/SNF2 (BRG1 or hbrm) and SNF5 (INI1/hSNF5). The complexes containing BRG1 have nucleosome-disruption activities similar to those of the yeast SWI/SNF complex (Wang et al. 1996). In this report we demonstrate that BAF170, BAF155, and BAF60 are homologs of yeast SWI3 and SWP73 (see also Cairns et al., this issue). These results and our earlier work establish the close relationship between mammalian and yeast SWI/SNF complexes and strongly support a conserved and universal function for the SWI/SNF complex in all eukaryotes.

Mammalian SWI/SNF complexes contain two different SWI3 homologs

BAF155 and BAF170 are different homologs of the yeast SWI3 protein. These two subunits are highly similar to each other (62% identical and 77% similar) and both are found in the same complex, because purification of the complex with antibody to one subunit contains the other (Fig. 8). The two proteins also appear to be in equimolar ratio based on visualization by silver staining or Coomassie-blue staining. The presence of a leucine zipper in both BAF155 and BAF170 suggests that these subunits could be contributing to a dimer-like structure by forming heterodimers.

The prediction that the complex has a dimer-like structure can be further supported by molecular weight considerations. The molecular masses of either yeast or mammalian SWI/SNF complex has been determined at 2 MD by Superose 6 sizing columns (Khavari et al. 1993; Peterson et al. 1994). The total molecular weight of all subunits of either the yeast complex (Cairns et al. 1994; Côté et al. 1994) or the mammalian complex is ~1 MD, half the size determined by sizing columns. One explanation for this apparent paradox is that some or all subunits of the complex are present in two copies. Our finding that the mammalian SWI/SNF complex contains two different homologs of SWI3 provides the strong evidence for such a model.

Wang et al.

Identification of a myb-like tryptophan repeat domain in the SWI3 family of proteins

We identified a myb-like tryptophan-repeat domain in all SWI3 proteins (Fig. 3c). This domain was identified originally as tandem repeats in the myb family of proteins (for review, see Luscher and Eisenman 1989) and

was recently termed the SANT domain and proposed to be a DNA binding domain (Aasland 1996). The domain is also present in components of two other known chromatin-remodeling complexes, the ADA2/ADA3/GCN5 histone acetyltransferase complex (Horiuchi et al. 1995; Brownell et al. 1996; Candau and Berger 1996) and the NURF complex (Tsukiyama et al. 1995; Tsukiyama and

hBAF60b	SIPKTPAC...	...APAQPPM	PAQRRLKRR	KMADKVLPR	IRELVESQA	YMDLLAFERK	LDQTTARKRM	ETQEALKKPL	TQK.....	RK	137	
mBAF60a	APQIQVQVQ.	QAV	QNRNHNNAKK	KMADKILPQR	IRELVESQA	YMDLLAFERK	LDQTTMRKRL	DIQEALKRPI	KQK.....	RK	142	
hBAF60a	APQIQVQVQ.	QAV	QNRNHNNAKK	KMADKILPQR	IRELVESQA	YMDLLAFERK	LDQTTMRKRL	DIQEALKRPI	KQK.....	RK	142	
hBAF60c	SSAPARQSQA	QSQQGPEPTA	PARSRSAKRR	KMADKILPQR	IRELVESQA	YMDLLAFERK	LDQTTMRKRV	DIQEALKRPM	KQK.....	RK	142	
cBAF60	APLPGQTAQI	QGNRGPPPK	KKK	RYADKILQPK	VRELVESQA	YMDLLAFERK	LDSTTTTKRI	DVQEALKRQP	KIK.....	KR	107	
ySWP73	NFFHAKKDP	VNQDKANNAS	QITPTVPESH	PSDMVIRPDL	A.ELIEELYS	EQQLVDSEKR	LDHFTHLRNL	HKRMVQAWE	RSKLSQEFLY	PHLNFPNVKF	124	
ySWP73b		MV	TQTNVPVPTY	PTDAYIPTYL	PDDKVSMLAD	KKKLIEMDSR	LDLYLTRRL	DTSINLPTNT	ETKDHP....	PNKEM	73	
Consen	AP---QVQVQ-	Q-Q---QPAV	Q-RN--AKKK	KMADKILPQR	IRELVESQA	YMDLLAFERK	LDQTTMRKRL	DIQEALKRPI	KQK-----	-----PN-RK		
hBAF60b	LRIYISNTFS	PSKAEQDSAG	TAGTPGGTPA	GDKVASWELR	VEGRLLDD..	.PS.....	KQKRKFSSFF	KSLVIEL..			204	
mBAF60a	LRIYISNTFN	PAKSDAED..	G	EGTVASWELR	VEGRLLDD..	.AALSKYDAT	KQKRKFSSFF	KSLVIEL..			205	
hBAF60a	LRIYISNTFN	PAKSDAED..	G	EGTVASWELR	VEGRLLDD..	.SALSKYDAT	KQKRKFSSFF	KSLVIEL..			205	
hBAF60c	LRIYISNTFN	PAKSDAED..	S	DGSIASWELR	VEGRLLDD..	.PS.....	KQKRKFSSFF	KSLVIEL..			198	
cBAF60	LRIYISHTFI	AGKEPEKEG.	D	DASVPMWELR	VEGRLLDDMQ	HPTVGANPRP	APKRKFSSFF	KSLVIEL..			174	
ySWP73	LRIYISNVSE	NQPMQMDTNN	EADLM....	ALENATWTR	TEGRLLDNVQ	AND.....	PAREKFSSFI	ESINVDKKNK	ENDNVPSTKF	NAAPEENATE	212	
ySWP73b	LRIYVYNTTE	SSPRS.DSGT	PAD.....	.SGKTTWTLR	TEGRLLDHE	SA.....	NCKHPFSEFL	EGYAVDEKRL	K.....	...PLGMGKK	142	
Consen	LRIYISNTFN	PAKSDA-DG-	-AD-----G	-G-VASWELR	VEGRLLDD-Q	-PALSKYDAT	KQKRKFSSFF	KSLVIELK--	-----	---P-----		
hBAF60b					..DKELYGPD	GHLVEWYWMP	TTQ.ETDGFQ	VKRPGDLNVR	CTILLML..D	HQPPQYKLDP	259	
mBAF60a					..DKELYGPD	NHLVEWHRTA	TTQ.ETDGFQ	VKRPGDVNVR	CTVLLML..D	YQPPQFKLDP	260	
hBAF60a					..DKELYGPD	NHLVEWHRTA	TTQ.ETDGFQ	VKRPGDVNVR	CTVLLML..D	YQPPQFKLDP	260	
hBAF60c					..DKELYGPD	NHLVEWHRTA	TTQ.ETDGFQ	VKRPGDLSVR	CTILLML..D	YQPPQYKLDP	253	
cBAF60					..DKELYGPD	NHLVEWHRTA	TTQ.ETDGFQ	VKRPGDRPVK	CTILLML..D	YQPMKFKLHP	229	
ySWP73	GPSDKKLNLN	LPLQFSLPNG	DNSTTTNTDQ	NNATMGEEA	KKDMSSTTPK	LESVKMAYDP	NNPVDFDGLD	IKRVGSENV	CTHSILRKSS	PPEPFMSYSP	312	
ySWP73b	RKRDSSLSLP	LNLQPEFYND	QDSTMGDNDN	GEDED	SASAEAREEI	VDALWNYDE	NNVVEFDGID	IKRQKDNLR	CSITITQLR.G	VDGKQVQYSP	246	
Consen	---D--L-L-	L-LQ---N-	--ST---D-	-----GE---	--DKELYGPD	NHLVEWHRTA	TTQVETDGFQ	VKRPGD-NVR	CTILLML--D	YQPPQYKLDP		
hBAF60b	RLARLLGVHT	QTRAAIMQAL	WLYIKHNOL					QD	GHREYVINC	RYFQIETSCG	RLRFSEIPMK	320
mBAF60a	RLARLLGIHT	QTRPVIIQAL	WQYIKTHKL					QD	PHREFVLCD	KYLQIIFESQ	RMKFSEIPQR	321
hBAF60a	RLARLLGIHT	QTRPVIIQAL	WQYIKTHKL					QD	PHREFVICD	KYLQIIFETQ	RMKFSEIPQR	321
hBAF60c	RLARLLGIHT	QSRSAIQAAL	WQYVKTNRL					QD	SHDKYINGD	KYFQIIFDCP	RLKFSEIPQR	314
cBAF60	RLARVLGIAA	ETRPRIIEAL	WQYIKTHKL					QD	PQDRDITNND	LFLEQCFGVS	KRMFMEIPQR	290
ySWP73	QLTATIGLKG	QTSDDAIFST	YKYIHLNELL		TNDESAFENL	MGNRNHNNSN	TSTSKMLDAA	SSQVSVVKLD	TQITLTLFSS	LKESPDPTMK	402	
ySWP73b	NLATLIGMOT	GSVNDAYYST	YKWEILNNLF	VTEQTEADQG	SNDAAEDSNE	NNNKNGAGDD	DGVEGSTPKD	KPELGEVKLD	SILQKVL...	..DTNAAHLP	331	
Consen	RLARLLGIHT	QTRP-IIQAL	WQYIKTNKL		-ND-----N-	--N-N-----	-----QD	PHERE-VNCD	KYLQIIFE--	RMKFSEIPQR		
hBAF60b	LAGE.....	..LQHPDPIV	INHVISVDPN	DQKKT.ACYD	IDVEVDPLK	AQMSNFLAST	TNQQE.....		TASLDVKIHE	TIESINQLKT	396	
mBAF60a	LHAL.....	..LMPPEPII	INHVISVDPN	DQKKT.ACYD	IDVEVDPLK	TQMSFLLST	ASQQE.....		TATLDNKIHE	TIESINQLKT	397	
hBAF60a	LHAL.....	..LMPPEPII	INHVISVDPN	DQKKT.ACYD	IDVEVDPLK	TQMSFLLST	ASQQE.....		TATLDN....		383	
hBAF60c	LTAL.....	..LLPPDPIV	INHVISVDPN	DQKKT.ACYD	IDVEVEPLK	QMSFLLST	ANQQE.....		TSPLDSKIHE	TIESINQLKI	390	
cBAF60	LHQE.....	..LQQPDPLV	LNHILQRPDD	GQDKTSACVD	IDVELEDPVK	QQMANFVHNO	TNAND.....		TQLLDQKIFD	LVDQINEMKL	367	
ySWP73	LTDLLSLINS	THLLPLOPIE	IDYTVRV...	DKASTYGEVL	LDIEVDVNA	LKFNNTQRES	QIGAAELNEN	ARELEQIKPK	TALQKELTS	VLSNLHESNK	499	
ySWP73b	LMNVVQTVNK	L.VSPLPEPI	IDYTTDL...	SKPTTYGATD	LDVDVSHI...	..LHQPPQPP	NLQKEE.ETD	AEDTAKL...	REITK	TALQNLSSAQ	414	
Consen	LHAL---N-	--L-PPDPI-	INHVISVDPN	DQKKTAYCYD	IDVEV-DPLK	-QM-SFLLST	ANQQEE----	A-----	IA-LD-KIHE	TIE-INQLKT		
hBAF60b	ORDFMLSFT	DEQDFIQEWL	RSQRDLKII	.TDVTGNPEE	ERRAEFYHQP	WAQEAAGRHI	FAKVQQRQE	LEQVLGIRLT	*	475		
mBAF60a	QREMLSLP.	ERSGFINDWL	QSCRCGLKTM	.TDVVGNPPEE	ERRAEFYHQP	WAQEAACRYT	YSKVQQRQE	LEQALGIRNT	*	475		
hBAF60a												
hBAF60c	QRDFMLSFSR	DFKGYVQDLI	RSQSRDLKVM	.TDVAGNPPEE	ERRAEFYHQP	WSQEAASRYT	YCKIQQRQE	LEQSLVVRNT	*	469		
cBAF60	RRDFELRFEN	ERSGFINKVV	VQSNLDLKTLL	.TESSGDGES	DRYATTYSTT	DTDEGVSRYM	YQKIQQRRAE	LEQSLGIRNN	*	446		
ySWP73	RYRFFKKISE	DPVKALNECT	ASTSNALKVL	SGDEGYNEDM	VRRANFY..K	ENEAMLRNT	EVILSNGRM*			566		
ySWP73b	KYQFFHELHL	HRPRTLTHYL	WSKQNELVL	QGDQYFNEDA	ARTSDIYSNN	NNDRLMGNT	SLLYSQGL*			483		
Consen	-RDF-LSFS-	DPSGFI--WL	-SQSRDLKV-	-TDVVGNPPEE	ERRAEFY-OP	WAQEA-VRY-	YSKVQQRQE	LEQ-LGIRNT	-			

Figure 5. (See facing page for legend.)

Figure 6. BAF60b fractionates as a separate complex on Superose 6 sizing column. Western blotting analysis after fractionation of nuclear extract of human YT cells (DE52/0.3M fraction) by a Superose 6 sizing column (Pharmacia). Antibodies against BRG1, BAF60a, BAF60b, and hSNF2L were used. The presence of these proteins is indicated by arrows.

Wu 1995), suggesting that it may play a common role in nucleosome-remodeling. However, we have not been able to demonstrate any DNA binding activity using a recombinant protein containing all three conserved regions of BAF170 (data not shown). Careful inspection of amino acid residues that should contact DNA based on the solution structure of c-myc (Ogata et al. 1992) revealed that three out of five residues in the SWI3 SANT domain are hydrophobic where these residues are charged or hydrophilic in c-myc (Fig. 3c). Furthermore, the DNA-protein cross-linking studies showed that the subunits in yeast SWI/SNF complex that contact DNA are not SWI3 (Quinn et al. 1996). Thus, the SANT do-

main in the SWI3 family of proteins may play a role other than DNA-binding.

BAF60 belongs to a large gene family that includes a novel yeast SWI/SNF subunit, SWP73

We cloned three closely related human *BAF60* homologs: *hBAF60a*, *b* and *c*. These human genes have two yeast homologs (Fig. 5). Cairns et al. (this issue) demonstrate that one of the two yeast homologs encodes a previously uncharacterized SWI/SNF subunit, SWP73. Null mutations in *SWP73* have a phenotype that is similar to other *SWI/SNF* deletions. Furthermore, yeast lacking *SWP73* are unable to activate transcription in response to glucocorticoids in yeast strains containing an introduced glucocorticoid receptor gene and a glucocorticoid responsive promoter. Thus, the work in both mammals and yeast indicates that BAF60a/SWP73 is an integral subunit of the SWI/SNF complex.

Mammalian SWI/SNF complexes and pol II holoenzyme

Recently, the yeast SWI/SNF complex was reported to copurify with SRB proteins and was suggested to be an integral part of the yeast pol II holoenzyme (Wilson et al. 1996). This result is different from previous reports from two other groups using different purification schemes for the SWI/SNF complex (Cairns et al. 1994; Côté et al. 1994). In those purifications, the SWI/SNF complex does not contain other components. To date we have isolated cDNA clones for seven out of nine subunits of mammalian SWI/SNF complexes and none is similar to the subunits of yeast pol II holoenzyme (W. Wang and G.R. Crabtree, unpubl.). The preparation of mammalian pol II holoenzyme has been described recently (Ossipow et al. 1995; Chao et al. 1996). Its pattern on silver-stained gels (Chao et al. 1996) appears to be distinct from that of the purified mammalian SWI/SNF complex. Thus, when us-

Figure 5. Sequence alignment of the *BAF60/SWP73* gene family. The sequence alignments of BAF60/SWP73 proteins from mammals, *C. elegans*, and *Saccharomyces cerevisiae*. The underlined sequences represent those that are either identical or similar to the peptide sequences obtained from microsequencing. The protein used for microsequencing is from calf thymus, and is therefore possibly not identical to the reported sequences from mouse or human. One peptide sequence is of high quality: RELVPE (1). The remaining sequences have ambiguity at some positions: KSL[V,g]IE (2); RLLGI (3); KLAGL[L,x]Q (4); KI(H,x)(T,d)IE(F,t)I (5); RAAF[Y,i](H,e)QP(W,l)A(Q,i)EA (6). The amino acids in parentheses represent differences between the cloned gene (uppercase) and the peptide sequence from microsequencing (lowercase). Peptide 3 is more similar to BAF60a, whereas peptide 4 and 6 are more similar to BAF60b. The remaining peptides do not distinguish between different BAFs. Thus, it appears that BAF60b is present in a large amount in BRG1 complex of calf thymus. The sequence mBAF60a refers to musd15kzl (Johnston et al. 1989). The published mouse sequence is 3101 bp, which is similar to the size of the mRNA (3.2 kb) as determined by Northern blot. It has no starting methionine and also contains a sequencing error that results in its carboxyl terminus being out of frame. Our human BAF60a clone is 2841 bp (mRNA ~4kb) but it contains an extra 150 bp at the 5' end. A stop codon was found in frame to the predicted starting methionine. However, the methionine at codon14 could be the starting codon because it has a better Kozak sequence (3/5 vs. 0/5). The human BAF60b cDNA clone is 1946 bp, smaller than its mRNA (~3 kb). A stop codon was found upstream in frame to the predicted starting methionine. There is also a methionine in between, but it is present in a GC-rich region and in a context of a poor Kozak sequence. The predicted starting codon make the BAF60b protein similar in size to BAF60a and BAF60c. The BAF60c cDNA clone is 1724 bp (the mRNA is ~2 kb). A stop codon was found upstream in frame with the predicted methionine. cBAF60 refers to an open reading frame identified by the *C. elegans* genome sequencing project, CEZK1128_5. ySWP73 refers to an open reading frame from *Saccharomyces cerevisiae*, SCPPETIT. This sequence was shown to encode the SWP73 subunit of the yeast SWI/SNF complex (Cairns et al., this issue). ySWP73b refers to the yeast sequence YCU2. These sequences were aligned using the Pileup and Pretty programs of UWGCG.

Table 2. The homology between mammalian BAF60 and yeast SWP73 gene families

Identity/similarity	mBAF60a	hBAF60a	hBAF60b	hBAF60c	cBAF60	yswp73
mBAF60a						
hBAF60a	98/99%					
hBAF60b	66/79%	66/78%				
hBAF60c	70/82%	71/82%	69/82%			
cBAF60	55/73%	56/73%	52/72%	55/74%		
yswp73	28/51%	27/51%	27/50%	27/50%	25/49%	
yswp73b	24/46%	26/49%	23/45%	26/49%	23/47%	32/54%

Each of the mammalian BAF60 homologs or yeast SWP73 homologs were aligned against each other using the Bestfit program of UWGCG. The calculated percentage of identical or similar amino acids are shown as % identical/% similar. mBAF60a refers to a known gene, musd15kzl. cBAF60 refers to a *C. elegans* open reading frame CEZK1128_5. ySWP73 refers to a yeast open reading frame SCPCPETIT. ySWP73b refers to a yeast open reading frame YCU2 (see text for data-bank access numbers).

ing stringent washing conditions (0.2 M Guanidine chloride, 0.5 M NaCl) for the purification of the of SWI/SNF complexes from mammalian cells, pol II holoenzyme components are not deleted.

Diversity and specialization of mammalian SWI/SNF complexes

Previously, we have purified two forms of BRG1 complexes and two forms of hbrm complexes from a single mammalian cell line. We showed that BRG1 and hbrm are present in separate complexes. The subunit composition of the BRG1 complex is also different when purified from several differentiated cell types (Wang et al. 1996). In this work, we find that of the three BAF60 homologs, at least BAF60a and BAF60b are in separate complexes. Thus, each mammalian SWI/SNF complex may contain a distinct BAF190 (BRG1 or hbrm) and a distinct BAF60 (a, b, or c) subunit. In contrast, the two SWI3 homologs, BAF170 and BAF155, together with the SNF5 homolog, BAF47, appeared to be the common sub-

units present in each complex we purified. They may form a common core with which the different homologs of BAF190 and BAF60 associate in a combinatorial manner. This would give rise to a diverse population of complexes consisting of similar but distinct subunits.

Why are there so many SWI/SNF-like complexes in higher eukaryotes? One possibility is that chromatin remodeling is a highly regulated developmental process in which different SWI/SNF-like complexes are molded to the needs of a differentiated cell, as suggested from the first studies mapping DNase sensitivity of chromatin (Weintraub and Van Lente 1974). The relatively selective expression of BAF60 homologs in different tissues suggests that there may well be tissue-specific chromatin remodeling activity. Further characterization of the chromatin remodeling complexes from different tissues should begin to clarify this issue.

Materials and methods

Purification of the mammalian SWI/SNF complexes

The purification procedure of different SWI/SNF complexes has been described in detail in Wang et al. (1996). Briefly, the nuclear extracts were fractionated over P-11 (Whatman) and DE52 (Whatman) ion-exchange columns (see Fig. 1a) and a S300 (Pharmacia) sizing column. The presence of the SWI/SNF complex was followed by Western blotting with antibodies against BRG1, hbrm, hSNF5, and BAF60b (Wang et al. 1996). The fractionated extract was then loaded onto antibody columns that were made by cross-linking different antibodies to Protein-Sepharose beads (Pharmacia) according to the protocols in Harlow and Lane (1988). The columns were thoroughly washed using 0.5 M buffer D [0.5 M KCl, 20 mM HEPES (pH 7.9), 0.25 mM EDTA, 10% glycerol, 0.1% Tween 20], and 0.2 M guanidine hydrochloride. The SWI/SNF complex was then eluted under mild acidic conditions (0.2 M glycine at pH 2.5) directly into neutralizing buffer, 1 M TrisHCl (pH 8.0). The purification of the BAF60b complex is similar to the purification of the BRG1 complex, except that the column was washed at 0.1 M guanidine hydrochloride.

Microsequencing of the subunits of the BRG1 complex

The peak fractions containing BRG1 complex A from BRG1 antibody columns were pooled and the proteins concentrated by

Figure 7. AF60b and BAF60c have different tissue-expression patterns. Northern blot analysis of mRNA from different human tissues as indicated on the top of each lane. This is the same blot used in the experiments of Fig. 4. The arrows indicate the presence of the mRNA for BAF60a, BAF60b, and BAF60c.

Figure 8. Mammalian SWI/SNF complex contains two SWI3 homologs and an SWP73 homolog. (a) Western blot analysis of load, flowthrough (FT), and eluted fractions of the BRG1 antibody column. Each fraction is indicated on the top of each lane. The arrows indicate BRG1, BAF170, BAF155, and BAF60a. (b) The silver-staining gel of complexes purified with different antibody columns as marked on top of each lane. The lines indicate the bands corresponding to each BAF. The arrows indicate BAF170 and BAF155, which are present in above-stoichiometric amounts in complexes purified by their own antibodies. This is in contrast to complexes purified with BRG1, IN11/hSNF5, or hbrm antibodies in which the amount of BAF170 and BAF155 are stoichiometric (see Fig. 1b, lanes 1,2; Fig. 8b, lanes 1,2). The data suggest that some BAF170 and BAF155 is not associated with BRG1 or hbrm complexes and could be immunopurified preferentially

by their antibodies. Another possibility is that complexes containing homodimers of either BAF170 or BAF155 are present in cells and are purified preferentially by the corresponding antibodies. Our previous findings that SWI/SNF complexes purified from some cell lines contain substoichiometric amounts of BAF170 support this view (Wang et al. 1996). A third possibility is that some BAF170 and 155 are associated loosely with mammalian SWI/SNF complexes, and they dissociate during purification. (c) Western blot analysis of complexes purified with different antibody columns as shown on the top of each lane. The arrows indicate the immunoreactivity of each BAF. We have frequently observed that the BAF47/hSNF5 antibody recognizes two bands. We suspect that the protein might be phosphorylated.

TCA precipitation. The proteins were separated by preparative SDS-PAGE (7.5%). The gel was then cut into two halves. The gel containing lower-molecular-weight proteins (smaller than the 66-kD marker) was transferred to PVDF membrane and visualized by Ponceau S-staining. The band containing BAF60 was excised and digested with trypsin (Boehringer-Mannheim, sequencing grade) overnight. The other half of the gel containing higher-molecular-weight proteins (larger than the 97 kD marker) was negatively stained with ZnCl_2 and developed in imidazole (Zhou and Admon 1995). The bands containing BAF170 and BAF155 were excised and digested with Lys-C endoproteinase (WAKO). The peptides eluted from the PVDF membrane or the gel slice were separated by reverse-phase HPLC and sequenced by automatic amino acid sequencer.

Cloning of the BAF155 subunit

BAF155 was purified using human KB cells as the source. The sequence of four peptides obtained after Lys-C digestion did not match any known genes in the NR databank at NCBI searched with the BLAST program. Luckily, one long peptide sequence (corresponding to amino acids 924–944, Fig. 2) was found to match a human expressed sequence tag (EST) sequence from The Institute of Genomic Research (TIGR) databank (ID no. EST18238, corresponding to amino acids 880–949 of the cloned BAF155). BLAST search with this EST sequence revealed that it is similar to the yeast SWI3 gene. Two oligonucleotides (nucleotides 3–21 and 214–196) were used as primers to amplify the EST sequence by PCR. Human cDNA from Jurkat T-cells was used as the template for the PCR reaction.

The PCR product obtained was then used as a template to make ^{32}P -labeled probes to screen cDNA libraries. The hybridized filters were washed twice under low stringency conditions

($2\times$ SSC and 0.1% SDS, room temperature) and twice under high-stringency conditions, $0.1\times$ SSC and 0.1% SDS at 65°C . Over 100 positive clones have been identified. They were analyzed by restriction digestion and PCR mapping and classified into different groups. The representative clones from each group were sequenced, and three overlapping clones were found to encode a single open reading frame containing all peptide sequences from microsequencing.

Cloning of BAF170

BAF170 was purified from human KB cells, and three high-quality peptide sequences were obtained after Lys-C endoproteinase digestion (Fig. 2a). These sequences do not match any known genes or EST sequences in the data bank. Two degenerate oligos were synthesized based on the longest peptide sequence (amino acids 728–748 of the cloned BAF170, Fig. 2a). The oligo sequences are: 5'-AA(G/A)GCIGA(C/T)CCIGCITT(C/T)GG-3' and 5'-GTGTICIGCIATICC-3'. They were used for internal PCR reactions using human cDNA from Jurkat T-cells as the template. After amplification (2 rounds, 25 cycles each round), the PCR products were separated by 8% nondenaturing PAGE. The DNA fragment with the expected size of 60 bp was excised and cloned into a PGEM-T vector (Promega). Four random clones containing the insert were sequenced and found to contain the predicted sequence based on the four middle amino acids. A 23-nucleotide oligo was then synthesized based on the amplified sequence: 5'-GGTCTGGAAAGCAGTGGCATCGC-3'. It was ^{32}P -labeled by T4 kinase and used as a probe to screen cDNA libraries under low-stringency washing conditions ($2\times$ SSC, 0.1% SDS, 42°C , $2\times$ 15 min; room temperature $2\times$ 15 min). Over 20 clones were obtained. One group of clones was found to contain the predicted sequence based on the peptide

Figure 9. The diversity of mammalian SWI/SNF complexes as evidenced by the presence of a distinct BAF60b complex. (a) Silver-staining gel of the affinity-purified complex using either BRG1 antibody (lane 2) or BAF60b antibody (lane 3) from human YT cells. The polypeptides that have similar mobilities in both complexes are marked. Since no microsequencing has been done for the BAF60b-associated polypeptides, their identities cannot be conclusively determined by simple use of Western blotting data. They were referred to as BAFx. However, the presence of strong immunoreactivity of BAF155, BAF170 and BAF47/hSNF5 suggest that these polypeptides are also subunits of the BAF60b complex. The BAF110 band is missing in this preparation of the BRG1 complex. We find that BAF110 can sometimes be washed away when using a BRG1 antibody column, but it is always present when using BAF47/hSNF5 antibody column. (b–d) Western blot analysis of the load, flowthrough (FT), and elute fractions of the BAF60b affinity column, respectively, as indicated on the top of each lane. The presence of the immunoreactivity was marked with an arrow. (*) The cross-linked IgG coming off the affinity column that was recognized by the secondary antibody. We noticed that the immunoreactivities of BAF60a, BAF155, BAF170, and BAF47/hSNF5 are all detectable in the flowthrough fraction of the BAF60b antibody column, whereas the reactivities for BRG1 and hbrm were not detected. The data imply that these subunits could be components of complexes other than BRG1 and hbrm. We found that the BRG1 and hbrm complexes are very “sticky” and can be precipitated nonspecifically by several unrelated antibodies. The complexes could then be washed off under high-stringency washing conditions. This could be the reason why BRG1 and hbrm complexes were depleted in the BAF60b antibody flowthrough fractions and were also present at low levels in the elute fractions.

sequence. The longest clone was sequenced and found to contain a cDNA of 4022 bp. A single open reading frame of 1213 amino acids was identified that contained all three peptide sequences obtained by microsequencing.

Cloning the BAF60 family of proteins

The BAF60 subunit was purified from calf thymus. The peptides obtained were either identical or similar to a published sequence in the data bank, musd15kzl (Fig. 4). To clone the human homologs of BAF60, the mouse sequence was used to search the human EST sequence data bank at TIGR and dbEST data bank at the National Center for Biotechnology Information (NCBI). Five EST sequences were found with high scores.

Human EST	Data bank	Probability	Gene
EST64722	TIGR	10^{-70}	BAF60a
THC48547	TIGR	10^{-87}	BAF60b
F05831	EMBL	10^{-50}	BAF60c
THC44070	TIGR	10^{-124}	BAF60c
EST04790/T06901	TIGR/dbest	10^{-18}	Unknown

Oligonucleotides were designed based on the first three EST sequences for PCR amplification of each sequence. The oligos for each sequence are: EST64722 (nucleotides 2–20 and 224–207), THC48547 (nucleotides 7–24 and 316–299), and F05831 (nucleotides 28–46 and 234–217). The cDNA from Jurkat cells was used as a template for PCR reaction. The amplified PCR products were used as probes to obtain cDNAs as described above (see cloning of BAF155).

Preparation of antibodies and immunoaffinity purifications

Rabbit polyclonal antibodies against BRG1, hbrm and hSNF5/INI1/BAF47 have been described previously (Wang et al. 1996).

The antibodies used in this work are all rabbit polyclonal antibodies. The immunogens are fusion proteins containing either glutathione *S*-transferase (GST) or the maltose binding protein (MBP) fused to a region of the cloned subunit. The region used for immunization for BAF155 is amino acids 872–949 (fused to MBP); for BAF170, amino acids 744–857 (fused to MBP); for BAF60a, two antibodies were made against two different regions: amino acids 80–276 (fused to MBP) and 280–475 (fused to GST); for BAF60b, amino acids 26–93 (fused to MBP). The antibodies were all affinity-purified using the corresponding immunogen as affinity ligands (Harlow and Lane 1988). All except BAF60a antibodies work on immunoprecipitation. They were cross-linked to Protein-A Sepharose (Pharmacia) with DMP (Harlow and Lane 1988) and were used for affinity purification of complexes (Wang et al. 1996).

Northern blot

The human multitissue blot was from Clontech and was probed with a 4.4-kb fragment of cDNA for *hBAF155* and a 2.6-kb fragment of *hBAF170* cDNA. The full-length cDNAs for *hBAF60a*, *b* and *c* were used as probes. The β -actin probe was provided by the manufacturer. The probes were labeled by 32 P-dCTP using Klenow and random hexamers as primers (1–2 million cpm/ml). After hybridization, the blot was washed at high-stringency conditions ($0.1\times$ SSC, 0.1% SDS, 2×10 min at 55°C) and exposed to Kodak X-ray films for 1–3 days. The same blot was washed and reprobable multiple times using different probes as described above.

Acknowledgments

We thank R. Tjian for his strong support throughout this project. W.W. has been supported by a Fellowship from the Cancer Research Fund of the Damon Runyon–Walter Winchell Foun-

dation, DRG-1182. This work has been supported by the Howard Hughes Medical Institute and a grant from the National Institutes of Health to G.R.C. We are very grateful to Dr. Robert Roeder for the hSNF2L antibody. We appreciate highly those scientists who have contributed to the EST data bases at NCBI and TIGR. We thank S. Ho, H. Bayle, and L. Timmerman for critical reading of the manuscript. We thank J. Oberlindacher for help in preparation of this manuscript. The genes cloned in this work have been submitted to GenBank under accession numbers U66615 (BAF155); U66616 (BAF170); U66617 (human BAF60a); U66618 (BAF60b); U66619 (BAF60c), and U66620 (mouse BAF60a).

The publication costs of this article were defrayed in part by payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 USC section 1734 solely to indicate this fact.

References

- Aasland, R. 1996. The SANT domain: a putative DNA-binding domain in the SWI-SNF and ADA complexes, the transcriptional co-repressor NCoR and TFIIB. *TIBS* **21**: 87–88.
- Breedon, L. and K. Nasmyth. 1987. Cell cycle control of the yeast HO gene: cis- and trans-acting regulators. *Cell* **48**: 389–397.
- Brizuela, B.J., L. Elfring, J. Ballard, J.W. Tamkun, and J.A. Kenison. 1994. Genetic analysis of the brahma gene of *Drosophila melanogaster* and polytene chromosome subdivisions 72AB. *Genetics* **137**: 803–813.
- Brownell, J.E., J. Zhou, T. Ranalli, R. Kobayashi, D.G. Edmondson, S.Y. Roth, and C.D. Allis. 1996. Tetrahymena histone acetyl transferase A: A homolog to yeast GCN5p linking histone acetylation to gene activation. *Cell* **84**: 843–851.
- Cairns, B.R., Y.-J. Kim, M.H. Sayes, B.C. Laurent, and R.D. Kornberg. 1994. A multisubunit complex containing the SWI1/ADR6, SWI2/SNF2, SWI3, SNF5, and SNF6 gene products isolated from yeast. *Proc. Natl. Acad. Sci.* **91**: 1950–1954.
- Cairns, B.R., N.L. Henry, and R.D. Kornberg. 1996a. TFG3/TF30/ANC1, a component of the yeast SWI/SNF complex that is similar to the leukemogenic proteins ENL and AF-9. *Mol. Cell. Biol.* **16**: 3308–3316.
- Cairns, B.R., R.S. Levinson, K.R. Yamamoto, and R.D. Kornberg. 1996b. Essential role of Swp73p in function of the yeast Swi/Snf complex. *Genes & Dev.* (this issue).
- Canda, R. and S.L. Berger. 1996. Structural analysis of yeast putative adaptors. *J. Biol. Chem.* **271**: 5237–5245.
- Carlson, M. and B.C. Laurent. 1994. The SNF/SWI family of global transcriptional activators. *Curr. Opin. Cell Biol.* **6**: 396–402.
- Chao, D.M., E.L. Gadbois, P.J. Murray, S.F. Anderson, M.S. Sonu, J.D. Parvin, and R.A. Young. 1996. A mammalian SRB protein associated with an RNA polymerase II holoenzyme. *Nature* **380**: 382–385.
- Chiba, H., M. Muramatsu, A. Nomoto, and H. Kato. 1994. Two human homologues of *Saccharomyces cerevisiae* SWI2/SNF2 and *Drosophila* brahma are transcriptional coactivators cooperating with the estrogen receptor and the retinoic acid receptor. *Nucleic Acids Res.* **22**: 1815–1820.
- Côté, J., J. Quinn, J.L. Workman, and C.L. Peterson. 1994. Stimulation of GAL4 derivative binding to nucleosomal DNA by the yeast SWI/SNF complex. *Science* **265**: 53–60.
- Courey, A.J., D.A. Holtzman, S.P. Jackson, and R. Tjian. 1989. Synergistic activation by the glutamine-rich domains of human transcription factor Sp1. *Cell* **59**: 827–836.
- Dingwall, A.K., S.J. Beek, C.M. McCallum, J.W. Tamkun, G.V. Kalpana, S.P. Goff, and M.P. Scott. 1995. The *Drosophila* snr1 and brm proteins are related to yeast SWI/SNF proteins and are components of a large protein complex. *Mol. Biol. Cell* **6**: 777–791.
- Harlow, E. and D. Lane. 1988. *Antibodies: A laboratory manual*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Hirschhorn, J.N., S.A. Brown, C.D. Clark, and F. Winston. 1992. Evidence that SNF2/SWI2 and SNF5 activate transcription in yeast by altering chromatin structure. *Genes & Dev.* **6**: 2288–2298.
- Hirschhorn, J.N., A.L. Bortvin, S.L. Ricupero-Hovasse, and F. Winston. 1995. A new class of histone H2A mutations in *Saccharomyces cerevisiae* causes specific transcriptional defects in vivo. *Mol. Cell. Biol.* **5**: 1999–2009.
- Horiuchi, J., N. Silverman, G.A. Marcus, and L. Guarente. 1995. ADA3, a putative transcriptional adaptor, consists of two separable domains and interacts with ADA2 and GCN5 in a trimeric complex. *Mol. Cell. Biol.* **15**: 1203–1209.
- Horlein, A.J., A.M. Naar, T. Heinzel, J. Torchia, B. Gloss, R. Kurokawa, A. Ryan, Y. Kamei, M. Sorderstrom, C.K. Glass, and M.G. Rosenfeld. 1995. Ligand-independent repression by the thyroid hormone receptor mediated by a nuclear receptor co-repressor. *Nature* **377**: 397–404.
- Imbalzano, A.N., H. Kwon, M.R. Green, and R.E. Kingston. 1994. Facilitated binding of TATA-binding protein to nucleosomal DNA. *Nature* **370**: 481–485.
- Johnson, P.F. and S.L. McKnight. 1989. Eukaryotic transcriptional regulatory proteins. *Ann. Rev. Biochem.* **58**: 799–839.
- Johnston, L.A., M.A. Kotarski, D.J. Jerry, and L.P. Kozak. 1989. An ubiquitously expressed gene 3.5 kilobases upstream of the glycerol-3-phosphate dehydrogenase gene in mice. *Mol. Cell. Biol.* **9**: 935–945.
- Kalpana, G.V., S. Marmon, W. Wang, and G.R. Crabtree. 1994. Binding and stimulation of HIV-1 integrase by a human homologue of yeast transcription factor SNF5. *Science* **266**: 2002–2006.
- Kassavetis, G.A., S.T. Nguyen, R. Kobayashi, A. Kumar, E.P. Geiduschek, and M. Pisano. 1995. Cloning, expression, and function of TFC5, the gene encoding the B' component of the *Saccharomyces cerevisiae* RNA polymerase III transcription factor TFIIB. *Proc. Natl. Acad. Sci.* **92**: 9786–9790.
- Kennison, J.A. and J.W. Tamkun. 1988. Dosage-dependent modifiers of polycomb and antennapedia mutations in *Drosophila*. *Proc. Natl. Acad. Sci.* **85**: 8136–8140.
- Khavari, P.A., C.L. Peterson, J.W. Tamkun, D.B. Mendel, and G.R. Crabtree. 1993. BRG1 contains a conserved domain of the SWI2/SNF2 family necessary for normal mitotic growth and transcription. *Nature* **366**: 170–174.
- Kruger, W., C.L. Peterson, A. Sil, C. Coburn, G. Arents, E.N. Moudrianakis, and I. Herskowitz. 1995. Amino acid substitutions in the structured domains of histones H3 and H4 partially relieve the requirement of the yeast SWI/SNF complex for transcription. *Genes & Dev.* **9**: 2770–2779.
- Kwon, H., A.N. Imbalzano, P.A. Khavari, R.E. Kingston, and M.R. Green. 1994. Nucleosome disruption and enhancement of activator binding by a human SWI/SNF complex. *Nature* **370**: 477–481.
- Laurent, B.C. and M. Carlson. 1992. Yeast SNF2/SWI2, SNF5, and SNF6 proteins function coordinately with the gene-specific transcriptional activators GAL4 and Bicoid. *Genes & Dev.* **6**: 1707–1715.
- Luscher, B. and R.N. Eisenman. 1990. New light on myc myb. Part II. myb. *Genes & Dev.* **4**: 2235–2241.
- McGhee, J.D., W.I. Wood, M. Dolan, J.D. Engel, and G. Felsen-

- feld. 1981. A 200 base pair region at the 5' end of the chicken adult beta-globin gene is accessible to nuclease digestion. *Cell* **27**: 45–55.
- Mermod, N., E.A. O'Neill, T.J. Kelly, and R. Tjian. 1989. The proline-rich transcriptional activator of CTF/NF-1 is distinct from the replication and DNA binding domain. *Cell* **58**: 741–753.
- Moehrle, A. and R. Paro. 1994. Spreading the silence: Epigenetic transcriptional regulation during *Drosophila* development. *Dev. Genetics* **15**: 478–484.
- Muchardt, C. and M. Yaniv. 1993. A human homologue of *Saccharomyces cerevisiae* SNF2/SWI2 and *Drosophila* brm genes potentiates transcriptional activation by the glucocorticoid receptor. *EMBO J.* **12**: 4279–4290.
- Muchardt, C., C. Sardet, B. Bourachot, C. Onufryk, and M. Yaniv. 1995. A human protein with homology to *Saccharomyces cerevisiae* SNF5 interacts with the potential helicase hbrm. *Nucleic Acids Res.* **23**: 1127–1132.
- Neugeborn, L. and M. Carlson. 1984. Genes affecting the regulation of SUC2 gene expression by glucose repression in *Saccharomyces cerevisiae*. *Genetics* **108**: 845–858.
- Ogata, K., H. Hojo, S. Aimoto, T. Nakai, H. Nakamura, A. Sarai, S. Ishii, and Y. Nishimura. 1992. Solution structure of a DNA-binding unit of Myb: A helix-turn-helix-related motif with conserved tryptophans forming a hydrophobic core. *Proc. Natl. Acad. Sci.* **89**: 6428–6432.
- Okabe, I., L.C. Bailey, O. Attree, S. Srinivasan, J.M. Perkel, B.C. Laurent, M. Carlson, D.L. Nelson, and R.L. Nussbaum. 1992. Cloning of human and bovine homologs of SNF2/SWI2: a global activator of transcription in yeast *S. cerevisiae*. *Nucleic Acids Res.* **20**: 4649–4655.
- Ossipow, V., J.-P. Tassan, E.A. Nigg, and U. Schibler. 1995. A mammalian RNA polymerase II holoenzyme containing all components required for promoter-specific transcription initiation. *Cell* **83**: 137–146.
- Peterson, C.L. and I. Herskowitz. 1992. Characterization of the yeast SWI1, SWI2, and SWI3 genes, which encode a global activator of transcription. *Cell* **68**: 573–583.
- Peterson, C.L., A. Dingwall, and M.P. Scott. 1994. Five SWI/SNF gene products are components of a large multisubunit complex required for transcriptional enhancement. *Proc. Natl. Acad. Sci.* **91**: 2905–2908.
- Peterson, C.L. and J.W. Tamkun. 1995. The SWI-SNF complex: A chromatin remodeling machine. *TIBS* **20**: 143–146.
- Prelich, G. and F. Winston. 1993. Mutations that suppress the deletion of an upstream activating sequence in yeast: Involvement of a protein kinase and histone H3 in repressing transcription in vivo. *Genetics* **135**: 665–676.
- Quinn, J., A.M. Fryberg, R.W. Ganster, M.C. Schmidt, and C.L. Peterson. 1996. DNA-binding properties of the yeast SWI/SNF complex. *Nature* **379**: 844–847.
- Siebenlist, U., D.B. Durand, P. Bressler, N.J. Holbrook, C.A. Norris, M. Kamoun, J.A. Kant, and G.R. Crabtree. 1986. Promoter region of interleukin-2 gene undergoes chromatin structure changes and confers inducibility on chloramphenicol acetyltransferase gene during activation of T cells. *Mol. Cell. Biol.* **6**: 3042–3049.
- Stern, M., R. Jensen, and I. Herskowitz. 1984. Five SWI genes are required for expression of the HO gene in yeast. *J. Mol. Biol.* **178**: 853–868.
- Tamkun, J.W., R. Deuring, M.P. Scott, M. Kissinger, A.M. Patatucci, T.C. Kaufman, and J.A. Kennison. 1992. *brahma*: a regulator of *Drosophila* homeotic genes structurally related to the yeast transcriptional activator SNF2/SWI2. *Cell* **68**: 561–572.
- Travers, A.A. 1992. The reprogramming of transcriptional competence. *Cell* **69**: 573–575.
- Treich, I., B.R. Cairn, T. Santos, E. Brewster, and M. Carlson. 1995. SNF11, a new component of the yeast SNF-SWI complex that interacts with conserved region of SNF2. *Mol. Cell. Biol.* **15**: 4240–4248.
- Tsukiyama, T., C. Daniel, J. Tamkun, and C. Wu. 1995. ISWI, a member of the SWI2/SNF2 ATPase family, encodes the 140 kd subunit of the nucleosome remodeling factor. *Cell* **83**: 1021–1026.
- Tsukiyama, T. and C. Wu. 1995. Purification and properties of an ATP-dependent nucleosome remodeling factor. *Cell* **83**: 1011–1020.
- Wang, W., J. Côté, Y. Xue, S. Zhou, P.A. Khavara, S.R. Biggar, C. Muchardt, G.V. Kalpana, S.P. Goff, M. Yaniv, J.L. Workman, and G.R. Crabtree. 1996. Purification and biochemical heterogeneity of the mammalian SWI/SNF complex. *EMBO J.* (in press).
- Weintraub, H. and F. Van Lente. 1974. Dissection of chromosome structure with trypsin and nucleases. *Proc. Natl. Acad. Sci.* **71**: 4249–4253.
- Weintraub, H. and M. Groudine. 1976. Chromosomal subunits in active genes have an altered conformation. *Science* **193**: 848–856.
- Wilson, C.J., D.M. Chao, A.N. Imbalzano, G.R. Schnitzler, R.E. Kingston, and R.A. Young. 1996. RNA polymerase II holoenzyme contains SWI/SNF regulators involved in chromatin remodeling. *Cell* **84**: 235–244.
- Winston, F. and M. Carlson. 1992. Yeast SNF/SWI transcriptional activators and the SPT/SIN chromatin connection. *TIG* **8**: 387–391.
- Wolffe, A.P. 1994. Transcription: In tune with the histones. *Cell* **77**: 13–16.
- Wu, C. and W. Gilbert. 1981. Tissue-specific exposure of chromatin structure at the 5' terminus of the rat preproinsulin II gene. *Proc. Natl. Acad. Sci.* **78**: 1577–1580.
- Yoshinaga, S.K., C.L. Peterson, I. Herskowitz, and K.R. Yamamoto. 1992. Roles of SWI1, SWI2, and SWI3 proteins for transcriptional enhancement by steroid receptors. *Science* **258**: 1598–1604.
- Zhou, S. and A. Admon. 1995. Enzymatic digestion of proteins in zinc chloride and Ponceau S stained gels. In *Techniques in protein chemistry VI* (ed. J.W. Crabbs), pp. 161–167. Academic Press, San Diego, CA.

Diversity and specialization of mammalian SWI/SNF complexes.

W Wang, Y Xue, S Zhou, et al.

Genes Dev. 1996, **10**:

Access the most recent version at doi:[10.1101/gad.10.17.2117](https://doi.org/10.1101/gad.10.17.2117)

References

This article cites 56 articles, 25 of which can be accessed free at:
<http://genesdev.cshlp.org/content/10/17/2117.full.html#ref-list-1>

License

Email Alerting Service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).