Wnt pathway components orient a mitotic spindle in the early *Caenorhabditis elegans* embryo without requiring gene transcription in the responding cell

Ann Schlesinger, Christopher A. Shelton, Julin N. Maloof,¹ Marc Meneghini, and Bruce Bowerman²

Institute of Molecular Biology, University of Oregon, Eugene, Oregon 97403 USA; ¹Department of Biochemistry and Biophysics, University of California, San Francisco, California 94143 USA

In a four-cell-stage *Caenorhabditis elegans* embryo, Wnt signaling polarizes an endoderm precursor called EMS. The polarization of this cell orients its mitotic spindle in addition to inducing endodermal fate in one daughter cell. Reducing the function of Wnt pathway genes, including a newly identified *GSK-3 β* homolog called *gsk-3*, disrupts endoderm induction, whereas only a subset of these genes is required for proper EMS mitotic spindle orientation. Wnt pathway genes thought to act downstream of *gsk-3* appear not to be required for spindle orientation, suggesting that *gsk-3* represents a branch point in the control of endoderm induction and spindle orientation. Orientation of the mitotic spindle does not require gene transcription in EMS, suggesting that Wnt signaling may directly target the cytoskeleton in a responding cell.

[**Key Words:** Cell polarity; endoderm; inductive signaling; mesoderm; cytoskeleton; *wingless*]

Received February 9, 1999; revised version accepted June 8, 1999.

During asymmetric cell division, correct partitioning of segregated cytoplasmic determinants must be coordinated with the axis of cell division (Lu et al. 1998; Hawkins and Garriga 1999). Such an asymmetric division occurs in the *Caenorhabditis elegans* embryo when a four-cell-stage embryonic cell, or blastomere, called EMS divides along the anterior/posterior (A/P) axis. After replication of the EMS centriole, the two centrosomes migrate to align initially along the left/right (L/R) axis (Fig. 1B). Shortly before mitosis, the EMS nucleus and centrosomes rotate 90°. The ensuing mitotic spindle forms along the A/P axis, and EMS divides into a posterior daughter, E, that produces endoderm, and an anterior daughter, MS, that makes mesoderm (Sulston et al. 1983; Hyman and White 1987; Fig. 1A).

Positioning of the centrosomes and polarization of endodermal potential within EMS both require signaling from the blastomere posterior to EMS, called P₂ (Schierenberg 1987; Goldstein 1992, 1993, 1995a,b; Fig. 1D). The role of P₂ signaling was demonstrated using blastomeres isolated from wild-type embryos and cultured in vitro. If EMS is isolated early in its cell cycle and cultured alone, it produces two MS-like daughters that make only mesodermal tissue, such as muscle. Only when P₂ is placed in contact with EMS does EMS make

endoderm in addition to mesoderm (Goldstein 1992). Moreover, if EMS is cultured alone, its mitotic spindle elongates along the axis established by centrosome migration, without any rotation (Goldstein 1995b). Contact with P₂ early in the EMS cell cycle induces rotation of the EMS nucleus and centrosomes, orienting the mitotic spindle perpendicular to the plane of contact between the two cells (Goldstein 1995b; Fig. 1D). Therefore P₂ signaling polarizes EMS such that a posterior daughter is born next to P₂ and produces endoderm. EMS remains competent to polarize endoderm potential in response to P₂ signaling for longer than it remains competent to orient its mitotic spindle (Goldstein 1992, 1995a,b), suggesting that the signals or their responses may be distinct.

Genetic insight into the nature of P₂ signaling has come, in part, from the identification of five *mom* genes required for endoderm specification. Mothers homozygous for a recessive mutation in any one of these essential genes produce mutant embryos that lack endoderm because both EMS daughters adopt an MS-like fate (Rochelleau et al. 1997; Thorpe et al. 1997). Another maternally expressed gene, *pop-1*, is required to repress endoderm fate in the anterior EMS daughter, as both EMS daughters in *pop-1* mutant embryos adopt an E fate (Lin et al. 1995). The identities of three *mom* genes and *pop-1* are known and all are related to Wnt pathway genes. *mom-1* is similar to porcupine (*porc*), *mom-2* to *wingless*

²Corresponding author.

E-MAIL bbowermann@molbio.uoregon.edu; FAX (541) 346-5891.

Figure 1. Proper EMS mitotic spindle orientation requires signaling from P_2 . All embryos are shown with anterior to the left. (A) Schematic lateral view of early *C. elegans* four- and eight-cell-stage embryos showing blastomere positions and lineages. P_2 , the smaller cell, is posterior to its sister EMS. On division, only the posterior EMS daughter, E, is in contact with P_2 . C and P_3 are the daughters of P_2 . Sister cells are connected by short lines. (B) Nomarski micrograph of a ventral view of a four-cell-stage embryo. EMS centrosomes are indicated by arrows; the centrosomes are visible by their exclusion of cytoplasmic yolk droplets. (Left) The centrosomes oriented along the L/R axis after migration; (middle) the centrosomes have rotated 90° to lie along the A/P axis and the EMS spindle has begun to elongate, oriented toward P_2 ; (right) E and MS after completion of cytokinesis. (C) Lateral views of four-cell-stage embryos. EMS centrosomes are indicated by arrows. (Left) A wild-type embryo with the EMS spindle oriented toward P_2 ; (right) a *mom-3* mutant embryo with the EMS spindle incorrectly oriented. (D) Isolated wild-type blastomeres in culture. EMS and P_2 blastomeres were separated immediately after P_1 division then recombined. P_2 is on the right. EMS mitotic spindle angles were measured as described in Material and Methods. (Left) Results from experiments in which P_2 and EMS were recombined early during the EMS cell cycle while EMS was competent to respond to the P_2 signal. All spindles in these experiments oriented correctly, with

spindle angles between 1° and 12°. (Right) Results from experiments in which P_2 and EMS were recombined after EMS could no longer respond to the P_2 signal. The range of spindle angles from 10° to 90° indicates that EMS spindles in these experiments were randomly oriented with respect to P_2 .

(*wg*), and *mom-5* to *frizzled* (*fz*) (Rocheleau et al. 1997; Thorpe et al. 1997). *pop-1* is related to *Drosophila pangolin/dTCF* (*pan*) and mammalian TCF/LEF family members that encode putative transcription factors regulated by Wnt signaling (Lin et al. 1995, 1998). Therefore, the widely conserved Wnt signal transduction pathway is required for endoderm induction during embryogenesis in *C. elegans* (for review, see Lu et al. 1998; Hawkins and Garriga 1999).

To determine whether Wnt signaling regulates mitotic spindle orientation in addition to polarizing endoderm potential in EMS, we asked if rotation of the EMS nucleus and its associated centrosomes requires the *mom* and other Wnt pathway genes. Previous studies have documented a mitotic spindle orientation defect in an eight-cell-stage blastomere called ABar in *mom* mutant embryos, as well as extensive defects in mitotic spindle orientation and blastomere positioning in *mom-1*; *mom-2* double mutants (Rocheleau et al. 1997; Thorpe et al. 1997). EMS spindle orientation has not been examined in mutant embryos lacking the function of individual *mom* genes. Using blastomeres isolated from mutant embryos and cultured in vitro, we show that a subset of the *C. elegans* Wnt pathway genes known to be

required for endoderm specification also are required for proper orientation of the EMS spindle. In addition, we have identified a *C. elegans GSK-3 β* gene that may function as a branchpoint for endoderm specification and spindle orientation. Finally, we present evidence that the induced rotation of the EMS mitotic spindle does not require gene transcription, suggesting that Wnt signaling may target the cytoskeleton directly in a responding cell.

Results

Wnt pathway components mom-1/porc and mom-5/fz, but not mom-2/wg, are required for proper spindle orientation

Both endoderm specification and proper EMS spindle orientation require signaling from P_2 (see introductory section). To determine if the *mom* genes, which are required for endoderm induction, also are required for rotation of the EMS nucleus and its pair of centrosomes, we first observed the orientation of the EMS spindle in developing mutant embryos using Nomarski optics. In wild-type embryos, the EMS centrosomes initially align along the L/R axis of the embryo. Before nuclear envelope break-

down, the EMS nucleus/centrosome complex rotates to form a spindle along the A/P axis (Hyman and White 1987; Fig. 1B). The spindle then elongates, with only the posterior daughter of EMS born in contact with P₂. In *mom-1/porc*, *mom-3*, and *mom-5/fz* mutant embryos, we observed defects in EMS spindle orientation. The axis of orientation varied, exhibiting significant dorsal/ventral or L/R components (Fig. 1C; data not shown). As the mitotic spindles elongated, however, they invariably assumed a mostly A/P orientation before the end of cytokinesis (data not shown). The A/P axis of EMS also is its longest axis, and therefore spindles may rotate passively to accommodate their length, making it somewhat difficult to quantitate this phenotype when observing intact embryos within their chitinous eggshells.

To facilitate scoring EMS mitotic spindle orientation in wild-type and mutant embryos, we isolated blastomeres in culture medium after removing the eggshells (Goldstein 1995b; see Materials and Methods). We isolated the smaller, posterior-most blastomere in a two-cell-stage embryo, called P₁, and allowed it to divide into its daughters, P₂ and EMS. The orientation of the mitotic spindle in EMS was then measured relative to the plane of contact between EMS and P₂. As illustrated in Figure 2, the range of wild-type EMS spindle angles is within 19° of a line perpendicular to the plane of contact with P₂. The mitotic spindles in EMS blastomeres from *mom-1/porc*, *mom-3*, and *mom-5/fz* mutant embryos, however, generated a much wider range of angles, showing nearly random orientations (Fig. 2). The EMS spindle in *mom-2/wg* mutant blastomeres oriented more normally (Fig. 2). *mom-4* experiments yielded a wide range of EMS spindle angles (0°–90°). Only 4/26 *mom-4* EMS spindles were abnormal, however, whereas the majority oriented

properly. Therefore *mom-1/porc*, *mom-5/fz*, *mom-3*, and perhaps to a lesser degree *mom-4* are required for both endoderm induction and proper EMS spindle orientation, but *mom-2/wg* appears to be required more specifically for endoderm induction.

A subset of Wnt pathway components are required for EMS spindle orientation

Because *mom-1/porc* and *mom-5/fz* are required for EMS spindle orientation, we examined mutant blastomeres lacking the function of other Wnt pathway components required for polarizing endoderm potential in EMS, as well as blastomeres lacking the function of a *C. elegans* GSK-3 β gene that we identified using degenerate PCR (Fig. 3; Materials and Methods). The predicted GSK-3 protein is 76% similar and 69% identical to the GSK-3 β encoded by *Drosophila zeste white 3* (*zw3*). We investigated the function of *gsk-3* using RNA-mediated interference (RNAi), a method that involves microinjecting double-stranded exonic RNA into the syncytial nematode ovary to reduce the function of the corresponding maternally expressed gene (Fire et al. 1998; see Materials and Methods). We refer to embryos from mothers microinjected with *gsk-3* RNA as *gsk-3*(RNAi) embryos. After isolation of *gsk-3*(RNAi) P₁ blastomeres, we observed that EMS spindles were randomly oriented (Fig. 2). We conclude that, in addition to *mom-1/porc* and *mom-5/fz*, the Wnt pathway component *gsk-3* also is required to orient the EMS mitotic spindle.

GSK-3 β is thought to interact with both β -catenin/Armadillo (Arm) and adenomatous polyposis coli (APC), possibly phosphorylating one or both proteins (Rubinfeld et al. 1996; Miller and Moon 1997). *C. elegans*

Figure 2. A subset of identified Wnt pathway genes are required for proper orientation of the EMS spindle. Isolated P₁ blastomeres from different mutant strains were cultured in vitro; on P₁ division, P₂ and EMS were left in contact and the angle of the EMS spindle was measured. Spindle angles were scored and recorded as described in Materials and Methods. The left-most quadrant shows wild-type control data, with spindle angles ranging from 0° to 19°. The range of spindle angles in EMS blastomeres from embryos mutant for *mom-1/porc* (1°–88°), *mom-5/fz* (0°–90°), *mom-3* (1°–90°), and *gsk-3* (1°–90°) are wider than wild-type, indicating a lack of orientation with respect to the plane of contact with P₂. The defects in EMS may represent a randomization of spindle orientation, or an incompletely penetrant failure to rotate the nucleus/centrosome complex. In some cases, endoderm induction occurred in the absence of correct EMS spindle orientation, and vice versa. Examples of such situations have been indicated for *mom-1/porc*, *mom-3*, and *mom-5/fz*. A (+) next to a spindle angle mark indicates the production of endoderm in spite of abnormal EMS spindle orientation and a (–) indicates the lack of endoderm in some partial embryos with normal EMS spindle angles. *mom-4* experiments yielded a wide range of EMS spindle angles (0°–90°). Only 4/26 were abnormal, whereas the majority of spindles oriented properly. A one-sided Student *t*-test comparing wild-type with *mom-4* yields a *P*-value of 0.18, indicating that the *mom-4* data set is not significantly different from wild type. Spindle angle ranges for *mom-2/wg* (0°–28°), *apr-1/APC* (3°–34°), *wrm-1/arm* (1°–17°), *pop-1/pan* (1°–32°), and the *apr-1/wrm-1* double (0°–13°) were similar to wild type. Mammalian and *Drosophila* genes names are aligned below their *C. elegans* homologs. *mom-3* and *mom-4* have not yet been cloned.

spindles are oriented normally in mutant embryos lacking maternal expression of *pop-1/pan* (Fig. 2), which encodes an HMG-domain protein targeted by Wnt signaling during endoderm induction (Lin et al. 1995, 1998; Rocheleau et al. 1997; Thorpe et al. 1997). Therefore, the genes acting downstream of *gsk-3* in the Wnt pathway are not required for proper EMS spindle orientation even when both *wrm-1/arm* and *apr-1/APC* functions are compromised simultaneously.

Next we examined the role of *gsk-3* during endoderm specification. Most *gsk-3*(RNAi) embryos produce intestine (82%; Table 1). Approximately one-fifth lack endoderm and instead make large clumps of cells similar in appearance to the extra pharyngeal tissue observed in *mom* mutant embryos (Table 1; data not shown). To determine the origins of the endoderm made in *gsk-*

Table 1. Cell fate in *gsk-3(RNAi)* embryos

A. Polarizing optics ^a		
	Percent producing endoderm (total)	
Wild type intact	100 (50/50)	
Wild type isolated E	100 (32/32)	
Wild type isolated MS	0 (0/10)	
Wild type isolated C	0 (0/16)	
Wild type C and E ablated	0 (0/6)	
<i>gsk-3</i> (RNAi)	82 (300/364)	
<i>gsk-3</i> (RNAi) isolated E	56 (14/25)	
<i>gsk-3</i> (RNAi) isolated MS	0 (0/8)	
<i>gsk-3</i> (RNAi) isolated C	71 (15/21)	
<i>gsk-3</i> (RNAi) C and E ablated	0 (0/8)	
B. Tissue production ^b		
	Percent making pharyngeal muscle	Percent making body wall muscle
Wild type isolated E	0 (0/9)	0
<i>gsk-3</i> (RNAi) isolated E	30 (4/12)*	38 (3/8)*

^aUsing polarizing optics to detect the presence of rhabditi granules, endoderm was scored in terminally differentiated embryos following ablation of indicated blastomeres. The total number of ablations scored is indicated in parenthesis.

^bTissue production was scored using tissue-specific antibodies on terminally differentiated, fixed embryos following ablation of indicated blastomeres. Embryos were stained with 3NB12, a monoclonal antibody that recognizes pharyngeal muscle cells, or 5.6, one that recognizes body wall muscle. An asterisk indicates that all positive staining occurred only in embryos that did not make endoderm.

3(RNAi) embryos, we used a laser microbeam to kill all but one early blastomere, subsequently scoring the ability of the isolated blastomere to produce endodermal or mesodermal cell types (Table 1). Whereas E always makes endoderm in wild-type embryos, 44% of *gsk-3(RNAi)* E blastomeres failed to make endoderm and instead produced pharyngeal and body wall muscle cells, consistent with E adopting an MS-like fate (Table 1). Therefore *gsk-3* is required positively for the specification of endoderm in *C. elegans*. As genes thought to act downstream of *gsk-3* are required only for endoderm specification, whereas *gsk-3* itself is required for both endoderm specification and EMS spindle orientation, we conclude that *gsk-3* represents a branchpoint by which Wnt signaling regulates both processes.

The production of endoderm by E in only 56% of *gsk-3(RNAi)* embryos does not account for the observation that 82% of intact mutant embryos produce endoderm. Accounting for this discrepancy, we found that a dorsal-posterior eight-cell-stage descendant of P₂ called C ectopically makes endoderm in most *gsk-3(RNAi)* embryos. These results indicate that in wild-type embryos *gsk-3* negatively regulates endoderm production in C while positively specifying endoderm in EMS (see Discussion).

EMS spindle orientation requires mom-1/porc and mom-3 in P₂, and mom-5/fz and gsk-3 in EMS

Having established a requirement for *mom-1/porc*, *mom-3*, *mom-5/fz* and *gsk-3* in spindle orientation, we constructed genetic chimeras to determine if these and other genes are required in P₂ to signal, or in EMS to respond to signaling. We isolated blastomeres from mutant and wild-type embryos, and in each experiment re-associated a mutant cell with a wild-type partner (see Materials and Methods). When placed in contact with a wild-type EMS, *mom-1/porc*, *mom-2/wg*, and *mom-3* mutant P₂ blastomeres failed to properly orient EMS mitotic spindles, whereas EMS cells from these mutant embryos were capable of responding to wild-type P₂ cells (Fig. 4). EMS blastomeres from *mom-5/fz* and *gsk-3* mutants did not respond competently to wild-type P₂ signaling, but *mom-5/fz* and *gsk-3* mutant P₂ blastomeres properly oriented the spindle of wild-type EMS cells (Fig. 4). Orientation of the EMS spindle was normal in the *mom-4* mosaics. Therefore, if *mom-4* is required for spindle orientation (Fig. 2), apparently either blastomere can provide the necessary function, as one cell is wild-type in the chimeras (Fig. 4).

The requirement for *mom-2/wg* in P₂ signaling in genetically mosaic partial embryos was surprising (Fig. 4), as *mom-2/wg* is not required for EMS mitotic spindle orientation after isolating P₁ blastomeres from *mom-2/wg* mutant embryos (Fig. 2). When assembling mosaic partial embryos in culture, however, we did not strictly control the age of P₂ when placed in contact with EMS (see Materials and Methods), as wild-type P₂ blastomeres remain competent to signal throughout their cell cycle (Goldstein 1995b; A. Schlesinger et al., unpubl.). Although the age of a wild-type P₂ does not affect its ability to orient an EMS mitotic spindle, *mom-2/wg* mutant P₂ blastomeres are able to signal only early in the P₂ cell cycle. When assembled into a genetic chimera within the first five minutes of their cell cycle, *mom-2/wg* mutant P₂ blastomeres properly orient the mitotic spindle of a wild-type EMS; however, older P₂ blastomeres fail to do so (Fig. 4B). Therefore, *mom-2/wg* mutant P₂ blastomeres prematurely lose their signaling ability. This defect in signaling from older P₂ blastomeres is specific for *mom-2/wg* mutant embryos and does not occur in genetic mosaic experiments with *mom-1/porc* or *mom-3* mutant P₂ blastomeres (data not shown). We conclude either that our *mom-2/wg* mutant alleles retain some maternal function that rapidly fades with time, or that other signals act in parallel with *mom-2/wg* to polarize EMS before cell division.

P₂ signaling orients the mitotic spindle in EMS independent of gene transcription

The induced rotation of the EMS nucleus and centrosomes involves a polarization of the cytoskeleton. P₂ signaling could polarize the cytoskeleton in EMS either directly, or as a secondary consequence of regulated gene

Figure 4. *mom-1/porc* and *mom-3* are required in P₂ and *mom-5/fz* and *gsk-3* in EMS for spindle orientation. (A) Mutant and wild-type EMS and P₂ blastomeres were isolated and then recombined to form genetic chimeras (Materials and Methods). (Left column) Schematic summaries of EMS spindle angle data from experiments with a mutant P₂ blastomere recombined with a wild-type EMS. For these experiments, EMS spindle ranges were as follows: *mom-1/porc*, 6°–83°; *mom-2/wg*, 0°–90°; *mom-3*, 1°–87°; *mom-4*, 0°–30°; *mom-5/fz*, 1°–13°; *gsk-3*, 3°–39°. (Right column) Data from experiments in which a wild-type P₂ was recombined with a mutant EMS. Spindle angle ranges for these experiments are *mom-1/porc*, 0°–14°; *mom-2/wg*, 1°–12°; *mom-3*, 3°–31°; *mom-4*, 3°–20°; *mom-5/fz*, 4°–87°; *gsk-3*, 1°–85°. (B) Genetic chimeras made with wild-type EMS and *mom-2/wg* mutant P₂. (Right) A 'young' EMS was placed into contact with an 'old' *mom-2/wg* mutant P₂ isolated from an older embryo such that P₂ divided before EMS (range of spindle angles, 12°–90°). (Left) A 'young' EMS was placed into contact with a *mom-2/wg* mutant P₂ of similar age (range of spindle angles: 0°–6°). For definition of young and old blastomeres, see text.

transcription in EMS. Because some components of the Wnt pathway in *Drosophila* are required for the proper orientation of epidermal bristles, and bristle orientation depends on the proper organization of the actin cytoskeleton, it has been long suspected that Wnt signaling might directly target the cytoskeleton (Wong and Adler 1993; see Discussion). Determining if the signal that induces endoderm in *C. elegans* primarily targets the cytoskeleton or the regulation of gene expression is difficult, as endoderm markers appear only after several subsequent developmental steps that clearly involve gene transcription. The EMS mitotic spindle rotation occurs

within minutes of the P₂ signal, however, and we have therefore been able to test if this response to Wnt signaling requires gene transcription in the responding cell.

Using cultured isolated blastomeres, we quantitatively assayed the induced rotation of the EMS centrosome/nucleus complex after blocking transcription using a chemical inhibitor and RNA interference. To provide an internal control for monitoring gene transcription, we used a transgenic *C. elegans* strain that expresses the green fluorescent protein (GFP) under control of the *pes-10* promoter (Fig. 5A). The *pes-10* gene is among the earliest expressed zygotic genes that have been identified in *C. elegans*, with *pes-10* RNA detectable in EMS and its descendants beginning at the four-cell stage (Seydoux and Fire 1994; Seydoux et al. 1996). Culturing blastomeres from *pes-10::GFP* embryos in actinomycin D, a chemical inhibitor of transcription, blocks detectable expression of GFP, as does RNA-mediated gene interference with *ama-1* RNA (see Materials and Methods). The *ama-1* gene encodes the largest subunit of RNA polymerase II (Pol II), and RNAi has been used previously to block its function (Powell-Coffman et al. 1996). In our most rigorous tests, we cultured blastomeres from *ama-1(RNAi)/pes-10::GFP* embryos in actinomycin D. Detectable GFP expression was absent, but in all cases EMS mitotic spindle orientation was normal (Fig. 5B). Previous studies have also suggested that cleavage patterns in early *C. elegans* embryos appear normal after blocking transcription (Edgar et al. 1994; Powell-Coffman et al. 1996). We conclude that components of the Wnt signal transduction pathway control EMS

Figure 5. Transcriptional inhibitors do not block proper EMS spindle orientation. (A) A P₁ blastomere from *pes-10::GFP* strain was isolated and cultured until the equivalent of the 28-cell stage. The fluorescence image was superimposed on a Nomarski micrograph. GFP fluorescence is visible in the descendants of EMS but not in the P₂-derived lineages. (B) Embryos were isolated from *pes-10::GFP* hermaphrodites injected with *ama-1* RNA. P₁ blastomeres from these embryos were cultured in media containing 0.5 µg/ml actinomycin D. EMS was allowed to divide and the EMS spindle angle was recorded. In all cases, although GFP fluorescence was never detected, EMS mitotic spindles oriented properly with respect to the plane of contact with P₂. Correct EMS orientation was also seen in EMS cells derived from *ama-1(RNAi)* P₁ blastomeres cultured in higher concentrations of actinomycin D (10–20 µg/ml, *n* = 6/6; see Materials and Methods).

spindle rotation by directly targeting the cytoskeleton in EMS, not by regulating gene expression.

Discussion

The polarization of endoderm potential in EMS and the orientation of its mitotic spindle both require Wnt pathway components

Our analysis indicates that Wnt signaling from P₂ coordinates the polarization of endoderm potential and orientation of the EMS mitotic spindle during *C. elegans* embryogenesis. Having both outcomes depend on shared genetic components perhaps ensures that differentially localized cytoplasmic determinants are appropriately segregated into the two daughters on cell division. Nevertheless, the two responses in EMS are separable—in some *mom* mutants, proper spindle orientation can occur in the absence of endoderm induction (Fig. 2). Moreover, endoderm can be induced during most of the EMS cell cycle, whereas its spindle can be oriented in response to P₂ signaling only if contact between the two blastomeres is made early in the EMS cell cycle (Goldstein 1995a,b). These differences in the two EMS responses may be due to P₂ expressing more than one signal, or perhaps to differences in the response pathways beyond a *gsk-3* branchpoint (see below). The temporally restricted ability of *mom-2/wg* P₂ blastomeres to properly orient EMS spindles (Fig. 4B) may reflect the presence of separate, functionally redundant signals. An unidentified signal for spindle orientation may be expressed by P₂ early in its cell cycle, masking any requirement for *mom-2/wg*.

C. elegans gsk-3 is a branchpoint for polarizing endoderm potential and orienting the EMS mitotic spindle

We have shown that *gsk-3* and all upstream Wnt pathway components so far identified in our system, with the possible exception of *mom-2/wg*, are required for both endoderm polarization and for mitotic spindle orientation in EMS. In contrast, the identified nematode Wnt pathway components predicted to act downstream of *gsk-3* (*apr-1/APC*, *wrm-1/arm*, and *pop-1/pan*) are required for endoderm induction but not for spindle orientation. We therefore propose a branchpoint at *gsk-3*, to explain how Wnt signaling from P₂ regulates the asymmetric division of EMS (Fig. 6). Pathway components beyond *gsk-3* that act in the spindle orientation branch remain unknown, although several genes with more general roles in spindle orientation during *C. elegans* embryogenesis have been identified (for review, see Kempthues and Strome 1997). Recent studies indicate that Wnt signaling in *Drosophila* also branches within responding cells to regulate both fate specification and tissue polarity (Axelrod et al. 1998; Boutros et al. 1998). This branchpoint occurs at *dishevelled* (*dsh*), however, before *zw3/GSK-3β*. Therefore the induction of multiple

Figure 6. Model for Wnt signaling in the early *C. elegans* embryo. The Wnt pathway in the early *C. elegans* embryo (B) is contrasted to the 'canonical' pathway derived from experimental data from other organisms (A; see Cadigan and Nusse 1997). In this diagram, the signaling cell is drawn above the responding cell. As shown in B, Wnt signaling is required for both EMS mitotic spindle orientation (left) and endoderm specification (right), although there may be separate signaling ligands for these two processes (Rocheleau et al. 1997; Thorpe et al. 1997; A. Schlesinger et al., unpubl.). *gsk-3* may represent a branchpoint in the EMS Wnt pathway, regulating both endoderm induction and mitotic spindle orientation. Not depicted but also possible are arrows from *gsk-3* directly to the EMS spindle, and directly to the cytoskeleton during endoderm induction. In contrast to the roles of *pan* and *LEF-1*, which are converted into activators by Wnt signaling, *pop-1* acts purely as a repressor in the *C. elegans* pathway (Cadigan and Nusse 1997). Release from POP-1-mediated repression of endoderm occurs not in EMS but in its posterior daughter, E.

responses by Wnt signaling may involve branches within responding cells that execute distinct outputs by varying downstream effectors.

gsk-3 acts positively to promote both endoderm specification and proper mitotic spindle orientation

We have shown that the *C. elegans* GSK-3β homolog, *gsk-3*, acts positively during both endoderm specification and mitotic spindle orientation in EMS. This positive role contrasts with the negative roles for GSK-3β during Wnt signal transduction as documented in *Drosophila*, *Xenopus*, and humans (Boyle et al. 1991; Siegfried et al. 1992; He et al. 1995). In these studies of the Wnt pathway, GSK-3β is thought to be active in the absence of Wnt signaling, promoting the degradation of β-catenin in a complex with APC and other proteins (Rubinfeld et al. 1996; Ikeda et al. 1998). In this canonical view, Wnt signaling blocks GSK-3β activity, thereby sta-

bilizing β -catenin, which subsequently can translocate to the nucleus and activate target gene expression by binding to and converting ubiquitous HMG domain repressors into transcriptional activators (Brannon et al. 1997; Fig. 6).

The response to Wnt signaling during endoderm induction in *C. elegans* appears to differ substantially from the canonical model. Instead of converting a repressor into a transcriptional activator, Wnt signaling from P_2 down-regulates the HMG domain protein POP-1 in the posterior daughter of the responding cell, thereby promoting endodermal fate (Rocheleau et al. 1997; Thorpe et al. 1997; Fig. 6). To reconcile these different observations, we suggest that Wnt signaling may induce the coincident degradation of POP-1/PAN and WRM-1/arm in E but not MS. This model invokes the established abilities of β -catenin/arm to form a complex with HMG domain proteins, and of GSK-3 β to promote the degradation of β -catenin. How Wnt signaling could be adapted to activate, instead of block, GSK-3 β function in this pathway remains entirely unclear, as does the ability of GSK-3 to positively promote mitotic spindle rotation, which occurs independently of *pop-1/pan* function. Our understanding of these issues may improve once the subcellular localizations of different Wnt pathway components are determined, and more interacting factors are identified.

Because all Wnt pathway components are required positively for endoderm induction in EMS, it has not been possible to order the function of different gene products relative to each other using double mutant analyses. Nevertheless, other data suggest that components of the Wnt pathway function in the same relative order during endoderm induction as they do in the canonical pathway. First, the excess endoderm phenotype of *pop-1/pan* mutants is epistatic to the *mom-2/wg*, *mom-5/fz*, *wrm-1/arm*, *gsk-3*, and *mom-4* mutant phenotypes, indicating that *pop-1/pan* acts downstream of these genes (Rocheleau et al. 1997; Thorpe et al. 1997; data not shown). Second, our analysis of genetically mosaic partial embryos shows that *mom-5/fz* and *gsk-3* are both required in EMS, the responding cell, and are therefore downstream of *mom-1/porc* and *mom-2/wg*, genes that are required in the signaling cell, P_2 . Third, we note that all Wnt pathway components up to and including *gsk-3*, except possibly *mom-2/wg*, are required both to polarize endoderm potential and to orient the EMS mitotic spindle, whereas all Wnt pathway components downstream of *gsk-3* are required only for endoderm induction. This latter observation is consistent with *gsk-3* functioning upstream of *wrm-1/arm* and *apr-1/APC*. Given these similarities to the canonical Wnt pathway, it seems likely that the relative order in which Wnt pathway components function is conserved during endoderm induction in *C. elegans* in spite of their different regulatory relationships.

Finally, our finding that *gsk-3* acts positively during endoderm induction is not without precedent for a GSK-3 β family member. A GSK-3 β homolog in *Saccharomyces cerevisiae*, Rim1p, positively regulates the expres-

sion of meiotic genes during sporulation by complexing with and phosphorylating two transcription factors, Ume6p and Ime1p (Mandel et al. 1994; Strich et al. 1994; Rubin-Bejerano et al. 1996; Malathi et al. 1997). Furthermore, we note that *gsk-3* in *C. elegans* appears to be required both positively and negatively to regulate the specification of endoderm in the early embryo. Whereas E fails to produce endoderm in fewer than half of *gsk-3* mutant embryos, a daughter of P_2 called C produces endoderm ectopically in approximately three-fourths of the mutant embryos (Table 1). Therefore, *gsk-3* is required to negatively regulate endoderm specification in P_2 , the sister of EMS. These results suggest that *gsk-3* function is capable of generating opposite outputs even within a single organism, perhaps through the intervention of other regulatory pathways.

The cytoskeleton as the primary target of cell signaling

Because rotational positioning of the EMS centrosomes appears to not require gene transcription, we suggest that P_2 to EMS signaling directly targets the cytoskeleton in EMS. Microtubules are the primary structural component of the mitotic spindle and are thought to interact with cortical microfilaments during rotation of the nucleus/centrosome complex in EMS (Hyman 1989; Goldstein et al. 1993; Waddle et al. 1994). Either microtubules or microfilaments could be targets of Wnt signaling, as both are required for correct orientation of the mitotic spindle in EMS (Hyman and White 1987; Goldstein 1995b). Wnt pathway genes have been implicated in cytoskeleton-related processes in other systems. Mammalian GSK-3 β can phosphorylate the microtubule-associated protein tau, and this activity is regulated by *dsh*, another Wnt pathway component (Wagner et al. 1997). Therefore P_2 signaling might influence microtubule dynamics directly through *gsk-3*. Alternatively, P_2 signaling might influence spindle orientation in EMS by locally activating microtubule motor complexes that directly interact with astral microtubules to orient the nucleus/centrosome complex before cell division.

It seems likely that Wnt signaling directly targets the cytoskeleton to control other developmental processes. For example, the polarized organization of hair cells in the *Drosophila* epithelium appears to be controlled by the orientation of actin microfilaments, which in turn are regulated by tissue polarity genes including *fz* and *dsh* (Wong and Adler 1993; Axelrod et al. 1998; Boutros et al. 1998). Wnt signaling in *C. elegans* may also target the cytoskeleton to polarize endoderm potential in EMS, as the polarization of endoderm potential appears to require functional microfilaments and microtubules (Goldstein 1995a). Alternatively, P_2 signaling may simply cause a localized activation of downstream Wnt pathway components, resulting in an asymmetric segregation of their activity to the daughters of EMS.

In conclusion, we emphasize that some developmental signals may influence cell fate by directly targeting cytoplasmic components in a responding cell and only in-

directly regulate gene transcription. As the Wnt pathway is widely conserved, it will be interesting to learn if direct targeting of the cytoskeleton in cells responding to Wnt signaling proves to be of general importance during animal development.

Materials and methods

Strains and alleles

Basic *C. elegans* culturing methods were used as described in Brenner (1974). N2 Bristol was used for the wild-type strain. The *mom* and *pop-1* alleles are recessive, maternal-effect lethal mutations. The mutant strains used were EU340 *pop-1(zu189) dpy-5(e61) I/hT1 (I;V)*; *him-5(e1490) IV*, EU308 *mom-1(or10) dpy-6(e14) X/szT1 (1;X)*, EU98 *mom-1(or10) lin-2(e1309) X/szT1 (1;X)*, EU278 *mom-2(or42) V/DnT1 (IV;V)*, EU95 *mom-2(or9) V/DnT1 (IV;V)*, EU365 *mom-2(or85) V/DnT1 (IV;V)*, EU376 *rol-6(e187) mom-3(or78)/mnC1 II*; *lin-2(e1309) X*, EU431 *rol-1(e91) mom-3(or78) lin-7(e1413)/mnC1 II*; *him-5(e1490) V*, EU309 *mom-4(or39) dpy-5(e61) I/hT2 (I;III)*, and EU452 *unc-13(e1091) mom-5(zu193) I/hT2 (I;III)*, EU340 *pop-1(zu189) dpy-5(e61) I/hT1 (I;V)*; *him-5(e1490) IV*. The JH103 transgenic strain containing the *pes-10::GFP* construct was provided by G. Seydoux (Seydoux and Fire 1994).

Microscopy and laser ablations

Light microscopy was performed with a Zeiss Axioscope equipped for Nomarski optics and epifluorescence. Data were either photographed with TechPan 200 film and developed with HC110, or recorded using a Dage-MTI Ve1000 digital camera linked to the Scion Image program run on a Macintosh PowerPC 8100/110. All images were processed using Adobe Photoshop 4.0.

Laser ablations were performed as described previously (Avery and Horvitz 1989). Embryos were fixed and stained with monoclonal antibodies 3NB12 and 5.6 to detect pharyngeal muscle and body wall muscle cells, respectively (Miller et al. 1983; Priess and Thomson 1987; Bowerman et al. 1992).

Blastomere isolation and spindle angle measurements

Early embryos were obtained by cutting open adult hermaphrodites and blastomeres were isolated and cultured as described previously (Edgar 1995; Shelton and Bowerman 1996). The blastomeres were mechanically separated by repeated mouth pipetting through a glass micropipette. For P_1 isolations, sister cells AB and P_1 were separated and P_1 was cultured alone. The orientation of the EMS mitotic spindle was determined as illustrated in Figure 1D—axes were designated (black lines) with the plane of contact between EMS and P_2 as 90° and the spindle angle (gray line) was measured relative to these axes. The angle of the EMS spindle was measured when the EMS cytokinetic furrow began to form. At this point in the cell cycle, the mitotic spindle has begun to elongate and the nuclear membrane has broken down. As in intact embryos, the EMS cell cycle was ~15 min long and P_2 divided ~4 min after EMS. For genetically mosaic partial embryos, wild-type and mutant early EMS and P_2 blastomeres were concurrently isolated. EMS and P_2 were recombined early in the EMS cell cycle allowing the cells to be in contact for at least 9 min before EMS cytokinesis. For all figures, data from multiple trials were schematically condensed into a single quadrant. Each line on the periphery of the quadrant, representing a data point, is placed at the angle of EMS division.

Identification of *C. elegans gsk-3*

Degenerate primers were made to regions conserved in known *gsk-3* genes but not conserved in other, related Ser/Thr kinases. cDNA was obtained from N2 worms by standard techniques using Trizol reagent (GIBCO-BRL) for RNA isolation and SuperScript II (GIBCO-BRL) for first-strand synthesis. PCR was performed with a MJ-Research PTC-100 thermocycler using *Taq* polymerase (Perkin Elmer), 1.5 mM $MgCl_2$, and an annealing temperature of $50^\circ C$. A single 900-bp band was obtained, cloned into pCRII (Invitrogen), and sequenced.

Sequence data were used to design primers for cloning additional cDNA fragments. For all PCR reactions described below, the high-fidelity Pwo polymerase (Boehringer Mannheim) was used. The 5' end of the cDNA was obtained by PCR with a primer to the *trans*-spliced leader sequence SL1 and internal primer ZW3C. The resulting 556-bp band was cloned into pGem-T (Promega) to yield pJMZWd. A fragment corresponding to bp 461–1227 was obtained by using primers ZW5E and ZW3F and cloned into pGem-T to yield pJMZWf. For sequencing, isolates from at least two independent PCR reactions were obtained, and each isolate was sequenced on both strands. The resulting sequence was used to search a *C. elegans* EST database, which revealed three cDNA clones corresponding to the *gsk-3* gene—yk307d8, yk387f2, and yk451g9. These clones were obtained and analyzed to determine the remaining 3' sequence.

RNA-mediated gene interference (RNAi)

gsk3 RNA was made from plasmid zw3/F1, which contains ~800 bp of the 5' coding region in pGEM-T (Promega). The insert was PCR-amplified with SP6 and T7 primers. The PCR product was used to make SP6 and T7 primed RNA, which was mixed to yield double-stranded RNA. *apr-1/APC* and *wrm-1/arm* RNAs were generated from cDNA phage clones using T7 and T3 primers for reverse transcription reactions (*apr-1/APC* was made from phage clone yk40c12 and *wrm-1/arm* from yk21a9). *ama-1* RNA was made with T7 primers from plasmid DB19, which contains ~4.7 kb of the *ama-1* coding region in pGEM-7ZF.

RNA was injected at concentrations between 1 and 5 $\mu g/ml$ into wild-type hermaphrodites, except for *ama-1* RNA, which was injected into the transgenic *pes-10::GFP* strain. After 24 hr, injected hermaphrodites were singled out and mated with wild-type males [except for *mom-2(RNAi)*] to increase the number of progeny. Twelve to 24 hr later, hermaphrodites laying embryos with the appropriate phenotype were used to isolate early embryos.

Transcriptional inhibition

In embryos from the JH103 strain, GFP fluorescence, driven by the *pes-10* promoter, is detectable at the 15-cell stage (after the daughters of EMS have divided), when it comes on in most somatic blastomeres (G. Seydoux, pers. comm.; data not shown). *pes-10* expression is detectable at the four-cell stage by RNA in-situ hybridization (Seydoux and Fire 1994; Seydoux et al. 1996). The lag in GFP detection is caused by a delay in translation and the long oxidation step required for GFP fluorescence (G. Seydoux, pers. comm.; Reid and Flynn 1997). GFP fluorescence was checked at the 15-cell stage as well as 24 hr later. The lack of GFP fluorescence was taken as an indication that inhibition of transcription occurred before transcription of the *pes-10::GFP* construct would have begun at the four-cell stage.

Transcriptional inhibition was achieved using two methods. First, RNA made from *ama-1*, the gene encoding the largest

subunit of RNA Pol II was used in RNA-mediated gene interference (Bird and Riddle 1989). *ama-1*(RNAi) embryos die with <100 cells (Powell-Coffman et al. 1996). RNAi was performed on *pes-10::GFP* hermaphrodites as described above and P₁ blastomeres were isolated from embryos collected 36–48 hr after injection. In addition, P₁ blastomeres were cultured in media supplemented with 0.5 µg/ml actinomycin D (Sigma, from an ~10 mg/ml stock in EtOH). These partial embryos arrested as a group of 12 cells, which is the equivalent of the 28-cell stage. With either method alone, GFP expression was not detected in blastomeres and EMS mitotic spindles oriented within the wild-type range (10/10 for *ama-1*(RNAi) and 10/10 for 0.5 µg/ml actinomycin D). For the experiments shown in Figure 5B, P₁ blastomeres from *ama-1*(RNAi) embryos were isolated and allowed to divide in culture media containing ~0.5 µg/ml actinomycin D. The division angle of the EMS spindle was measured as indicated above. Correct EMS spindle orientation was also seen in experiments done using *ama-1*(RNAi) blastomeres cultured in higher concentrations of actinomycin D [10–20 µg/ml (*n* = 6)]. At the higher concentrations, however, blastomeres sometimes failed to divide even though the EMS spindle oriented normally, apparently attributable to toxic levels of the ethanol solvent [data not shown].

Acknowledgments

We thank Geraldine Seydoux for the *pes-10::GFP* strain; Jennifer Knight and Bill Wood for the *ama-1* plasmid; Craig Mello and Kevin Fitzgerald for sharing *gsk-3* sequence information; and David Miller and Julie Ahringer for antibodies. We are grateful to Yuji Kohara for providing cDNA clones, to the *C. elegans* Genetic Stock Center [funded by the National Institutes of Health (NIH) National Center for Research Resources] for sending strains, to Matt Thomas and J. Clayton Carter for technical help, and to Chris Doe and Danielle Hamill for helpful advice and comments on the manuscript. We especially thank Cynthia Kenyon for her generosity and support of J.N.M. A.S. and M.M. were supported by training grants from the NIH (GM07413 and HD07348); C.A.S. by a postdoctoral fellowship from the NIH (GM16981); J.N.M. in part by a Howard Hughes Medical Institute predoctoral fellowship, and B.B. by a grant from the NIH (GM49869).

The publication costs of this article were defrayed in part by payment of page charges. This article must therefore be hereby marked 'advertisement' in accordance with 18 USC section 1734 solely to indicate this fact.

Note added in proof

The GenBank accession no. for *gsk-3* is AF159950. A deletion allele of this gene has been isolated by Michael Basson at Axys Pharmaceuticals, Inc. In future publications the gene *sgg-1* will replace *gsk-3*.

References

- Avery, L. and H.R. Horvitz. 1989. Pharyngeal pumping continues after laser killing of the pharyngeal nervous system of *C. elegans*. *Neuron* **3**: 473–485.
- Axelrod, J.D., J.R. Miller, J.M. Shulman, R.T. Moon, and N. Perrimon. 1998. Differential recruitment of Dishevelled provides signaling specificity in the planar cell polarity and wingless signaling pathways. *Genes & Dev.* **12**: 2610–2622.
- Bird, D.M. and D.L. Riddle. 1989. Molecular cloning and sequencing of *ama-1*, the gene encoding the largest subunit of *Caenorhabditis elegans* RNA polymerase II. *Mol. Cell. Biol.* **9**: 4119–4130.
- Boutros, M., N. Paricio, D.I. Strutt, and M. Mlodzik. 1998. Dishevelled activates JNK and discriminates between JNK pathways in planar polarity and wingless signaling. *Cell* **94**: 109–118.
- Bowerman, B., B.A. Eaton, and J.R. Priess. 1992. *skn-1*, a maternally expressed gene required to specify the fate of ventral blastomeres in the early *C. elegans* embryo. *Cell* **68**: 1061–1075.
- Boyle, W.J., T. Smeal, L.H. Defize, P. Angel, J.R. Woodgett, M. Karin, and T. Hunter. 1991. Activation of protein kinase C decreases phosphorylation of c-Jun at sites that negatively regulate its DNA-binding activity. *Cell* **64**: 573–584.
- Brannon, M., M. Gomperts, L. Sumoy, R.T. Moon, and D. Kimelman. 1997. A beta-catenin/XTcf-3 complex binds to the *siamois* promoter to regulate dorsal axis specification in *Xenopus*. *Genes & Dev.* **11**: 2359–2370.
- Brenner, S. 1974. The genetics of *Caenorhabditis elegans*. *Genetics* **77**: 71–94.
- Cadigan, K.M. and R. Nusse. 1997. Wnt signaling: A common theme in animal development. *Genes & Dev.* **11**: 3286–3305.
- Edgar, L.G. 1995. Blastomere culture and analysis. In *Caenorhabditis elegans: Modern biological analysis of an organism* (ed. H.F. Epstein and D.C. Shakes), pp. 303–321. Academic Press, San Diego, CA.
- Edgar, L.G., N. Wolf, and W.B. Wood. 1994. Early transcription in *Caenorhabditis elegans* embryos. *Development* **120**: 443–451.
- Fire, A., S. Xu, M.K. Montgomery, S.A. Kostas, S.E. Driver, and C.C. Mello. 1998. Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. *Nature* **391**: 806–811.
- Goldstein, B. 1992. Induction of gut in *Caenorhabditis elegans* embryos. *Nature* **357**: 255–257.
- . 1993. Establishment of gut fate in the E lineage of *C. elegans*: The roles of lineage-dependent mechanisms and cell interactions. *Development* **118**: 1267–1277.
- . 1995a. An analysis of the response to gut induction in the *C. elegans* embryo. *Development* **121**: 1227–1236.
- . 1995b. Cell contacts orient some cell division axes in the *Caenorhabditis elegans* embryo. *J. Cell Biol.* **129**: 1071–1080.
- Goldstein, B., S.N. Hird, and J.G. White. 1993. Cell polarity in early *C. elegans* development. *Development Suppl.* 279–287.
- Hawkins, N. and G. Garriga. 1999. Asymmetric cell division: From A to Z. *Genes & Dev.* **12**: 3625–3638.
- He, X., J.-P. Saint-Jeannet, J.R. Woodgett, H.E. Varmus, and I.B. Dawid. 1995. Glycogen synthase kinase-3 and dorsoventral patterning in *Xenopus* embryos. *Nature* **374**: 617–622.
- Hyman, A.A. 1989. Centrosome movement in the early divisions of *Caenorhabditis elegans*: A cortical site determining centrosome position. *J. Cell Biol.* **109**: 1185–1193.
- Hyman, A.A. and J.G. White. 1987. Determination of cell division axes in the early embryogenesis of *Caenorhabditis elegans*. *J. Cell Biol.* **105**: 2123–2135.
- Ikeda, S., S. Kishida, H. Yamamoto, H. Murai, S. Koyama, and A. Kikuchi. 1998. Axin, a negative regulator of the Wnt signaling pathway, forms a complex with GSK-3β and β-catenin and promotes GSK-3β-dependent phosphorylation of β-catenin. *EMBO J.* **17**: 1371–1384.
- Kemphues, K. and S. Strome. 1997. Fertilization and establishment of polarity in the embryo. In *Caenorhabditis elegans II* (ed. D.L. Riddle, T. Blumenthal, B.J. Meyer and J.R. Priess),

- pp. 335–360. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Lin, R., S. Thompson, and J.R. Priess. 1995. *pop-1* encodes an HMG box protein required for the specification of a mesoderm precursor in early *C. elegans* embryos. *Cell* **83**: 599–609.
- Lin, R., R.J. Hill, and J.R. Priess. 1998. POP-1 and anterior-posterior fate decisions in *C. elegans* embryos. *Cell* **92**: 229–239.
- Lu, B., L.Y. Jan, and Y. N. Jan. 1998. Asymmetric cell division: Lessons from flies and worms. *Curr. Opin. Genet. Dev.* **8**: 392–399.
- Malathi, K., Y. Xiao, and A.P. Mitchell. 1997. Interaction of yeast repressor-activator protein Ume6p with glycogen synthase kinase 3 homolog Rim1p. *Mol. Cell Biol.* **17**: 7230–7236.
- Mandel, S., K. Robzyk, and Y. Kassir. 1994. *IME1* gene encodes a transcription factor which is required to induce meiosis in *Saccharomyces cerevisiae*. *Dev. Genet.* **15**: 139–147.
- Miller, D.M., III, I. Ortiz, G.C. Berliner, and H.F. Epstein. 1983. Differential localization of two myosins within nematode thick filaments. *Cell* **34**: 477–490.
- Miller, J.R. and R.T. Moon. 1997. Analysis of the signaling activities of localization mutants of β -catenin during axis specification in *Xenopus*. *J. Cell Biol.* **139**: 229–243.
- Powell-Coffman, J.A., J. Knight, and W.B. Wood. 1996. Onset of *C. elegans* gastrulation is blocked by inhibition of embryonic transcription with an RNA polymerase antisense RNA. *Dev. Biol.* **178**: 472–483.
- Priess, J.R. and J.N. Thomson. 1987. Cellular interactions in early *C. elegans* embryos. *Cell* **48**: 241–250.
- Reid, B.G. and G.C. Flynn. 1997. Chromophore formation in green fluorescent protein. *Biochemistry* **36**: 6786–6791.
- Rocheleau, C.E., W.D. Downs, R. Lin, C. Wittmann, Y. Bei, Y.H. Cha, M. Ali, J.R. Priess, and C.C. Mello. 1997. Wnt signaling and an APC-related gene specify endoderm in early *C. elegans* embryos. *Cell* **90**: 707–716.
- Rubin-Bejerano, I., S. Mandel, K. Robzyk, and Y. Kassir. 1996. Induction of meiosis in *Saccharomyces cerevisiae* depends on conversion of the transcriptional repressor Ume6 to a positive regulator by its regulated association with the transcriptional activator Ime1. *Mol. Cell Biol.* **16**: 2518–2526.
- Rubinfeld, B., I. Albert, E. Porfiri, C. Fiol, S. Munemitsu, and P. Polakis. 1996. Binding of GSK3 β to the APC- β -catenin complex and regulation of complex assembly. *Science* **272**: 1023–1026.
- Schierenberg, E. 1987. Reversal of cellular polarity and early cell-cell interactions in the embryo of *Caenorhabditis elegans*. *Dev. Biol.* **122**: 452–463.
- Seydoux, G. and A. Fire. 1994. Soma-germline asymmetry in the distributions of embryonic RNAs in *Caenorhabditis elegans*. *Development* **120**: 2823–2834.
- Seydoux, G., C.C. Mello, J. Pettitt, W.B. Wood, J.R. Priess, and A. Fire. 1996. Repression of gene expression in the embryonic germ lineage of *C. elegans*. *Nature* **382**: 713–716.
- Shelton, C.A. and B. Bowerman. 1996. Time-dependent responses to *glp-1*-mediated inductions in early *C. elegans* embryos. *Development* **122**: 2043–2050.
- Siegfried, E., T.-B. Chou, and N. Perrimon. 1992. *wingless* signaling acts through *zeste-white 3*, the *Drosophila* homolog of *glycogen synthase kinase-3*, to regulate *engrailed* and establish cell fate. *Cell* **71**: 1167–1179.
- Strich, R., R.T. Surosky, C. Steber, E. Dubois, F. Messenguy, and R.E. Esposito. 1994. *UME6* is a key regulator of nitrogen repression and meiotic development. *Genes & Dev.* **8**: 796–810.
- Sulston, J.E., E. Schierenberg, J.G. White, and J.N. Thomson. 1983. The embryonic cell lineage of the nematode *Caenorhabditis elegans*. *Dev. Biol.* **100**: 64–110.
- Thorpe, C.J., A. Schlesinger, J.C. Carter, and B. Bowerman. 1997. Wnt signaling polarizes an early *C. elegans* blastomere to distinguish endoderm from mesoderm. *Cell* **90**: 695–705.
- Waddle, J.A., J.A. Cooper, and R.H. Waterston. 1994. Transient localized accumulation of actin in *Caenorhabditis elegans* blastomeres with oriented asymmetric divisions. *Development* **120**: 2317–2328.
- Wagner, U., J. Brownlee, N.G. Irving, F.R. Lucas, P.C. Salinas, and C.C.J. Miller. 1997. Overexpression of the mouse dishevelled-1 protein inhibits GSK-3 β -mediated phosphorylation of tau in transfected mammalian cells. *FEBS Lett.* **411**: 369–372.
- Wong, L.L. and P.N. Adler. 1993. Tissue polarity genes of *Drosophila* regulate the subcellular location for prehair initiation in pupal wing cells. *J. Cell Biol.* **123**: 209–221.

Wnt pathway components orient a mitotic spindle in the early *Caenorhabditis elegans* embryo without requiring gene transcription in the responding cell

Ann Schlesinger, Christopher A. Shelton, Julin N. Maloof, et al.

Genes Dev. 1999, **13**:

References

This article cites 44 articles, 22 of which can be accessed free at:
<http://genesdev.cshlp.org/content/13/15/2028.full.html#ref-list-1>

License

Email Alerting Service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).