An HMG1-like protein facilitates Wnt signaling in *Caenorhabditis elegans*

Lily I. Jiang and Paul W. Sternberg¹

Howard Hughes Medical Institute and Division of Biology, California Institute of Technology,
Pasadena, California 91125 USA

We show that during *Caenorhabditis elegans* male spicule development, the specification of a glial versus neuronal cell fate in a canonical neurogenic sublineage is dependent on Wnt signaling. Inactivation of a Wnt signaling pathway mediated by the Wnt receptor LIN-17 transforms the SPD sheath cell into its sister, the SPD neuron. We discovered a new mutant, *son-1*, that displays this same cell fate transformation. The *son-1* mutation enhances the phenotypes of reduction-of-function *lin-17* mutants in several developmental processes, including vulva development, somatic gonad development, and male tail patterning. *son-1* encodes an HMG1/2-like DNA-binding protein and is localized in all cell nuclei through development as revealed by a GFP reporter construct. Disruption of *son-1* function by RNA-mediated interference results in the same spicule defect as caused by overexpression of POP-1, a TCF/LEF class HMG protein known to act downstream of the Wnt signaling pathway. Our results provide *in vivo* evidence for the functional involvement of an HMG1/2-like protein, SON-1, in Wnt signaling. The sequence nonspecific HMG protein SON-1 and the sequence specific HMG protein POP-1 might both act in the Wnt responding cells to regulate gene transcription in opposite directions.

[Key Words: Wnt; HMG-1; TCF/LEF; cell fate specification; *C. elegans*]

Received December 28, 1998; revised version accepted February 19, 1999.

The Wnt signaling pathway is one of the major signaling pathways controlling animal development. It controls cell proliferation, cell and tissue polarity, and cell fate specification in many developmental processes of many organisms. In recent years, several components of the Wnt signaling pathway have been identified and the mechanisms by which these factors act have begun to be elucidated (for review, see Cadigan and Nusse 1997; Moon et al. 1997). Wnt family genes, which encode secreted glycoproteins, act as intercellular signals. The Frizzled (Fz) family seven transmembrane proteins are putative Wnt receptors that transduce Wnt signals to downstream targets. Activation of the Wnt signaling pathway leads to accumulation of β -catenin in the cytoplasm. β -Catenin interacts with the TCF/LEF family HMG box-containing DNA-binding protein and the complex accumulates in the nucleus, whereby it regulates gene expression of Wnt pathway targets. Although this simplified model appears to be conserved among diverse organisms, the discoveries of new components and the multiplicity of Wnt pathway components have added complexity to this model.

In *Caenorhabditis elegans*, there are several Wnt signaling pathways that are involved in several aspects of development such as cell fate specification during em-

bryogenesis, vulval formation, somatic gonad development, and tail patterning. Some components are shared by several or all of these processes. During embryogenesis, Wnt signaling pathway genes, *mom-2* (Wnt), *mom-5* (Fz), *wrm-1* (β -catenin), and *pop-1* (TCF/LEF), mediate interactions between the P2 and EMS cells (Lin et al. 1995, 1998; Rocheleau et al. 1997; Thorpe et al. 1997). Signaling from the P2 cell promotes asymmetric division of the EMS cell to produce an anterior MS fate and a posterior E fate. Inactivation of the Wnt signaling pathway results in an E-to-MS fate transformation. The Wnt pathway also plays a role in vulval development. Mutants of several Wnt pathway components show various vulva lineage defects, for example, *lin-17* (Fz), *bar-1* (β -catenin), and *mom-1* and *mom-3* mutants (Sternberg and Horvitz 1988; Sawa et al. 1996; Thorpe et al. 1997; Eisenmann et al. 1998). In somatic gonad development, *lin-17* has been also shown to control several asymmetric cell divisions (Sternberg and Horvitz 1988). In the tail region, a Wnt signal LIN-44 and its putative receptor LIN-17 control the polarity of several asymmetric cell divisions, including B, T, F, and U (Herman and Horvitz 1994; Chamberlin and Sternberg 1995; Herman et al. 1995; Sawa et al. 1996). *lin-44*, *lin-17*, and *bar-1* also are involved in the specification of P12 fate in the tail region (Eisenmann et al. 1998; Jiang and Sternberg 1998). A common feature of most of these Wnt signaling pathways in *C. elegans* appears to be that Wnt signaling is

¹Corresponding author.

E-MAIL pws@cco.caltech.edu; FAX (626) 568-8012.

required for posterior cell fates, and inactivation of signaling results in posterior to anterior fates transformation.

The TCF/LEF class HMG box proteins have been shown to regulate Wnt target gene expression on pathway activation (van de Wetering et al. 1997). The *C. elegans* TCF/LEF class protein is encoded by the *pop-1* gene (Lin et al. 1995). *pop-1* negatively regulates Wnt signaling, as loss-of-function mutation in *pop-1* confers a phenotype opposite that of inactivation of Wnt signaling. Moreover, the expression level of POP-1 protein appears to be up-regulated on inactivation of Wnt signaling (Rocheleau et al. 1997; Lin et al. 1998). Therefore, it was proposed that activation of the Wnt signaling pathway relieves the inhibitory effects of POP-1 on gene transcriptional regulation. Yet it is not clear how Wnt target gene expression is regulated and what other unidentified factors are involved in this process.

We have been studying cell-cell interactions and cell fate specification during *C. elegans* male spicule development. Previous studies have shown that cell fate specification in spicule development is mediated by multiple signaling pathways. The *C. elegans* male spicule is generated by a single male specific B blast cell (Fig. 1). The first division of B cell is asymmetric and is controlled by the Wnt signaling pathway genes, *lin-44* and *lin-17* (Sternberg and Horvitz 1988; Herman and Horvitz 1994; Chamberlin and Sternberg 1995). At the early third larval stage (L3; B lineage 10-cell stage), the specification of B cell progeny involves a RAS signaling pathway and a *C. elegans* Notch homolog *lin-12* (Chamberlin and Sternberg 1994). During the L3 stage, those cells continue to divide to generate spicule neurons, non-neuronal sheath and socket cells, and connective tissues (Fig. 1). For example, ζ sublineage generates the neuron SPD and its associated sheath (SPDsh) and socket (SPDso) cells.

Figure 1. The B lineage. The B cell division pattern from 1- to 10-cell stage and the neurogenic sublineages β and ζ are shown (adapted from Sulston et al. 1980; Chamberlin and Sternberg 1993). (sh) Sheath cell; (so) socket cell. The SPV neuron is derived from β sublineage and the SPD neuron from ζ sublineage. The SPD neuron and SPD sheath cell are sister cells derived from ζ .pa.

Using a molecular marker that is specifically expressed in ζ -derived SPD neurons, we studied neuronal versus non-neuronal cell fate specification during male spicule development. Although the cell division pattern of the ζ sublineage resembles the *Drosophila* sensory organ precursor (SOP) lineage in which the Notch pathway plays a crucial role, we found that *lin-12* is not involved in the specification of the SPD neuron fate. Rather, the *lin-17*-mediated Wnt signaling pathway plays an important role in this neuronal versus non-neuronal fate decision in ζ lineage. Loss-of-function mutants of *lin-17* display a SPDsh to neuron fate transformation. To identify new genes involved in this cell fate decision, we conducted a genetic screen and discovered a mutant, *son-1*(sy549), which shows the same cell fate transformation. Moreover the *son-1* (sheath-to-neuron fate transformation) mutation enhances the phenotypes of a *lin-17* hypomorph, *n698*, to *lin-17* null-like, with respect to vulval development, somatic gonad development, and male tail patterning. We cloned the *son-1* locus and found it encodes an HMG1/2-like DNA-binding protein, which is distinct from the TCF/LEF class HMG proteins. Disruption of *son-1* function through RNA-mediated interference (RNAi) leads to defects in several Wnt pathway-mediated developmental processes. Overexpression of POP-1 causes the same spicule defect as loss-of-function *son-1*. Our results provide in vivo evidence that the HMG1/2-like protein encoded by *son-1* plays a specific role in Wnt signaling. SON-1 and POP-1 may both act in the Wnt-responding cells, but in opposite directions, to regulate gene transcription. Taken together, these data provide evidence for further complexity in Wnt responses during development.

Results

lin-17 mutants cause a SPDsh to neuron fate transformation

To study terminal cell fate specification in *C. elegans* male spicule development, we used a molecular marker, *gpa-1::lacZ*. In transgenic animals bearing a *lacZ* reporter gene driven by the *gpa-1* promoter, expression is found in male spicules as well as phasmid neurons in the tail region and several cells in the head (J. Mendel and P.W. Sternberg, unpubl.). In wild-type animals, *gpa-1::lacZ* marker is expressed in one cell per spicule. Cell ablation experiments identified this cell as the spicule neuron SPD (Fig. 2A).

Using this SPD neuron specific marker, we examined spicule terminal cell fates in several classes of known mutants. The Notch pathway controls neuronal versus non-neuronal cell fate specification in both the central and peripheral nervous systems during *Drosophila* embryonic development (Hartenstein and Posakony 1990; Parks and Muskavitch 1993). In the development of *Drosophila* SOP lineage, the Notch pathway mediates interactions between two daughter cells to ensure one adopt neuronal fate, the other one non-neuronal fate. As the cell division pattern of the spicule neurogenic lineage

resembles that of *Drosophila* SOP lineage (Fig. 1), we examined the possible involvement of the *C. elegans* Notch homolog *lin-12* in the specification of SPD neuron versus SPDsh fate. Neither gain-of-function nor loss-of-function *lin-12* mutants displayed abnormal marker expression pattern ($n = 50$). Activation of the *lin-12* pathway at a critical developmental time by overexpressing the cytoplasmic ankyrin repeat region of GLP-1, which has been shown to act as an activated *lin-12* or *glp-1* in many developmental processes (Roehl and Kimble 1993), did not affect marker expression ($n = 50$).

The Wnt pathway genes, *lin-44* and *lin-17*, are involved in the first asymmetric division of the B cell. Mutant males bearing strong *lin-44* and *lin-17* alleles have early B lineage defects, resulting in severely crumpled spicules. From a few escapers in which the first division of B cell is not disrupted, we noticed extra expression of the *gpa-1::lacZ* marker in the spicule region (Fig. 2B,C). Using a weak *lin-17* allele, *n698*, wherein the first B cell division is usually not disrupted, we found that 80% of the animals show two cells per spicule expressing the SPD marker (Fig. 2D; Table 1).

To determine which cell is responsible for the ectopic expression, we ablated specific cells in the B lineage and examined marker expression in operated animals. Because the ϵ lineage develops abnormally in *lin-17* mutants (Chamberlin and Sternberg 1995; this paper), we

Table 1. Both *son-1(sy549)* and *lin-17(n698)* confer a SPD sheath to neuron fate transformation

Cells ablated	<i>lin-17(n698)</i>		<i>son-1(sy549)</i>	
	no. ^a	marker expression ^b	no. ^a	marker expression ^b
Intact	15/17	2	39/52	2
β	5/6	2	4/6	2
ζ	15/16	0	12/15	0
ζ a	13/15	2	5/5	2
ζ p	5/5	0	6/9	0
ζ pp	4/7	2	1/2	2
ζ pa				
(SPD+SPDsh)	11/11	0	12/14	0
ζ paa (SPD)	10/13	1	13/16	1
ζ pap (SPDsh)	13/21	1	6/7	1

Cells were ablated at L3 stage and animals recovered to allow reaching adulthood. Adult males were then examined for *gpa-1::lacZ* marker expression. There are two ζ cells per male; both were ablated.

^aNumbers of animals operated on and scored that showed the corresponding phenotypes. The rest of the animals showed 0, 1, or 2 cells per spicule expressing the marker due to incomplete penetrance or ablation error or damage.

^bNumbers indicate number of cells per spicule expressing the *gpa-1::lacZ* marker.

Figure 2. (A) In wild-type males, the *gpa-1::lacZ* marker is expressed in one cell per spicule. Cell ablation experiment identifies this cell being SPD neuron. Both spicules are shown on this focal plane. (B–D) *gpa-1::lacZ* is expressed in two cells per spicule in *lin-44* and *lin-17* mutant males. (B) *lin-44(n1792)*; (C) *lin-17(n671)*; (D) *lin-17(n698)*, early B lineage defect is rare in this weak allele and ~80% males show a two cells per spicule marker expression pattern. Lateral view, only one spicule is in focus. (E–H) The *gpa-1::lacZ* marker is expressed in two cells per spicule in *son-1(sy549)* mutants. Cell ablation experiment indicated that *son-1(sy549)* confers a SPD sheath to neuron fate transformation. (E) *sy549* intact animal, lateral view showing the left spicule. (F) When the precursor of the SPD neuron and the SPD sheath is ablated, no expression is observed in the spicule region. (G) When the SPD neuron is ablated, the SPD marker is expressed in one cell per spicule. (H) When the SPD sheath is ablated, the SPD marker is expressed in one cell per spicule. Scale bar, 20 μ m.

first ablated the ϵ cells at the 10-cell stage and examined the marker expression pattern in operated animals. Ablation of ϵ does not eliminate the extra *gpa-1::lacZ* marker-expressing cell ($n = 9$). Further ablation experiments indicate that the extra expression results from SPDsh, the sister cell of the SPD neuron (Table 1). Ablation of the precursor cell of SPD and SPDsh eliminated marker expression in the spicule region; ablation of either cell resulted in one cell per spicule expressing the marker. We also used other cell type-specific markers to examine the terminal cell fates in *lin-17(n698)* mutants. *osm-6::GFP* is normally expressed in SPD and SPV neurons of wild-type spicules and in male sensory ray neurons (Collet et al. 1998). In the *lin-17(n698)* mutant, *osm-6::GFP* is expressed in an extra cell associated with the spicules (four of six animals examined), consistent with the extra expression observed with the *gpa-1::lacZ* marker. Expression of a socket cell marker, *egl-17::GFP* (Burdine et al. 1998; L.I. Jiang and P.W. Sternberg, in prep.), was not affected by the *lin-17(n698)* mutation ($n = 10$), suggesting that the socket cells are correctly specified in *lin-17(n698)* mutants. Therefore, *lin-17* mutants confer a sheath to neuron fate transformation.

son-1 mutant confers the same sheath to neuron fate transformation

To discover additional loci involved in the neurogenic sublineage cell fate specification, we performed an F_1 clonal screen looking for mutations that alter *gpa-1::lacZ* expression specifically in the spicule region. From the screen, we recovered one mutant, *sy549*, in which two cells per spicule express the *gpa-1::lacZ* marker (Fig. 2E). Cell ablation experiments revealed that the extra cell that expresses the *gpa-1::lacZ* marker is the SPDsh cell, as in *lin-17* mutants (Fig. 2E–G; Table 1).

sy549 mutant males have wild-type spicules and wild-type B-cell division pattern. Expression patterns of other cell-type-specific markers examined are consistent with the conclusion that *sy549* confers a SPDsh to SPD neuron fate transformation (*osm-6::GFP*: 7 of 11 animals showed extra marker expression in the spicule region; *egl-17::GFP*: 10 of 10 wild-type expression). Thus, we named this mutant *son-1*.

son-1(sy549) displays several phenotypes resembling *lin-17* mutants

In addition to the male spicule phenotype, *son-1(sy549)* mutants also display a protruding vulva phenotype with high penetrance. Hermaphrodites retain eggs in uterus, some animals have a disorganized gonad and explode at the vulva, and the brood size is low [~ 50 in a *syIs20 him-5(e1490)* background]. Abnormal vulval lineages and vulval morphology are observed in $\sim 10\%$ of animals ($n > 50$). The abnormal vulval lineage occurs in 2° cells; it can occur at either P5.p or P7.p. In most cases, the correct number of cells were generated, but the vulval morphology was abnormal, and an extra vulva invagina-

tion was observed in several animals. This phenotype is similar to the vulval defect observed in *lin-17* mutants (Ferguson and Horvitz 1985; Ferguson et al. 1987). Occasionally, *son-1(sy549)* hermaphrodites have gonads with a single arm instead of two, as observed in *lin-17* mutants (Sternberg and Horvitz 1988). In addition, like *lin-17* mutant males, *son-1(sy549)* mutant males have ray defects and cannot mate.

son-1(sy549) enhances *lin-17(n698)* phenotypes to *lin-17* null like

To examine the interactions between *son-1* and *lin-17*, we constructed strains defective in both *son-1* and *lin-17*. We first examined the *gpa-1::lacZ* expression pattern in the double-mutant animals. The SPD neuron marker is expressed in no more than two cells per spicule in the double mutants, confirming the conclusion from the ablation experiment that both confer the same sheath-to-neuron fate transformation.

lin-17 mutants are defective in vulva formation, somatic gonad development, and phasmid and spicule development in the tail. *n698* is a reduction-of-function allele of *lin-17* (Sawa et al. 1996) and displays weaker phenotypes than a null allele, *n671*. About 50% of *n698* animals have abnormal vulval invaginations, but the extra posterior invagination phenotype is rarely seen (Table 2). In contrast, all *n698; sy549* double-mutant animals examined have abnormal vulval invaginations, and 58% of animals show an extra posterior invagination, which is typically seen only in strong *lin-17* alleles (Table 2). Approximately 73% of the *n698* animals have a T-lineage defect, such that their phasmids fail to take up DiO (Sawa et al. 1996). *son-1(sy549)* mutants are wild type for DiO staining, whereas 98% of *n698; sy549* double-mutant animals fail to uptake DiO (Table 2). Double-mutant hermaphrodites often have a single gonad arm instead of the normal two. Double-mutant males show ectopic meiosis, indicative of no distal tip cells (Sternberg and Horvitz 1988). Double-mutant males also show more severely crumpled spicules and have no rays. Taken together, *lin-17(n698); son-1(sy549)* double-mutant animals behave like strong *lin-17* mutant animals with respect to vulva development, somatic gonad development, and male tail patterning. However, neither *lin-17(n698)* mutants nor the double mutants exhibit a P12 fate specification defect (Table 2).

Next, we constructed a double-mutant strain carrying *son-1(sy549)* and a *lin-17* null allele, *n671*. We examined two relatively low penetrance phenotypes, the ectopic posterior vulva invagination phenotype and P12 fate specification. We did not observe any phenotypic enhancement in the double-mutant animals (Table 2). Therefore, *son-1* and *lin-17* likely act in the same pathway.

son-1 encodes an HMG1/2-like protein

son-1(sy549) is recessive, and on the basis of the spicule phenotype, we mapped it to chromosome III, between

Genotype		Phenotypes			
<i>lin-17</i>	<i>son-1</i>	bivulva ^a	hermaphrodite gonad ^b	P12.p → P11.p ^c	abnormal phasmids ^d
+	<i>sy549</i>	2% (<i>n</i> = 108)	1% (<i>n</i> = 108)	0 (<i>n</i> = 100)	4% (<i>n</i> = 53)
<i>n698</i>	+	2% (<i>n</i> = 46)	8% (<i>n</i> = 46)	0 (<i>n</i> = 100)	73% (<i>n</i> = 346)
<i>n698</i>	<i>sy549</i>	58% (<i>n</i> = 98)	28% (<i>n</i> = 101)	0 (<i>n</i> = 100)	98% (<i>n</i> = 57)
<i>n671</i>	+	64% (<i>n</i> = 66)	23% (<i>n</i> = 66)	29% ^e (<i>n</i> = 41)	95% ^f (<i>n</i> = 400)
<i>n671</i>	<i>sy549</i>	64% (<i>n</i> = 76)	20% (<i>n</i> = 69)	22% (<i>n</i> = 92)	N.D.

^fData from Sawa et al. (1996).

We examined the expression pattern of *son-1* by generating both transcriptional and translational fusions of *son-1* to a GFP reporter gene. The constructs contain either 1- or 3- to 4-kb upstream genomic sequence of

Figure 4. Sequence of the *son-1* gene. (A) *son-1* cDNA sequence and deduced amino acid sequence. The boxed sequence is derived from alternative splicing at the end of exon 1, which leads to additional two amino acids as indicated in parentheses. This alternative splicing does not change the overall protein structure. (Shaded regions) HMG boxes; (***) stop codon; (arrows) molecular lesion of the *sy549* allele. (B) Comparison of the HMG box A and box B sequences between SON-1 and HMG1-like proteins of *Drosophila melanogaster*, *Mus musculus*, and *Homo sapiens*. Consensus residues between SON-1 and other HMG1-like proteins are shaded. (*) Unusual valine residue found in the HMG box A of SON-1 protein.

son-1. We found that *son-1* is expressed in every somatic cell through development (Fig. 5). The transcriptional fusion construct contains only part of HMG box A and does not show nuclear localization of GFP. The translational fusion construct is predicted to encode a fusion protein of full-length SON-1, lacking the last 6 amino acids, followed by GFP. This construct is able to rescue *son-1(sy549)* phenotypes and displays nuclear localized GFP expression. This expression pattern is consistent with the predicted role for HMG-box containing proteins as DNA-binding proteins.

son-1(sy549) mutants are not generally defective in gene transcription

The HMG1/2-like proteins have been traditionally considered as architectural components of chromatin that have general roles in regulating chromosomal functions (Bustin and Reeves 1996). Although several recent studies have started to suggest specific roles for the HMG1/2-like proteins in regulating gene transcription (Lehming et al. 1994; Zwilling et al. 1995; Zappavigna et al. 1996; Boonyaratanakornkit et al. 1998; Jayaraman et al. 1998), the in vivo functions of this class of HMG proteins during development have not been studied. In spite of the broad expression pattern revealed by the *son-1::GFP* reporter gene, we do not think that *son-1(sy549)* generally disrupts transcription. First, *son-1(sy549)* mutants display specific phenotypes in several developmental processes that resemble *lin-17* mutants, and *son-1(sy549)* genetically interacts with *lin-17*.

Second, *son-1(sy549)* mutant males display ectopic transgene expression specifically in the spicule region. If it were a general regulator of transcription, this pheno-

type would suggest that loss-of-function *son-1* might also up-regulate expression of the *gpa-1::lacZ* reporter elsewhere. Yet, we did not observe ectopic or stronger expression of the *gpa-1::lacZ* marker in the phasmid neurons or in the head region. Furthermore, other transgenic lines examined did not show any up-regulation or down-regulation of reporter gene expression by *son-1(sy549)*.

Third, *son-1(sy549)* enhances the phenotypes of a weak allele of *lin-17* to null-like phenotypes. This suggests that loss-of-function *son-1* may down-regulate gene expression. To further test this possibility, we used a sensitized genetic assay that is able to detect a twofold difference in gene expression (Meneely and Wood 1987). We used a dosage-sensitive allele of the *let-2* gene that is involved in developmental processes unrelated to those mediated by *son-1*. The *let-2* gene encodes a type IV $\alpha 2$ collagen, and its different alleles cause animal lethality with varying severity (Sibley et al. 1994). The allele *mn114* causes a strong dosage-dependent phenotype: Animals homozygous for *mn114* can survive to adulthood but are sterile, whereas hemizygous animals die at larval stages (Meneely and Wood 1987). If *son-1* is a general regulator of transcription, it should enhance the phenotype of *let-2(mn114)*. Animals homozygous for both *let-2(mn114)* and *son-1(sy549)* can still survive to adulthood as do *let-2(mn114)* homozygotes. Taken together, our results provide the first evidence that the HMG1/2-like proteins have specific developmental functions in vivo.

Disruption of *son-1* function by RNAi results in defects in several *lin-17*-mediated developmental processes

To obtain possible null phenotypes of *son-1*, we dis-

Figure 5. A GFP reporter construct reveals that *son-1* is expressed in all somatic cell nuclei through development. (A) An L3 male; (B) an L1 hermaphrodite; (C) male tail at L4 molt. Scale bar, 20 μ m.

rupted *son-1* function by RNA-mediated interference. dsRNA were synthesized with a full-length cDNA clone as template and were injected into adult hermaphrodites. We found an embryonic lethal phenotype as well as post-

embryonic abnormalities. The embryonic lethality is ~11% penetrant. The reason for the embryonic lethality is not apparent to us. The dead eggs did not show lack of endoderm fates as seen in *mom* mutants (Rocheleau et al. 1997). Viable hermaphrodites and males grow to adulthood and show a sterile phenotype at ~30% penetrance. Hermaphrodites often have a single gonad arm, resembling the *lin-17* phenotype (Fig. 6A; Sternberg and Horvitz 1988). In males, ectopic meiosis is seen in germ line, indicative of loss of distal tip cells (Fig. 6B; Sternberg and Horvitz 1988). This defect is also seen in *lin-17* mutant males a result of Z1 and Z4 lineage defect in which more linker cells are generated at the expense of distal tip cells (Sternberg and Horvitz 1988). Disruption of *son-1* function by RNAi also resulted in a vulval lineage defect and bursting at the vulva region at low penetrance. Abnormal *gpa-1::lacZ* marker expression at the spicule region mimicking *sy549* phenotype was also observed (Fig. 6C). Therefore, the phenotypes produced by *son-1* RNAi are consistent with those seen in *son-1(sy549)* mutants. The *sy549* allele likely reduces but does not fully eliminate SON-1 functions, consistent with the deficiency result. The strong resemblance of *son-1* and *lin-17* mutant phenotypes further suggests that *son-1* is specifically involved in the *lin-17*-mediated Wnt signaling.

Overexpression of POP-1 results in the same spicule defect as *son-1* RNAi

The TCF/LEF class of HMG box-containing proteins, different from the HMG1/2 class of proteins, has been implicated as a component of Wnt signaling pathway to regulate Wnt target gene expression (van de Wetering et al. 1997). The *pop-1* gene encodes a *C. elegans* homolog of the TCF/LEF class HMG protein and has been implicated in Wnt signaling during embryogenesis (Lin et al. 1995, 1998; Rocheleau et al. 1997; Thorpe et al. 1997). As *pop-1* is thought to negatively regulate Wnt signaling, we wondered if overexpression of POP-1 would result in

Figure 6. Phenotypes of disruption of *son-1* function by RNAi and overexpression of POP-1. (A–C) Phenotypes observed from *son-1* RNAi experiment. (A) Hermaphrodite gonad showing abnormally short gonad arm; (arrow) meiosis at the distal end of the gonad arm. (B) Male gonad (white arrow-head) ectopic germ-line meiosis. (C) *gpa-1::lacZ* is expressed in two cells per spicule. All four blue cells can be seen on this focal plane, two from the left spicule and two from the right one. Cells from the same side are indicated by linkage of two arrow-heads. (D) When POP-1 is overexpressed in wild-type males, *gpa-1::lacZ* expression is observed in two cells per spicule (black arrowheads). Lateral view, showing only the left spicule. Scale bar, 20 μ m.

similar defects such as loss-of-function of *son-1*. When POP-1 is overexpressed at early to mid L3 stage in wild-type males, they show two cells per spicule expressing the SPD marker, resembling the *son-1(sy549)* mutant phenotype (Fig. 6D). Thus, the two HMG box-containing proteins, POP-1 and SON-1, appear to have opposite roles in regulating Wnt signaling. POP-1 negatively regulates Wnt signaling, whereas SON-1 positively regulates Wnt signaling.

Although *pop-1(zu189)* has a maternal effect embryonic lethal phenotype, this mutant does not display post-embryonic defects. Homozygous animals produced by a heterozygous mother are viable and develop normally. Using the spicule SPD neuron-specific marker, we found that *pop-1* mutant males have a wild-type marker expression pattern in the spicules. Mutant animals defective in both *pop-1* and *son-1* genes display additive phenotypes. They still exhibit maternal effect embryonic lethality, whereas all of the postembryonic phenotypes resemble *son-1(sy549)* mutants.

son-1 does not act upstream of Wnt

To determine how *son-1* is involved in Wnt signaling, we first tested whether *son-1* acts in the Wnt signaling cells or in the responding cells. The Wnt signal LIN-44 is expressed in the tail region and plays a role in tail patterning (Herman and Horvitz 1994; Herman et al. 1995). We think that LIN-44 is the signal that acts through the LIN-17 receptor during male spicule development. First, during the first division of B lineage, LIN-44 acts through LIN-17 to mediate the asymmetric cell division (Chamberlin and Sternberg 1995). Second, *lin-44* mutants exhibit a similar SPD neuron marker expression pattern in the spicule region as *lin-17* and *son-1* mutants. We reasoned that if *son-1* acts in the signaling cells to regulate *lin-44* expression, we should be able to rescue *son-1* mutant phenotypes by overexpression of the Wnt signal LIN-44. To test this hypothesis, we used a construct containing full-length *lin-44* cDNA driven by a heat-shock promoter. This construct has been shown to rescue *lin-44* phenotypes (Herman et al. 1995). When LIN-44 overexpression is induced during the L2/L3 stage in *son-1(sy549)* mutant background, it does not affect the spicule phenotype of *son-1(sy549)* mutants. This result indicates that *son-1* does not act upstream of the Wnt signal to regulate its expression. Because *son-1* is expressed in cell nuclei, it is likely that *son-1* acts in the Wnt-responding cells to regulate Wnt pathway gene expression. However, we cannot rule out the possibility that *son-1* is involved in the post-translational regulation of the Wnt signal.

We then tested whether overexpression of SON-1 is able to suppress *lin-17* mutant phenotypes. We engineered a *hs-son-1* construct that contains a genomic clone encoding full-length *son-1* driven by a heat-shock promoter. In a *son-1(sy549)* background, overexpression of SON-1 by a single pulse of heat shock at the L3 stage within 4 hr prior to the cell division is able to rescue the spicule sheath to neuron fate transformation phenotype

of the mutant. In a *lin-17(n698)* mutant background, we induced SON-1 expression during embryogenesis to the L1/L2 stage and at the L2/L3 stage and examined the phasmid and spicule phenotypes, respectively. We did not observe any suppression of *lin-17* phenotypes by SON-1 overexpression. *son-1* might act at the level of Wnt receptor to regulate *lin-17* function. Another possibility is that *son-1* activity is not rate limiting for Wnt pathway function. Consistent with this hypothesis, simply overexpressing SON-1 in a wild-type background by heat shock or injection of the wild-type gene at high dosage does not result in any gain-of-function phenotype.

Structure/function analysis of SON-1 protein

Both HMG boxes A and B of the HMG1/2-like proteins have been shown to be able to bind bent DNA structures and to interact with certain transcription factors independently in vitro (Bianchi et al. 1992; Zappavigna et al. 1996). To examine the in vivo functions of SON-1 HMG boxes A and B, we engineered deletion constructs lacking either individual box and generated transgenic animals bearing these constructs (Fig. 7). In a wild-type background, we did not observe any dominant-negative or gain-of-function effect caused by these constructs. When these transgenes are crossed into a *son-1(sy549)* mutant background, only the construct containing both HMG boxes is able to rescue the mutant phenotypes. None of the constructs containing a single HMG box were able to rescue the mutant phenotypes (Fig. 7). This result suggests that both HMG boxes A and B are required for the function of the SON-1 protein in vivo.

Discussion

Wnt signaling in spicule development

During *C. elegans* male spicule development, the Wnt

Figure 7. Structure/function analysis of HMG boxes. HMG boxes are indicated as open box and labeled. Bold lines indicate the acidic tail. All constructs contain additional 1-kb upstream and 1-kb downstream genomic sequences. (a) Constructs were tested in wild-type background for any dominant-negative or gain-of-function effect. (b) Constructs were tested in *son-1(sy549)* mutant background for any rescuing activity. The number of transgenic lines tested in which >75% of the animals showed wild-type SPD marker expression is indicated; these were considered positive for rescuing activity. (+) Positive for tested activity; (–) negative for tested activity.

pathway genes *lin-44* and *lin-17* control the first cell division of the spicule lineage, namely, the B lineage (Sternberg and Horvitz 1988; Herman and Horvitz 1994; Chamberlin and Sternberg 1995; Sawa et al. 1996). The first division of the B cell, B.a versus B.p, is asymmetric. Inactivation of the Wnt pathway results in reversed polarity or posterior B.p to anterior B.a fate transformation. Additional functions for *lin-17* have also been implicated in several asymmetric cell divisions in both B.a and B.p lineages after the 10-cell stage (Chamberlin and Sternberg 1995). Using a terminal cell fate marker, we discovered that the Wnt pathway genes *lin-17* and likely *lin-44* are involved in the fate specification of SPD neuron (ζ .xxa) versus SPDsh (ζ .xpx) during the last round cell division of the B lineage. Loss-of-function mutation in *lin-17* results in a posterior sheath to anterior neuron fate transformation. Therefore, the *lin-17*-mediated Wnt signaling pathway is used at the first and the last cell division of B cell development to mediate cell fate specification. Like most of the other Wnt-mediated developmental processes, the Wnt pathway acts to promote the posterior fate in the *C. elegans* male spicule lineage. We envision that the source of Wnt signal LIN-44, the hyp8-11 cells at the tip of the tail (Herman et al. 1995), sets up the anterior and posterior polarity for general tail patterning.

In *C. elegans*, several Wnt pathways mediate several distinct developmental processes. Some components are unique, others are shared in several processes. We examined the possible involvement of other Wnt pathway components in this last cell division of male spicule development. In *lin-44* mutant males, the abnormal SPD neuron marker expression in the spicule region is observed, resembling that of the *lin-17* mutants. However, all of the *lin-44* alleles available are null or strong alleles, wherein the early B lineage defect is highly penetrant, making it difficult to rigorously examine its effect on the later stage cell fate specification. *mom-1* and *mom-3* homozygous males display a severely crumpled spicules phenotype at high penetrance, indicating an early B lineage defect. SPD neuron marker expression appeared to be normal in *bar-1(sy324)* and *bar-1(mu349)* mutant males. *pop-1* mutant males have wild-type spicules and normal marker expression pattern. However, overexpression of POP-1 at the L3 stage results in extra cell expressing *gpa-1::lacZ* in the spicules, similar to loss-of-function *son-1* mutants.

During the development of *C. elegans* male spicules, the neurogenic sublineages β and ζ resemble the *Drosophila* SOP lineage in that a progenitor cell divides twice and generates neuron, sheath, and socket cells. The mechanism used for ζ lineage terminal cell fate specification appears to be different from that for the SOP lineage. In the development of *Drosophila* SOP lineage, sister cell interaction mediated by the *Notch/Delta* pathway plays an important role for the specification of neuronal versus non-neuronal cell fate (for review, see Greenwald 1998). However, we did not observe any effect of *lin-12* on the specification of SPD versus SPDsh fate in *C. elegans* male spicule development. Rather, the

lin-17-mediated Wnt signaling pathway controls this fate decision. Thus the two different signaling pathways, Notch and Wnt, are used in different organisms to achieve a similar cell fate patterning in a conserved neurogenic lineage.

However, the Wnt pathway is not the sole player for spicule development. The β lineage closely resembles ζ lineage in generating the neuron and non-neuronal structural cells. Yet the Wnt pathway does not seem to be involved to the neuron/sheath fate specification derived from β lineage as revealed by other molecular markers (data not shown). In contrast to the ζ lineage, cell-cell interactions within the β lineage do play an important role during the development of the β lineage, because β lineage can be easily perturbed by cell ablations or mutants of the *lin-3/let-23* EGF/EGFR pathway components (Chamberlin and Sternberg 1993, 1994). Therefore, multiple signaling pathways must be utilized during male spicule development.

An HMG1/2-like protein is involved in Wnt signaling

We identified a new gene, *son-1*, involved in the SPD neuron versus SPDsh fate specification in a similar fashion as *lin-17* during *C. elegans* male spicule development. *son-1* encodes an HMG1/2-like protein. A GFP reporter gene revealed that *son-1* is broadly expressed through development and its expression is localized in cell nuclei, consistent with its being an HMG-box containing DNA-binding protein.

Although the HMG1/2 class proteins are traditionally considered as architectural components of chromatin, the specific phenotypes of *son-1(sy549)* suggest that this gene may be specifically involved in Wnt signaling. First, *son-1(sy549)* causes a similar SPDsh to neuron fate transformation as does a *lin-17* hypomorph. The vulva and gonad defects observed in *son-1(sy549)* mutants, although at low penetrance, do resemble what are seen in *lin-17* mutants. Second, we observed specific postembryonic phenotypes that resemble *lin-17* mutant phenotypes from *son-1 RNAi* experiments. Third, *son-1(sy549)* enhances *lin-17(n698)* phenotypes to *lin-17* null-like but does not enhance phenotypes of other unrelated mutants such as *let-2(mn114)*. Fourth, overexpression of POP-1 gave similar spicule phenotype as disruption of *son-1* function. Taken together, *son-1* is involved in several Wnt-mediated developmental processes, including vulva formation, somatic gonad development, and male spicule development. Our results provide the first evidence for the specific functions of the HMG1/2 class proteins in vivo.

The HMG1/2-like protein encoded by *son-1* has several interesting features that are different from what is known about the HMG1/2 class proteins. The HMG1/2 class of proteins is classified as sequence nonspecific HMG proteins in comparison to the sequence-specific HMG proteins, such as TCF/LEF (Bianchi et al. 1992). The canonical HMG box found in sequence nonspecific HMG proteins and the generalized HMG box found in sequence-specific HMG proteins share substantial se-

quence homology and identical protein structure. The sequence specificity is thought to reside in the first 12 amino-terminal residues and the last 25 carboxy-terminal residues of the HMG box. In particular, the amino acids at positions 7 and 12 have been suggested to be crucial for sequence-specific binding (Read et al. 1994). The residues at positions 7 and 12 are valine (V) or isoleucine (I) and asparagine (N) in all sequence-specific HMG boxes. Instead, all presently known sequence non-specific HMG boxes have proline (P) and serine (S) at these two positions. The box B of SON-1 protein appears to be consistent with this finding (positions 102 and 107 of the protein, Fig. 4B). Interestingly, the box A of SON-1 protein has a valine (V) residue at position 7 (position 14 of the protein, Fig. 4B). This raises the possibility that the *C. elegans* HMG1/2-like protein SON-1 has more specific functions than the HMG1/2-like proteins in other organisms. Furthermore, our data suggest that both HMG boxes are required *in vivo* for SON-1 functions, in contrast to the *in vitro* observations that each individual HMG box is able to recognize bent DNA structure and to interact with certain transcription factors in a similar fashion as is the whole protein (Bianchi et al. 1992; Zappavigna et al. 1996). These features of SON-1 may be indicative of its specific functions.

Two classes of HMG proteins, SON-1 and POP-1, play opposite roles in Wnt signaling

The TCF/LEF class sequence-specific HMG proteins have been implicated in Wnt signaling. They can either positively or negatively regulate Wnt target gene expression depending on their transcription cofactors (van de Wetering et al. 1997; Waltzer and Bienz 1998). The *C. elegans* TCF/LEF homolog, *pop-1*, appears to negatively regulate Wnt signaling as loss-of-function in *pop-1* causes an opposite phenotype as inactivation of the Wnt signaling during *C. elegans* embryogenesis. In the specification of SPD neuron versus SPDsh fate during male spicule development, overexpression of POP-1 results in the same defect as loss-of-function mutation in *lin-17* and *son-1*. Therefore, a POP-1-like activity also negatively regulates Wnt signaling during *C. elegans* postembryonic development. Both the sequence-specific TCF/LEF and sequence-nonspecific HMG1/2 class of proteins are involved in *lin-17*-mediated Wnt signaling and they play opposite roles in regulating Wnt pathway activity.

pop-1 has been shown to act downstream of the Wnt receptor and its expression is negatively regulated by Wnt signaling (Rocheleau et al. 1997; Thorpe et al. 1997; Lin et al. 1998). Inactivation of the Wnt pathway upregulates POP-1 protein level. Here we show that *son-1* also acts in the Wnt responding cells. Therefore, both the POP-1-like protein and SON-1 act in the same cell nucleus to regulate gene transcription. The POP-like activity inhibits gene transcription, and SON-1 facilitates gene transcription (Fig. 8). As SON-1 is an HMG-1/2-like protein, we think that SON-1 could activate gene transcription in two ways. First, SON-1 could alter chromatin conformation in favor of gene expression. Second,

Figure 8. Model for SON-1 function. SON-1 and a POP-1-like protein antagonistically regulate gene expression in Wnt-responding cells. The POP-1-like activity inhibits gene transcription, whereas SON-1 facilitates gene transcription. SON-1 might facilitate gene transcription after the inhibitory effect of the POP-1-like activity is relieved on activation of the Wnt pathway. SON-1 could change the chromatin conformation in favor of gene expression. Alternatively, SON-1 could interact with other transcription factors to help stimulate gene expression. (POP-1) A POP-1-like protein; (TF) transcription factor.

SON-1 could physically interact with other transcription factors and help stimulate specific gene expression. Proteins of the HMG1/2 class have been shown to interact with POU domain proteins or HOX proteins *in vitro* and enhance their DNA-binding and transcriptional activities (Zwilling et al. 1995; Zappavigna et al. 1996). Our data suggest that *son-1* function may not be rate limiting for Wnt signal transduction. Moreover, *son-1::GFP* expression does not seem to be altered in a *lin-17* mutant background. Therefore, it is possible that SON-1 acts to activate gene transcription after the inhibitory effect of POP-1 is relieved by activation of the Wnt pathway.

Materials and methods

Strains and alleles

Routine culturing, maintenance and genetic manipulations of *C. elegans* strains were performed according to standard procedures (Brenner 1974). N2 (Bristol, UK) strain was used as wild type (Brenner 1974). The following strains were used and are referenced in Riddle et al. (1997) or as indicated: LG I-*lin-17*(n671), *lin-17*(n698) (Ferguson and Horvitz 1985), *pop-1*(zu189) (Lin et al. 1995), *dpy-5*(e61), *hT1*, *hT2*; LG II-*dpy-10*(e128), *rol-1*(e91) *mom-3*(or78) (Thorpe et al. 1997), *mnC1*[*dpy-10*(e128) *unc-52*(e444)]; LG III-*dpy-17*(e164), *lon-1*(e185), *unc-36*(e251), *unc-32*(e189), *dpy-18*(e364), *sDf125*, *sDf127*, *sDf135*, *nDf16*, *nDf20*, *nDf21*, *pha-1*(e2123ts) (Granato et al. 1994), *hT2*, *qC1*, *dpy-19*(e1259) *lin-12*(n137)/*unc-32*(e189) *lin-12*(n676n909); LG IV-*dpy-20*(e1282), *dpy-13*(e184); LG V-*syIs20*[*gpa-1::lacZ*+*dpy-20*(+)], *him-5*(e1490), *dpy-11*(e224); LG X-*bar-1*(sy324) (A. Golden and P.W. Sternberg, unpubl.), *bar-1*(mu349) (kindly provided by J. Whangbo and C. Kenyon, University of California, San Francisco), *lon-2*(e678), *mom-1*(or10) *unc-6*(n102) (Thorpe et al. 1997).

Assay for β -galactosidase activity was performed as described previously (Fire 1992).

Anatomical and lineage analysis/cell ablation/photographs

Animal anatomy and cell division patterns were examined under Nomarski optics at 20°C as described by Sulston and Horvitz (1977). For laser ablation, L3 males were mounted on 5% agar pad and anesthetized with 1%–3% sodium azide. Laser killing cells was performed as described (Sulston and White 1980; Avery and Horvitz 1987, 1989). Worms were then recovered from the agar pad and allowed to grow to adulthood, at which time we examined the *gpa-1::lacZ* marker expression. Photographs of Nomarski images were taken on Kodak Tech Pan film. Photographs of β -galactosidase expression images were taken on Kodak Royal Gold ASA 25 films. GFP expression was examined with a Chroma High Q GFP LP filter set (450 nm excitation/505 nm emission) and photos were taken on Fuji Provia ASA 400 film.

Isolation of the son-1(sy549) allele

By use of the SPD neuron marker, *gpa-1::lacZ*, an F₁ clonal screen was designed to look for mutants that specifically disrupt the terminal cell fate specification during male spicule development. Standard EMS mutagenesis was used (Brenner 1974). The *sy549* allele was recovered from a screen of ~6300 haploid genomes.

Cloning of son-1

son-1(sy549) was mapped to chromosome III by two-factor mapping. Three-factor mapping put *son-1(sy549)* between *lon-1* and *unc-36*. Markers used for three-factor mapping were *dpy-17(e164) unc-32(e189)*, *dpy-17(e164) unc-36(e251)*, and *lon-1(e185) unc-32(e189)*. Deficiency mapping indicated that both *sDf125* and *sDf127* delete *son-1(sy549)*, but *nDf16*, *nDf20*, and *nDf21* do not. Therefore, *son-1* mutation lies between the left end point of *sDf127* and the left end point of *nDf16*. Cosmid pools or a single cosmid in that region were injected into *son-1(sy549) pha-1(ts); syIs20 him-5* strain at a concentration of 25 ng/μl each to test for rescue. Those cosmids include C23G10, T12A2, C18F10, F47D12, C56G2, C16A3, C05D11, T26A5, F23F12, K03F8, C09E7, F27B3, F20H11, C13B9, M01G4, F37A4, K07E12, R05H11, F28F5, and R13F6. *pha-1* was used as a germline transformation marker (Granato et al. 1994). Rescuing activity was scored primarily on the basis of the spicule phenotype and other phenotypes were examined as well. A single cosmid C18F10 rescued *son-1(sy549)* phenotypes in all respects. Using primers LJ1(5'-CCCAGTTCTTACTGCTTCGCATC-3') and LJ2(5'-CATGAAAACCTCCACTGCAACGG-3'), we amplified a 3-kb fragment by PCR from the 3' end of C18F10 which contains a single gene *F47D12.4*. Three independent PCR clones were injected into *son-1(sy549)* mutants and all of them were able to rescue the mutant phenotypes.

son-1 cDNA was obtained by RT-PCR with wild-type N2 total RNA as template and using primers complementary to *son-1* start and stop codon regions on the basis of GeneFinder prediction. The ~600-bp products from five independent RT-PCR clones were cloned into pBluescript and sequenced. Three clones are consistent with GeneFinder prediction, the other two have an alternative splice site at the end of the first exon and result in an additional 2 amino acids without changing the rest of the protein coding. The 5' end of the gene was confirmed by RT-PCR by use of SL1 primer and an internal gene-specific primer (LJ6: 5'-ACAGGGCTCTTTTGGTGC-3'). A cDNA clone yk74h7 has been reported recently on the web database by Dr. Y. Kohara (National Institute of Genetics, Mishima, Japan). It may predict an extra exon of 32 amino acids at the 5' end of *son-1*.

The 1-kb genomic region of the *F47D12.4* gene was PCR amplified from small numbers of *sy549* mutants (Aroian et al. 1994) and sequenced on an automated sequencing machine (Applied Biosystems). Four independent PCR clones showed the same nucleotide change of g → a at nucleotide position 228 (two plus strands, two minus strands). This change results in a substitution of asparagine for aspartic acid 72.

Molecular biology

son-1-GFP The transcriptional fusion GFP constructs contain either 1- or 4.5-kb upstream genomic sequence and the first 64 amino acids of *son-1* protein (at the *EcoRV* site). The upstream sequence was PCR amplified from cosmid C18F10, the coding region was derived from the 3-kb rescuing fragment. The translational fusion GFP constructs (pLJ33/pJ35) contain PCR fragment of either 1- or 3.2-kb upstream genomic sequence and full length, but the last 6 amino acids of SON-1 protein cloned into GFP vectors from the Fire laboratory (Carnegie Institute of Washington).

hs-son-1 A 1.4-kb *EcoRI* fragment of *son-1* genomic sequence was cloned into vectors containing heat shock promoter *hsp16-2* or *hsp16-48* to generate pLJ25 and pLJ26, respectively.

son-1 deletion constructs To create *son-1* deletion constructs, we started from the 3-kb rescuing fragment. The 3-kb fragment was cloned into pCRII vector (Invitrogen, pLJ13). For the box A deletion construct (pLJ41), a *SmaI-EcoRV* fragment was removed and religated, which deleted amino acids 7–76. For the box B deletion construct (pLJ40), sequences between *Bam*HI and *Spe*I was deleted, thereby removing amino acids 99–162. For the box A construct only (pLJ39), a stop codon was generated after the first 98 amino acids. All constructs still contain 1-kb upstream and 1-kb downstream genomic sequence of *son-1*.

Generation of transgenic animals

Transgenic lines carrying an extrachromosomal array of testing plasmids were generated by standard germ-line injection and a DNA transformation method (Mello and Fire 1995). GFP constructs were injected into *pha-1(ts); him-5(e1490)* strain at a concentration of 20 ng/μl together with *pha-1* rescuing plasmid pBX-1 at 100 ng/μl. Heat shock constructs and *son-1* deletion constructs were injected into *pha-1(ts); syIs20 him-5* strain. Injection concentrations for heat-shock constructs containing *son-1* genomic DNA, *lin-44* cDNA, and *pop-1* cDNA are 50, 100, and 20 or 100 ng/μl, respectively. *son-1* deletion constructs were injected at 50 and 100 ng/μl. To test the ability to rescue *son-1* phenotypes, transgenic lines were further crossed into *son-1(sy549) pha-1(ts); syIs20 him-5* strain.

RNA interference experiments

Double-stranded RNA was synthesized as described (Fire et al. 1998) by use of full-length *son-1* cDNA as template. Approximately 40–50 ng/μl dsRNA was injected into *syIs20 him-5* hermaphrodites. Eggs laid between 3 and 51 hr after injection were collected at 12-hr intervals. Phenotypes of F₁s were analyzed in detail. Unrelated dsRNA and DEPC-H₂O were used as controls for the experiment.

Heat shock experiments

Worms were staged under Nomarski optics and heat shocked in a 33°C water bath for 30, 45, or 60 min.

Acknowledgments

We thank Jane Mendel for the *syIs20 him-5* strain, Mike Herman for the *lin-17(n698)* allele and the *hs-lin44* constructs, Helen Stewart for the *sDf* strains and *sDf* break-point mapping data, Rueylin Lin for the *hs-pop-1* construct, Jennifer Whangbo for the *bar-1(mu349)* allele, the CGC for strains, and the Sanger center for cosmid and sequencing information. We thank members of our laboratory for discussion and comments on the manuscript. This work was supported by a U.S. Public Health Service grant HD23690 to P.W.S. and by the Howard Hughes Medical Institute, with which P.W.S. is an investigator.

The publication costs of this article were defrayed in part by payment of page charges. This article must therefore be hereby marked 'advertisement' in accordance with 18 USC section 1734 solely to indicate this fact.

References

- Aroian, R.V., G.M. Lesa, and P.W. Sternberg. 1994. Mutations in the *Caenorhabditis elegans* *let-23* EGFR-like gene define elements important for cell-type specificity and function. *EMBO J.* **13**: 360–366.
- Avery, L. and H.R. Horvitz. 1987. A cell that dies during wild-type *C. elegans* development can function as a neuron in a *ced-3* mutant. *Cell* **51**: 1071–1078.
- Avery, L. and H.R. Horvitz. 1989. Pharyngeal pumping continues after laser killing of the pharyngeal nervous system of *C. elegans*. *Neuron* **3**: 473–485.
- Bianchi, M.E., L. Falcioni, S. Ferrari, and D.M.J. Lilley. 1992. The DNA binding site of HMG1 protein is composed of two similar segments (HMG boxes), both of which have counterparts in other eukaryotic regulatory proteins. *EMBO J.* **11**: 1055–1063.
- Boonyaratankornkit, V., V. Melvin, P. Prendergast, M. Altmann, L. Ronfani, M.E. Bianchi, L. Taraseviciene, S.K. Nordeen, E.A. Allegretto, and D.P. Edwards. 1998. High-mobility group chromatin proteins 1 and 2 functionally interact with steroid hormone receptors to enhance their DNA binding in vitro and transcriptional activity in mammalian cells. *Mol. Cell. Biol.* **18**: 4471–4487.
- Brenner, S. 1974. The genetics of *Caenorhabditis elegans*. *Genetics* **77**: 71–94.
- Burdine, R.D., C.S. Branda, and M.J. Stern. 1998. *egl-17(fgf)* expression coordinates the attraction of the migrating sex myoblasts with vulval induction in *C. elegans*. *Development* **125**: 1083–1093.
- Bustin, M. and R. Reeves. 1996. High mobility group chromosomal proteins: Architectural components that facilitate chromatin function. *Prog. Nucleic Acid Res. Mol. Biol.* **54**: 35–100.
- Cadigan, K.M. and R. Nusse. 1997. Wnt signaling: A common theme in animal development. *Genes & Dev.* **11**: 3286–3305.
- Chamberlin, H.M. and P.W. Sternberg. 1993. Multiple cell interactions are required for fate specification during male spicule development in *Caenorhabditis elegans*. *Development* **118**: 297–324.
- . 1994. The *lin-3/let-23* pathway mediates inductive signaling during male spicule development in *Caenorhabditis elegans*. *Development* **120**: 2713–2721.
- . 1995. Mutations in the *Caenorhabditis elegans* gene *vab-3* reveal distinct roles in fate specification and unequal cytokinesis in an asymmetric cell division. *Dev. Biol.* **170**: 679–689.
- Collet, J., C.A. Spike, E.A. Lundquist, J.E. Shaw, and R.K. Herman. 1998. Analysis of *osm-6*, a gene that affects sensory cilium structure and sensory neuron function in *Caenorhabditis elegans*. *Genetics* **148**: 187–200.
- Eisenmann, D.M., J.N. Maloof, J.S. Simske, C. Kenyon, and S.K. Kim. 1998. The β -catenin homolog BAR-1 and LET-60 RAS coordinately regulate the *hox* gene *lin-39* during *Caenorhabditis elegans* vulval development. *Development* **125**: 3667–3680.
- Ferguson, E.L. and H.R. Horvitz. 1985. Identification and characterization of 22 genes that affect the vulval cell lineages of the nematode *Caenorhabditis elegans*. *Genetics* **110**: 17–72.
- Ferguson, E.L., P.W. Sternberg, and H.R. Horvitz. 1987. A genetic pathway for the specification of the vulval cell lineages of *Caenorhabditis elegans*. *Nature* **326**: 259–267.
- Fire, A. 1992. Histochemical techniques for locating *Escherichia coli* beta-galactosidase activity in transgenic organisms. *Genet. Anal.* **9**: 151–158.
- Fire, A., S. Xu, M.K. Montgomery, S.A. Kostas, S.E. Driver, and C.C. Mello. 1998. Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. *Nature* **391**: 806–811.
- Greenwald, I. 1997. Development of the vulva. In *Caenorhabditis elegans II* (ed. D.L. Riddle, T. Blumenthal, B.J. Meyer, and J.R. Priess) pp. 519–541. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Granato, M., H. Schnabel, and R. Schnabel. 1994. *pha-1*, a selectable marker for gene-transfer in *C. elegans*. *Nucleic Acid Res.* **22**: 1762–1732.
- Hartenstein, V. and J.W. Posakony. 1990. A dual function of the *Notch* gene in *Drosophila* sensillum development. *Dev. Biol.* **142**: 13–30.
- Herman, M.A. and H.R. Horvitz. 1994. The *Caenorhabditis elegans* gene *lin-44* controls the polarity of asymmetric cell divisions. *Development* **120**: 1035–1047.
- Herman, M.A., L.L. Vassilieva, H.R. Horvitz, J.E. Shaw, and R.K. Herman. 1995. The *C. elegans* gene *lin-44*, which controls the polarity of certain asymmetric cell divisions, encodes a Wnt protein and acts cell nonautonomously. *Cell* **83**: 101–110.
- Hill, R.J. and P.W. Sternberg. 1992. The gene *lin-3* encodes an inductive signal for vulval development in *C. elegans*. *Nature* **358**: 470–476.
- Jayaraman, L., N.C. Moorthy, K.G. Murthy, J.L. Manley, M. Bustin, and C. Prives. 1998. High mobility group protein-1 (HMG-1) is a unique activator of p53. *Genes & Dev.* **12**: 462–472.
- Jiang, L.I. and P.W. Sternberg. 1998. Interactions of EGF, Wnt and HOM-C genes specify the P12 neuroectoblast fate in *C. elegans*. *Development* **125**: 2337–2347.
- Lehming, N., D. Thanos, J.M. Brickman, J. Ma, T. Maniatis, and M. Ptashne. 1994. An HMG-like protein that can switch a transcriptional activator to a repressor. *Nature* **371**: 175–179.
- Lin, R.L., S. Thompson, and J.R. Priess. 1995. *pop-1* encodes an HMG box protein required for the specification of a mesoderm precursor in early *C. elegans* embryos. *Cell* **83**: 599–609.
- Lin, R.L., R.J. Hill, and J.R. Priess. 1998. POP-1 and anterior-posterior fate decisions in *C. elegans* embryo. *Cell* **92**: 229–239.
- Mello, C.C. and A. Fire. 1995. *Methods in Cell Biology: C. elegans*. Academic Press, New York, NY.
- Meneely, P.M. and W.B. Wood. 1987. Genetic analysis of X-chromosome dosage compensation in *Caenorhabditis elegans*. *Genetics* **117**: 25–41.
- Moon, R.T., J.D. Brown, and M. Torres. 1997. Wnts modulate

- cell fate and behavior during vertebrate development. *Trends Genet.* **13**: 157–162.
- Parks, A.L. and M.A. Muskavitch. 1993. *Delta* function is required for bristle organ determination and morphogenesis in *Drosophila*. *Dev. Biol.* **157**: 484–496.
- Read, C.M., P.D. Cary, N.S. Preston, M. Lenicek-Allen, and C. Crane-Robinson. 1994. The DNA-sequence specificity of HMG boxes lies in the minor wing of the structure. *EMBO J.* **13**: 5639–5646.
- Riddle, D.L., T. Blumenthal, B.J. Meyer, and J.R. Priess, eds. 1997. *C. elegans* II. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Rocheleau, C.E., W.D. Downs, R. Lin, C. Wittmann, Y. Bei, Y.-H. Cha, M. Ali, J.R. Priess, and C.C. Mello. 1997. Wnt signaling and an APC-related gene specify endoderm in early *C. elegans* embryos. *Cell* **90**: 707–716.
- Roehl, H. and J. Kimble. 1993. Control of cell fate in *C. elegans* by a GLP-1 peptide consisting primarily of ankyrin repeats. *Nature* **364**: 632–635.
- Sawa, H., L. Lobel, and H.R. Horvitz. 1996. The *Caenorhabditis elegans* gene *lin-17*, which is required for certain asymmetric cell divisions, encodes a putative protein similar to seven-transmembrane protein similar to the *Drosophila* Frizzled protein. *Genes & Dev.* **10**: 2189–2197.
- Sibley, M.H., P.L. Graham, N. Vonmende, and J.M. Kramer. 1994. Mutations in the alpha-2(IV) basement-membrane collagen gene of *Caenorhabditis elegans* produce phenotypes of differing severities. *EMBO J.* **13**: 3278–3285.
- Sternberg, P.W. and H.R. Horvitz. 1988. *lin-17* mutations of *Caenorhabditis elegans* disrupt certain asymmetric cell divisions. *Dev. Biol.* **130**: 67–73.
- Sulston, J.E. and H.R. Horvitz. 1977. Postembryonic cell lineages of the nematode *Caenorhabditis elegans*. *Dev. Biol.* **56**: 110–156.
- Sulston, J.E. and J.G. White. 1980. Regulation and cell autonomy during postembryonic development of *Caenorhabditis elegans*. *Dev. Biol.* **78**: 577–597.
- Thorpe, C., A. Schlesinger, J.C. Carter, and B. Bowerman. 1997. Wnt signaling polarizes an early *C. elegans* blastomere to distinguish endoderm from mesoderm. *Cell* **90**: 695–705.
- van de Wetering, M., R. Cavallo, D. Dooijes, M. van Beest, J. van Es, J. Loureiro, A. Ypma, D. Hursh, T. Jones, A. Bejsovec, M. Peifer, M. Mortin, and H. Clevers. 1997. Armadillo coactivates transcription driven by the product of the *Drosophila* segment polarity gene *DTCF*. *Cell* **88**: 789–799.
- Waltzer, L. and M. Bienz. 1998. *Drosophila* CBP represses the transcription factor TCF to antagonize Wingless signaling. *Nature* **395**: 521–525.
- Zappavigna, V., L. Falciola, M.H. Citterich, F. Mavilio, and M.E. Bianchi. 1996. HMG1 interacts with HOX proteins and enhances their DNA binding and transcriptional activation. *EMBO J.* **15**: 4981–4991.
- Zwilling, S., H. König, and T. Wirth. 1995. High mobility group protein 2 functionally interacts with the POU domains of octamer transcription factors. *EMBO J.* **14**: 1198–1208.

An HMG1-like protein facilitates Wnt signaling in *Caenorhabditis elegans*

Lily I. Jiang and Paul W. Sternberg

Genes Dev. 1999, **13**:

References

This article cites 44 articles, 14 of which can be accessed free at:
<http://genesdev.cshlp.org/content/13/7/877.full.html#ref-list-1>

License

Email Alerting Service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).