RESEARCH COMMUNICATION

Dll4, a novel Notch ligand expressed in arterial endothelium

John R. Shutter,¹ Sheila Scully,² Wei Fan,³
William G. Richards,³ Jan Kitajewski,⁴
Gisele A. Deblandre,⁵ Chris R. Kintner,⁵
and Kevin L. Stark^{1,6}

¹Departments of Molecular Genetics, ²Pathology, and ³Neurobiology, Amgen, Inc., Thousand Oaks, California 91320 USA; ⁴Department of Pathology and OB/GYN, Columbia University, College of Physicians and Surgeons, New York, New York 10032 USA; ⁵Molecular Neurobiology Laboratory, The SALK Institute for Biological Studies, La Jolla, California 90037 USA

We report the cloning and characterization of a new member of the Delta family of Notch ligands, which we have named *Dll4*. Like other *Delta* genes, *Dll4* is predicted to encode a membrane-bound ligand, characterized by an extracellular region containing several EGF-like domains and a DSL domain required for receptor binding. In situ analysis reveals a highly selective expression pattern of *Dll4* within the vascular endothelium. The activity and expression of *Dll4* and the known actions of other members of this family suggest a role for *Dll4* in the control of endothelial cell biology.

Received December 16, 1999; revised version accepted April 7, 2000.

Cell-to-cell communication is required for many biological processes such as differentiation, proliferation, and homeostasis. One system utilized by a wide range of eukaryotes is the Notch-signaling pathway. The *Notch* gene family consists of structurally conserved cell surface receptors that are activated by one or more membrane-bound ligands of the emerging *Delta/Serrate/Lag-2* (DSL) gene family (Fleming 1998). Previous studies with invertebrates have shown that the Notch-signaling pathway allows a subset of cells within an equipotent precursor population to acquire a specific cell fate in a spatially and temporally restricted manner. In this process, known as lateral inhibition or specification, cells destined for one fate inhibit neighboring cells from adopting the same fate (Artavanis-Tsakonas et al. 1999). In *Drosophila*, for example, equipotent cells in the embryonic ectoderm give rise to both neural and epidermal cells. As cells assume a neural fate, expression of neurogenic genes results in increased expression of Delta on the cell surface, leading to an inhibition of these same

genes in neighboring cells via activation of their Notch receptors. Notch-receptor activation results in the transactivation of the *Enhancer of Split complex* [*E(spl)-C*] which, in turn, suppresses the expression of downstream genes, including *Delta*, necessary for neurogenesis (Artavanis-Tsakonas et al. 1999). Accordingly, mutant flies lacking Notch have excessive numbers of neuronal cells, whereas activating mutations of Notch lead to an inhibition of neuron production (Struhl et al. 1993). Although Delta–Notch interactions in vertebrates are less well understood, the identification of several family members suggests that these genes play an important role in mammalian development.

Four mammalian Notch receptors have been identified, *Notch1/TAN-1* (Ellisen et al. 1991; del Amo et al. 1993), *Notch2* (Weinmaster et al. 1992), *Notch3* (Lardelli et al. 1994), and *Notch4/int-3* (Uyttendaele et al. 1996). Dysregulated Notch signaling has been shown to result in several developmental abnormalities and diseases. Chromosomal translocations resulting in the truncation of the human *Notch1* homolog, *TAN1*, have been found in certain T-cell acute lymphoblastic leukemias (T-ALL) (Ellisen et al. 1991). A frequent site for insertional activation by the mouse mammary tumor virus (MMTV) in malignant mammary carcinomas is the *Notch4* locus. The insertion of MMTV results in an aberrantly expressed intracellular form of *Notch4* originally identified as *Int-3* (Gallahan and Callahan 1987). Cerebral autosomal-dominant arteriopathy with subcortical infarcts and leucoencephalopathy (CADASIL) is a disorder that leads to ischaemic strokes and dementia in adults, and has been traced recently to missense mutations of the *Notch3* receptor (Joutel et al. 1997).

Notch ligands are divided into two subclasses, the Delta family and the Serrate family. All Notch ligands share some structural features (Fleming 1998) including epidermal growth factor (EGF)-like repeats, a characteristic DSL domain necessary for Notch binding (Muska-vitch 1994) and a transmembrane region. However, an extracellular cysteine-rich domain and insertions that interrupt some EGF-like repeats are common only to the Serrate family. It is these structural differences that categorize a Notch ligand as a Delta or Serrate family member. Mammalian ligands reported include two members of the Serrate family, *Jagged1/JAG1* (Lindsell et al. 1995; Oda et al. 1997a) and *Jagged2* (Shawber et al. 1996), and two members of the Delta family, murine *Dll1* (Bettenhausen et al. 1995) and murine *Dll3* (Dunwoodie et al. 1997).

Ligand mutations can also result in the disruption of the Notch-signaling pathway, leading to developmental abnormalities. Mutations of human *JAG1* have been attributed to the development of Alagille syndrome (AGS), an autosomal dominant disorder characterized by developmental abnormalities of the heart, skeleton, muscle, liver, and eyes (Li et al. 1997; Oda et al. 1997b). Mice homozygous for the pudgy (*pu*) mutation exhibit severe deformities of the ribs and vertebrae, and this mutation

[Key Words: Notch signaling pathway; endothelium]

⁶Corresponding author.

E-MAIL kstark@amgen.com; FAX (805) 480-1368.

has been mapped to the *Dll3* locus (Kusumi et al. 1998). *Dll1*-deficient mice generated by homologous recombination exhibit neonatal lethality (Hrabe de Angelis et al. 1997), and disruption of murine *Jagged2* revealed an essential role for this gene in limb, craniofacial, and thymic development (Jiang et al. 1998).

The pleiotropic effects of dysregulated Notch signaling validate this pathway as an important regulator of development, and the existence of multiple ligands and receptors suggests a more specialized role for these genes in mammals than in lower eukaryotes. Here we report the molecular cloning and characterization of *Dll4*, the third mammalian member of the Delta family of Notch ligands. The ability to activate Notch receptors and the specific expression pattern of *Dll4* implicate a role for this gene in the regulation of vascular biology.

Results and Discussion

cDNA cloning

An EST database from a murine, white adipose cDNA library was found to contain a clone with homology to known members of the DSL family of Notch ligands. This EST consisted of 450 bp of sequence encoding a DSL domain (Tax et al. 1994) followed by an EGF-like repeat. RACE of double-stranded DNA synthesized from white adipose tissue yielded a 3.4-kb cDNA. Analysis of the insert sequences from multiple clones revealed a 2058-bp ORF which was predicted to encode a 686 amino acid protein (Fig. 1). These clones also contained ~300 bp of 5' untranslated region (UTR) and 1000 bp of 3' UTR, including a polyadenylation signal sequence, and a poly(A) tail (data not shown). The human *DLL4* gene was also isolated and the ORF is ~86% identical at the nucleotide level and 87% identical at the amino acid level to murine *Dll4*.

Dll4 is a novel Delta family member

The predicted protein sequences for both human *DLL4* and murine *Dll4* exhibit all of the hallmarks of the Delta family of Notch ligands. The extracellular region contains 8 EGF-like repeats, as well as a DSL domain that is conserved among all Notch ligands and is necessary for receptor binding. The predicted protein also contains a transmembrane region, and a cytoplasmic tail lacking any catalytic motifs. The absence of an extracellular cysteine-rich domain and the lack of insertions between the EGF-like repeats clearly exclude *Dll4* from the Jagged family of ligands. Sequence comparisons between *Dll4*

and the mu-rine Dll1 and Dll3, and *Xenopus* X-Delta1 (Chitnis et al. 1995) and X-Delta2 (Jen et al. 1997) proteins are shown in Figure 2. The Dll1 and X-Delta1 homologs are >70% identical at the amino acid level, whereas each of these genes is ~50% identical to Dll4. Similarly, Dll4 is ~50% identical to X-Delta2; suggesting *Dll4* is not a mammalian homolog of the *X-Delta2* gene, but rather a novel fourth member of the *Delta* gene family.

Dll4 is expressed in arterial endothelium

Northern blots containing mRNA from a variety of murine tissues were hybridized with a *Dll4* probe. Expression of the 4.3-kb *Dll4* message was highest in lung, followed by heart, kidney, skeletal muscle, and brain (Fig. 3). Transcripts were barely detectable in spleen and testis, but hybridization with GAPDH showed little detectable mRNA in these samples. Northern analysis of whole mouse embryos demonstrated increased expression of *Dll4* from E7 through E17 of development. Analysis by RT-PCR revealed a low level of *Dll4* expression in all tissues examined with higher levels in lung, brown and white adipose, and adrenal (data not shown).

In situ hybridization was performed on a variety of mouse tissues with an antisense *Dll4* riboprobe, and a strikingly specific expression pattern was observed. In both embryonic and adult tissues, the predominant site of *Dll4* expression was the vasculature (Fig. 4). Although it is difficult to rule out expression in the underlying smooth muscle, the expression of *Dll4* in the choroid plexus (Fig. 4a), which lacks smooth muscle cells, suggests that the endothelium is the predominant site of *Dll4* expression. In brain, *Dll4* expression was observed in blood vessels (Fig. 4b), although some non-vascular expression was also seen in neural epithelium (not shown). Non-vascular expression was also observed in adrenal cortex, spleen, and lymph node (not shown). The precise cell types that express *Dll4* in these tissues requires further investigation. Expression of *Dll4* was observed in the glomeruli of the kidney (Fig. 4c) and the vasculature of the lung (Fig. 4d). Within the vasculature, *Dll4* is relatively restricted to arteries, arterioles, and capillaries. This is illustrated strikingly in the umbilical cord (Fig. 4e), in which high expression is observed within the umbilical artery but not in the adjacent vein. The arterial expression of *Dll4* is further illustrated by whole-mount in situ analysis of developing embryos (see the accompanying paper by Krebs et al.). In situ analysis

	X-Delta2	M-Dll1	M-Dll3	M-Dll4
X-Delta1	51%	74%	34%	50%
X-Delta2	-	51%	41%	52%
M-Dll1	-	-	36%	51%
M-Dll3	-	-	-	31%

Figure 2. Comparison of amino acid identity between vertebrate *Delta* genes reveals *Dll4* is a novel family member. Murine Dll4 is ~30%–50% identical to known *Delta* genes, whereas homologs (i.e., X-Delta1 and murine Dll1) are >70% identical. Dll3 remains the most divergent family member.

Figure 3. Murine *Dll4* is expressed in most tissues. Northern analysis shows a transcript of ~4.3 kb in several adult tissues, with lung expressing the highest levels. Expression is also seen in embryos, with increasing levels from embryonic day 9 through day 17. GAPDH hybridization provides for variations in loading and helps explain the lack of signal in spleen and testis.

of human *DLL4* was also performed, and a similar expression pattern was seen (data not shown).

Chromosomal localization

To determine whether the human or mouse gene mapped near any human disease syndromes or pre-exist-

Figure 4. Murine *Dll4* is highly specific for arterial endothelial cells. In situ analysis was performed on murine tissues with an antisense murine *Dll4* riboprobe. *Dll4* expression in (a) choroid plexus; (b) brain; (c) kidney; (d) lung; (e) umbilical cord. The bright field and dark field views are displayed on the left and right, respectively, for each tissue section. Endothelial cells expressing *Dll4* are indicated with arrows. See text for details.

ing mouse mutants, the chromosomal localization of both genes was performed. The human gene was mapped using the Stanford G3 human-hamster radiation hybrid panel, and the mouse gene was localized by FISH. The human *DLL4* gene was located on 15q21.1, and the mouse *Dll4* gene was mapped to chromosome 2E3, a region that shows conservation of synteny with human chromosome 15q. A search of human syndromes (OMIM 1999) and mouse mutants (MGD 1999) that map near these regions did not reveal any phenotypes apparently attributable to the Notch-signaling pathway.

Dll4 activates *Notch1* and *Notch4*

In developing *Xenopus* embryos, activation of the Notch-signaling pathway results in the inhibition of neurogenesis (Coffman et al. 1993), which can be visualized by staining for the presence of N-tubulin. The ability of *Dll4* to activate the *Xenopus* Notch receptor (X-Notch) was examined by injecting *Dll4* RNA into a single blastomere of two-celled albino embryos, which were subsequently labeled for N-tubulin expression at stage 14. Consistent with X-Notch activation, *Dll4* suppresses N-tubulin formation in approximately one-half of the embryos tested when compared with uninjected controls (Fig. 5a,b). To further verify Notch activation by *Dll4*, expression levels of *ESR-1* (Wettstein et al. 1997) and *ESR-7* (Deblandre et al. 1999) in neural ectoderm were quantified in animal cap experiments in the presence or absence of *Dll4* (Fig. 5c, columns 1–6). The *Xenopus* *ESR* genes are related to *Drosophila E(spl)-C*, and are transcriptionally activated by the Notch-signaling pathway. *Dll4* activated endogenous X-Notch, leading to increases in *ESR-1* and *ESR-7* levels. The coexpression of murine *Notch1* with *Dll4* resulted in higher *ESR* levels, similar to those resulting from activation of X-Notch by X-Delta1. This effect is most likely due to a higher affinity between the murine genes, as coexpression of murine *Notch1* with X-Delta1 did not significantly increase *ESR* gene expression. The *Notch4* receptor is highly expressed in the vascular endothelium of both embryos and adults (Uyttendaele et al. 1996). To investigate the potential of *Dll4* as a *Notch4* ligand, we performed animal cap experiments with murine *Notch4* (Fig. 5c, columns 7–10). Similar to the *Notch1* signaling seen previously, the activation of *ESR* gene expression upon *Dll4/Notch4* coinjection was >3.5-fold higher than injection of *Dll4* alone. Although the specificity of *Dll4* for each of the Notch receptors has yet to be elucidated, *Dll4* clearly functions as a ligand for *Notch1* and *Notch4*.

The isolation and characterization of *Dll4* further expands the DSL family of mammalian Notch ligands. The structural characteristics of the protein clearly establish *Dll4* as a novel *Delta* gene, and the expression pattern implicates Notch signaling in the regulation of blood vessel biology. The specific expression of *Dll4* in arterial endothelium is reminiscent of *ephrin-B2*; another membrane-bound ligand (Wang et al. 1998). During vascular development, *ephrin-B2* is expressed in arterial endothelial cells, whereas its cognate receptor, *Eph-B4*, is ex-

Figure 5. *Dll4* acts as a Notch ligand in *Xenopus* embryos. One blastomere of 2-cell albino embryos was injected with either (a) *nLacZ* RNA and 0.1–0.25 ng of *Dll4* RNA, or (b) *nLacZ* RNA only. At stage 14, the embryos were fixed, stained in whole mount with X-gal (light blue reaction product), and then labeled by in situ hybridization for the expression of N-tubulin (dark blue). Embryos are presented anterior to the left with the injected side on the top. Note that N-tubulin expression is suppressed in the area of *Dll4* injection (as visualized with *lacZ*). (c) Results of RNase protection assay to measure the levels of *ESR-1* and *ESR-7*, targets of Notch signaling, in neural ectoderm isolated from embryos injected with the indicated RNAs. Ectoderm was neuralized by injection of *Noggin*. RNAs were extracted at stage 13 and the ability of *Dll4* to activate Notch targets were examined. For quantification, specific band intensities in each lane were normalized to the amount of *EF1-α* RNA to control for sample recovery, and the gene expression results are represented relative to the uninjected controls. Note that *Dll4* weakly activates *ESR-1* and *ESR-7* expression via endogenous X-Notch activation (column 3), whereas *ESR* gene activation by *Dll4* is strongly potentiated by coinjection of murine *Notch1* (column 4) to reach the levels induced by X-Delta1 (column 5). In a separate experiment (columns 7–10), the ability of *Dll4* to signal through the murine *Notch4* receptor was tested. The activation of *ESR-1* and *ESR-7* expression by *Dll4* (column 9) is strongly potentiated by coinjection with murine *Notch4* (column 10). In both experiments, activation of targeted genes was increased >3.5-fold upon coinjection of *Dll4* with either murine *Notch1* or *Notch4*.

pressed only in venous endothelium. Targeted disruption of the murine *ephrin-B2* gene affects angiogenesis, but not vasculogenesis, in developing embryos. Because the Notch-signaling pathway is an important regulator of neurogenesis and myogenesis (Nye and Kopan 1995) a similar involvement in the regulation of vascularization could be envisioned. In fact, the study of receptor mutants and downstream genes is providing evidence that Notch signaling is pivotal in vascular development. The *E(spl)-C* genes are regulated by Notch activation, and a new *E(spl)-C* family member in zebrafish, *gridlock*, is

GENES & DEVELOPMENT 1317

the *EF1- α* probe to control for sample recovery during the assay. Quantification was carried out on a PhosphorImager (Molecular Dynamics), and for each lane, specific band intensities were normalized to the amount of *EF1- α* RNA.

Acknowledgments

We thank Bethany Sutton for DNA sequencing, Tom Gridley, Melissa Graham, Laarni Ulias, Gerry Weinmaster, and Carol Hicks for helpful discussions, and the support of the Amgen Genome Project and the Dll4 Project Team. J.K. was supported in part by NIH grant no. HL62454.

The publication costs of this article were defrayed in part by payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 USC section 1734 solely to indicate this fact.

References

- Artavanis-Tsakonas, S., M.D. Rand, and R.J. Lake. 1999. Notch Signaling: Cell fate control and signal integration in development. *Science* **284**: 770–776.
- Bettenhausen, B., M. Hrabe de Angelis, D. Simon, J.L. Guenet, and A. Gossler. 1995. Transient and restricted expression during mouse embryogenesis of Dll1, a murine gene closely related to Drosophila Delta. *Development* **121**: 2407–2418.
- Chitnis, A., D. Henrique, J. Lewis, D. Ish-Horowicz, and C. Kintner. 1995. Primary neurogenesis in *Xenopus* embryos regulated by a homologue of the Drosophila neurogenic gene Delta. *Nature* **375**: 761–766.
- Coffman, C.R., P. Skoglund, W.A. Harris, and C.R. Kintner. 1993. Expression of an extracellular deletion of Xotch diverts cell fate in *Xenopus* embryos. *Cell* **73**: 659–671.
- Deblandre, G.A., D.A. Wettstein, N. Koyano-Nakagawa, and C. Kintner. 1999. A two-step mechanism generates the spacing pattern of the ciliated cells in the skin of *Xenopus* embryos. *Development* **126**: 4715–4728.
- del Amo, F.F., M. Gendron-Maguire, P.J. Swiatek, N.A. Jenkins, N.G. Copeland, and T. Gridley. 1993. Cloning, analysis, and chromosomal localization of Notch-1, a mouse homolog of Drosophila Notch. *Genomics* **15**: 259–264.
- Dunwoodie, S.L., D. Henrique, S.M. Harrison, and R.S. Beddington. 1997. Mouse Dll3: A novel divergent Delta gene which may complement the function of other Delta homologues during early pattern formation in the mouse embryo. *Development* **124**: 3065–3076.
- Ellisen, L.W., J. Bird, D.C. West, A.L. Soreng, T.C. Reynolds, S.D. Smith, and J. Sklar. 1991. TAN-1, the human homolog of the Drosophila notch gene, is broken by chromosomal translocations in T lymphoblastic neoplasms. *Cell* **66**: 649–661.
- Fleming, R.J. 1998. Structural conservation of Notch Receptors and ligands. *Semin. Cell Dev. Biol.* **9**: 599–607.
- Gallahan, D. and R. Callahan. 1987. Mammary tumorigenesis in feral mice: Identification of a new int locus in mouse mammary tumor virus (Czech II)-induced mammary tumors. *J. Virol.* **61**: 66–74.
- Harland, R.M. 1991. In situ hybridization: An improved whole-mount method for *Xenopus* embryos. *Methods Cell Biol.* **36**: 685–695.
- Hrabe de Angelis, M., J. McIntyre, II, and A. Gossler. 1997. Maintenance of somite borders in mice requires the Delta homologue Dll1. *Nature* **386**: 717–721.
- Jen, W.C., D. Wettstein, D. Turner, A. Chitnis, and C. Kintner. 1997. The Notch ligand, X-Delta-2, mediates segmentation of the paraxial mesoderm in *Xenopus* embryos. *Development* **124**: 1169–1178.
- Jiang, R., Y. Lan, H.D. Chapman, C. Shawber, C.R. Norton, D.V. Serreze, G. Weinmaster, and T. Gridley. 1998. Defects in limb, craniofacial, and thymic development in Jagged2 mutant mice. *Genes & Dev.* **12**: 1046–1057.
- Joutel, A., K. Vahedi, C. Corpechot, A. Troesch, H. Chabriot, C. Vaysiere, C. Cruaud, J. Maciazek, J. Weissenbach, M.G. Bousser et al. 1997. Strong clustering and stereotyped nature of Notch3 mutations in CADASIL patients. *Lancet* **350**: 1511–1515.
- Kintner, C.R. and D.A. Melton. 1987. Expression of *Xenopus* N-CAM RNA in ectoderm is an early response to neural induction. *Development* **99**: 311–325.
- Kintner, C.R. and J. Dodd. 1991. Hensen's node induces neural tissue in *Xenopus* ectoderm. Implications for the action of the organizer in neural induction. *Development* **113**: 1495–1505.
- Knecht, A.K., P.J. Good, I.B. Dawid, and R.M. Harland. 1995. Dorsal-ventral patterning and differentiation of noggin-induced neural tissue in the absence of mesoderm. *Development* **121**: 1927–1935.
- Kopan, R., E.H. Schroeter, H. Weintraub, and J.S. Nye. 1996. Signal transduction by activated mNotch: Importance of proteolytic processing and its regulation by the extracellular domain. *Proc. Natl. Acad. Sci.* **93**: 1683–1688.
- Kusumi, K., E.S. Sun, A.W. Kerrebrock, R.T. Bronson, D.C. Chi, M.S. Bulotsky, J.B. Spencer, B.W. Birren, W.N. Frankel, and E.S. Lander. 1998. The mouse pudgy mutation disrupts Delta homologue Dll3 and initiation of early somite boundaries. *Nat. Genet.* **19**: 274–278.
- Lamb, T.M., A.K. Knecht, W.C. Smith, S.E. Stachel, A.N. Economides, N. Stahl, G.D. Yancopoulos, and R.M. Harland. 1993. Neural induction by the secreted polypeptide noggin. *Science* **262**: 713–718.
- Lardelli, M., J. Dahlstrand, and U. Lendahl. 1994. The novel Notch homologue mouse Notch 3 lacks specific epidermal growth factor-repeats and is expressed in proliferating neuroepithelium. *Mech. Dev.* **46**: 123–136.
- Li, L., I.D. Krantz, Y. Deng, A. Genin, A.B. Banta, C.C. Collins, M. Qi, B.J. Trask, W.L. Kuo, J. Cochran et al. 1997. Alagille syndrome is caused by mutations in human Jagged1, which encodes a ligand for Notch1. *Nat. Genet.* **16**: 243–251.
- Lindsell, C.E., C.J. Shawber, J. Boulter, and G. Weinmaster. 1995. Jagged: A mammalian ligand that activates Notch1. *Cell* **80**: 909–917.
- Mouse Genome Database (MGD), Mouse Genome Informatics. The Jackson Laboratory, Bar Harbor, Maine. World Wide Web (URL: <http://www.informatics.jax.org/>). October, 1999.
- Muskavitch, M.A. 1994. Delta-notch signaling and Drosophila cell fate choice. *Dev. Biol.* **166**: 415–430.
- Nieuwkoop, P.D. and J. Faber. 1967. Normal table of *Xenopus laevis*. Amsterdam, North Holland.
- Nye, J.S. and R. Kopan. 1995. Vertebrate ligands for Notch. *Curr. Biol.* **5**: 966–969.
- Oda, T., A.G. Elkahoul, P.S. Meltzer, and S.C. Chandrasekharappa. 1997a. Identification and cloning of the human homolog (JAG1) of the rat Jagged1 gene from the Alagille syndrome critical region at 20p12. *Genomics* **43**: 376–379.
- Oda, T., A.G. Elkahoul, B.L. Pike, K. Okajima, I.D. Krantz, A. Genin, D.A. Piccoli, P.S. Meltzer, N.B. Spinner, F.S. Collins et al. 1997b. Mutations in the human Jagged1 gene are responsible for Alagille syndrome. *Nat. Genet.* **16**: 235–242.
- Online Mendelian Inheritance in Man, OMIM (TM). Center for Medical Genetics, Johns Hopkins University (Baltimore, MD) and National Center for Biotechnology Information, National Library of Medicine (Bethesda, MD). World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>. 1999.
- Oschwald, R., K. Richter, and H. Grunz. 1991. Localization of a nervous system-specific class II beta-tubulin gene in *Xenopus laevis* embryos by whole-mount in situ hybridization. *Int. J. Int. Biol.* **35**: 399–405.
- Shawber, C., J. Boulter, C.E. Lindsell, and G. Weinmaster. 1996. Jagged2: A serrate-like gene expressed during rat embryogenesis. *Dev. Biol.* **180**: 370–376.
- Struhl, G., K. Fitzgerald, and I. Greenwald. 1993. Intrinsic activity of the Lin-12 and Notch intracellular domains in vivo. *Cell* **74**: 331–345.
- Tax, F.E., J.J. Yeagers, and J.H. Thomas. 1994. Sequence of *C. elegans* lag-2 reveals a cell-signalling domain shared with Delta and Serrate of Drosophila. *Nature* **368**: 150–154.
- Turner, D.L. and H. Weintraub. 1994. Expression of achaete-scute homologue 3 in *Xenopus* embryos converts ectodermal cells to a neural fate. *Genes & Dev.* **8**: 1434–1447.
- Uyttendaele, H., G. Marazzi, G. Wu, Q. Yan, D. Sassoon, and J. Kitajewski. 1996. Notch4/int-3, a mammary proto-oncogene, is an endothelial cell-specific mammalian Notch gene. *Development* **122**: 2251–2259.
- Wang, H.U., Z.-F. Chen, and D.J. Anderson. 1998. Molecular distinction and angiogenic interaction between embryonic arteries and veins revealed by ephrin-B2 and its receptor Eph-B4. *Cell* **93**: 741–753.
- Weinmaster, G., V.J. Roberts, and G. Lemke. 1992. Notch2: A second mammalian Notch gene. *Development* **116**: 931–941.
- Wettstein, D.A., D.L. Turner, and C. Kintner. 1997. The *Xenopus* homologue of Drosophila Suppressor of Hairless mediates Notch signaling during primary neurogenesis. *Development* **124**: 693–702.
- Zhong, T.P., M. Rosenberg, M.-A.P.K. Mohideen, B. Weinstein, and M. Fishman. 2000. gridlock, an HLH gene required for assembly of the aorta in Zebrafish. *Science* **287**: 1820–1824.

***Dll4*, a novel Notch ligand expressed in arterial endothelium**

John R. Shutter, Sheila Scully, Wei Fan, et al.

Genes Dev. 2000, **14**:

Access the most recent version at doi:[10.1101/gad.14.11.1313](https://doi.org/10.1101/gad.14.11.1313)

References

This article cites 37 articles, 17 of which can be accessed free at:
<http://genesdev.cshlp.org/content/14/11/1313.full.html#ref-list-1>

License

Email Alerting Service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).