p53-independent functions of the p19^{ARF} tumor suppressor

Jason D. Weber,^{1,5} John R. Jeffers,² Jerold E. Rehg,³ David H. Randle,^{1,4} Guillermina Lozano,⁶ Martine F. Roussel,^{1,4} Charles J. Sherr,^{1,4,5} and Gerard P. Zambetti^{2,4,7}

Departments of ¹Tumor Cell Biology, ²Biochemistry, and ³Pathology, St. Jude Children's Research Hospital, Memphis, Tennessee 38105, USA; ⁴Department of Biochemistry, University of Tennessee, Memphis, Tennessee 38163, USA;

⁵Howard Hughes Medical Institute, St. Jude Children's Research Hospital, Memphis, Tennessee 38105, USA;

⁶Department of Molecular Genetics, The University of Texas M.D. Anderson Cancer Center, Houston, Texas 77030, USA

The p19^{ARF} tumor suppressor antagonizes Mdm2 to induce p53-dependent cell cycle arrest. Individual TKO (triple knock out) mice nullizygous for *ARF*, *p53*, and *Mdm2* develop multiple tumors at a frequency greater than those observed in animals lacking both *p53* and *Mdm2* or *p53* alone, demonstrating that p19^{ARF} can act independently of the Mdm2-p53 axis in tumor surveillance. Reintroduction of *ARF* into TKO mouse embryo fibroblasts (MEFs), but not into those lacking both *p53* and *ARF*, arrested the cell division cycle in the G1 phase. Inhibition of the retinoblastoma protein had no effect on the ability of ARF to arrest TKO MEFs. Thus, in the absence of Mdm2, p19^{ARF} interacts with other targets to inhibit cell proliferation.

[Key Words: ARF; p53; Mdm2; DMP1; tumor suppression]

Received June 14, 2000; revised version accepted August 1, 2000.

Two proteins encoded by the *INK4a/ARF* locus, p16^{INK4a} and p19^{ARF}, functionally interact with the retinoblastoma protein (RB) and p53 tumor suppressors, respectively. These four proteins are among the most frequently disrupted genes in human cancer. The p53 monomer forms a labile homotetrameric transcription factor that is activated in response to DNA damage and inappropriate mitogenic signaling (Ko and Prives 1996). Among its transcriptional targets are p21^{Cip1}, an inhibitor of cyclin-dependent kinases, and Mdm2, a potent negative regulator of p53. Mdm2 is a multifunctional protein that inhibits p53 activity in several ways: It binds to the p53 transactivation domain to antagonize its activity on DNA (Momand et al. 1992; Oliner et al. 1993), and it fosters p53 degradation (Haupt et al. 1997; Kubbutat et al. 1997), acting as an E3 ligase that directly ubiquitinates p53 (Honda et al. 1997) and enhancing its shuttling to cytoplasmic proteasomes (Roth et al. 1998). In response to cellular stress, stabilization of p53 occurs when its association with Mdm2 is disrupted by post-translational modifications of the proteins (Giaccia and Kastan 1998) or by interference of Mdm2 function by p19^{ARF} (Kamijo et al. 1998; Pomerantz et al. 1998; Stott et al. 1998; Zhang et al. 1998).

The *ARF* tumor suppressor acts as a sensor of hyperproliferative signals emanating from oncoproteins and inducers of S-phase entry, such as Myc, E1A, Ras, and E2F-1 (Bates et al. 1998; De Stanchina et al. 1998; Palm-

ero et al. 1998; Zindy et al. 1998). In turn, p19^{ARF} triggers p53-dependent growth arrest in the G1 and G2 phases of the cell cycle or, in the presence of appropriate collateral signals, sensitizes cells to apoptosis (Quelle et al. 1995; Kamijo et al. 1997; De Stanchina et al. 1998; Zindy et al. 1998; Eischen et al. 1999; Jacobs et al. 1999; Schmitt et al. 1999). ARF binds directly to Mdm2, sequestering it in the nucleolus and enabling transcriptionally active p53 to accumulate in the nucleoplasm (Tao and Levine 1999; Weber et al. 1999). ARF also limits the ability of Mdm2 to ubiquitinate p53 in vitro (Honda and Yasuda 1999; Midgley et al. 2000) and in vivo (Xirodimas and Lane, pers. comm.). Binding of ARF to Mdm2 involves two separate domains of both the mouse p19^{ARF} and human p14^{ARF} proteins that interact cooperatively with a central acidic segment of Mdm2 (Lohrum et al. 2000a; Weber et al. 2000). Although ARF localizes to the nucleolus on its own, ARF binding unmasks a cryptic nucleolar localization signal within the carboxy-terminal Mdm2 RING domain, the integrity of which is required for localization of the binary complex to the nucleolus (Lohrum et al. 2000b; Weber et al. 2000). Disruption of the Mdm2 nucleolar localization signal enables Mdm2 to retain ARF in the nucleoplasm. Therefore, the ARF-Mdm2 interaction appears to be bi-directional in that each protein can regulate the subnuclear localization of the other.

Emerging evidence suggests that the ARF-p53-Mdm2 pathway is not strictly linear. Mice engineered to overexpress a *c-Myc* transgene under the control of the immunoglobulin heavy chain enhancer (E_H) develop pre-B and B-cell lymphomas, with a majority of the resulting

⁷Corresponding author.

E-MAIL gerard.zambetti@stjude.org; FAX (901) 525-8025.

Article and publication are at www.genesdev.org/cgi/doi/10.1101/gad.827300.

tumors sustaining *ARF* deletion, *p53* mutation, or *Mdm2* overexpression (Eischen et al. 1999). However, several tumors that lacked p19^{ARF} or *p53* function overexpressed *Mdm2*, allowing argument against a simple epistatic relationship between the genes. Moreover, pre-malignant B cells expressing Eμ-Myc and lacking both *ARF* and *p53* proliferated at a faster rate in vitro than cells lacking *ARF* or *p53* alone, implying that p19^{ARF} may functionally interact with proteins other than *p53*. To address this issue, we generated mice nullizygous for *ARF*, *p53*, and *Mdm2* and studied the effects of reintroduced ARF in cells lacking its only known target.

Results and Discussion

ARF and *p53* collaborate in tumor surveillance

Both *p53*-null and *ARF*-null mice are highly tumor-prone with mean latencies for survival of 19 and 32 weeks, respectively (Donehower et al. 1992; Kamijo et al. 1999). In mice lacking *p53*, T-cell lymphomas predominate (~70%), with the remainder being sarcomas (Jacks et al. 1994). In contrast, *ARF*-null animals have an appreciably lower incidence of lymphomas (25%) and primarily develop poorly differentiated sarcomas (50%), with carcinomas and gliomas appearing in the others (Kamijo et al. 1997, 1999). These results are compatible with the idea that ARF acts upstream from *p53* and is selectively activated in response to only a subset of stress signals that affect *p53* function (Sherr 1998). The increased latency for tumor formation observed in *ARF*-null versus *p53*-null animals may shift the tumor spectrum away from rapidly lethal lymphomas, allowing the emergence of tumor types not normally seen in *p53*-null mice.

We generated mice deficient for *ARF/p53* and *ARF/p53/Mdm2* (designated triple knock-out, or TKO, mice) in an attempt to determine whether *ARF* and *p53* might synergize in tumor suppression, either in the presence or absence of *Mdm2*. We observed no differences in the mean latencies of tumor formation (~16 weeks) in TKO mice or *ARF/p53*- or *p53/Mdm2*-deficient mice, indicating that combinatorial loss of *ARF* and *p53* had little or no effect on survival as compared with animals lacking *p53* alone (Fig. 1). Disruption of *Mdm2* leads to embryonic lethality, which is rescued on a *p53*-null background (Jones et al. 1995; Montes de Oca Luna et al. 1995). The spectrum of tumors seen in mice lacking both *p53* and *Mdm2* is similar to that of mice lacking *p53* alone (McDonnell et al. 1999). In contrast, mice lacking both *ARF* and *p53*, with or without *Mdm2*, showed a wider range of tumor types than animals lacking either gene alone, and many developed multiple primary tumors (Table 1). *ARF*-null animals rarely show hemangiosarcomas and osteogenic sarcomas, but we have never observed teratomas, rhabdomyosarcomas, hepatocellular carcinomas, intestinal adenocarcinomas, or renal cell carcinomas in these animals throughout three years of continuous observation. Four of the 19 TKO mice developed primary tumors arising from three distinct cell types.

Although we cannot rigorously exclude the contribu-

Figure 1. Survival of mice lacking *ARF*, *p53*, and/or *Mdm2*. *p53/Mdm2*-null (○), *ARF/p53*-null (dotted line), *ARF/p53*-null *Mdm2*^{+/-} (●), and *ARF/p53/Mdm2*-null (solid line) mice were observed for 33 weeks. Tumors arising in *ARF/p53*-null and *ARF/p53/Mdm2*-null animals are listed in Table 1.

tion of strain-specific modifiers on the emergence of different tumor types, all parental mice used in our study were originally derived from strain 129/SvJ ES cells inoculated into C57BL/6 blastocysts, generating chimeric animals that were backcrossed onto a C57BL/6 background. Therefore, the emergence of tumors in *ARF/p53*-null and TKO mice not normally observed in *ARF*-null, *p53*-null, or *p53/Mdm2* double-null animals, taken together with the similar latencies for tumor development, suggests that *ARF* and *p53* loss contribute independently to some aspects of tumor suppression.

Mdm2- and *p53*-independent functions of ARF

Early-passage mouse embryo fibroblasts (MEFs) explanted from *p53*-null embryos or cell lines that sustained *p53* mutations during establishment are resistant to ARF-induced arrest, underscoring a requirement of functional *p53* for ARF activity (Kamijo et al. 1997, 1998; Pomerantz et al. 1998; Stott et al. 1998; Zhang et al. 1998). In agreement with these findings, MEFs lacking both *ARF* and *p53* were resistant to ARF overexpression, whereas 90% of wild-type MEFs arrested within 48 hours of infection with a p19^{ARF}-expressing retrovirus (Fig. 2A). Unexpectedly, >90% of *p53/Mdm2*-null or *ARF/p53/Mdm2*-null MEFs failed to incorporate 5-bromodeoxyuridine (BrdU) after ARF was introduced (Fig. 2A); however, the kinetics of ARF-induced arrest were significantly delayed.

Cells lacking *p53* express unusually high levels of endogenous p19^{ARF} (Fig 2C, lane 1) (Quelle et al. 1995), which can be suppressed by reintroduction of wild-type *p53* (Kamijo et al. 1998; Stott et al. 1998). Surprisingly, endogenous p19^{ARF} was expressed at much lower levels in MEFs lacking both *p53* and *Mdm2* (Fig. 2C, lane 2; Fig. 2D, panel 3), implying that *Mdm2* somehow regulates the *p53*-ARF feedback loop. Use of cells lacking *p53* and *Mdm2* therefore provided an opportunity to determine

Table 1. Tumor spectrum in ARF/p53-null and ARF/p53/Mdm2-null mice

Genotype	Tumor type ^a	Number
ARF/p53-null	Single tumors	13 (72%)
	Lymphoma	9
	Meningioma	1
	Osteogenic sarcoma	1
	Teratoma	1
	Spindle cell carcinoma (skin)	1
	Multiple primary tumors	5 (28%)
	Lymphoma	
	+ Osteogenic sarcoma	1
	+ Hepatocellular carcinoma	1
	+ Hemangiosarcoma	1
	+ Fibrosarcoma	1
ARF/p53/ Mdm2-null	Single tumors	10 (53%)
	Lymphoma	5
	Hemangiosarcoma	2
	Nerve sheath sarcoma	1
	Rhabdomyosarcoma	1
	Osteosarcoma	1
	Multiple primary tumors	9 (47%)
	Undifferentiated sarcoma	
	+ Intestinal adenocarcinoma	1
	Lymphoma	
	+ Rhabdomyosarcoma ^b	2
	+ Intestinal adenocarcinoma	1
	+ Dermal sarcoma, unknown type	1
	+ Hemangiosarcoma	1
	+ Osteogenic sarcoma	1
	+ Skin basal cell carcinoma	1
	+ Intestinal adenocarcinoma	1

^aMice with multiple tumors are grouped under the subheading of one of the primary tumors followed by the diagnosis of the second or third tumor types (indicated by + signs and hierarchical indentation).

^bLymphomas were of the widely disseminated T-cell type except for one mouse that exhibited lymphoma of distinct B- and T-cell types plus rhabdomyosarcoma.

the effects of ARF protein induction on cell cycle progression in response to DMP1, a transcriptional ARF activator (Inoue et al. 1999). Enforced DMP1 expression induced p19^{ARF} to levels that were significantly lower than those observed in p53-null cells (Fig. 2C). Induced ARF was readily detected within the nucleoli of wild-type (not shown) and p53/Mdm2-null MEFs (Fig. 2D, panels 6 and 12). In turn, DMP1 inhibited BrdU incorporation in both cell types by 96 hours postinfection (Fig 2B; Fig 2D, panel 11). In contrast, DMP1 was without effect in ARF/p53-null and TKO MEFs (Fig. 2B), underscoring the dependence of DMP1 on ARF in arresting proliferation (Inoue et al. 1999).

The amino-terminal 14 amino acids of mouse and human ARF are the most conserved residues in the polypeptide and comprise one of two binding sites for Mdm2 (Lohrum et al. 2000a; Weber et al. 2000). In mouse p19^{ARF}, the second Mdm2 interaction domain resides within residues 26–37, the integrity of which is also re-

quired for ARF nucleolar localization (Weber et al. 1999). Hence, a p19^{ARF} mutant lacking residues 1–14 (ARF Δ1–14) retains its ability to correctly localize to nucleoli but is unable to sequester Mdm2 and does not activate p53-dependent growth arrest (Weber et al. 2000). To determine whether the same protein interaction domain of ARF was required to arrest proliferation of MEFs lacking Mdm2, we introduced ARF Δ1–14 into TKO cells by retroviral gene transfer. In the experiment shown, the level of ARF Δ1–14 expressed in infected TKO MEFs was higher than that of the full-length ARF protein (Fig. 3A). Both wild-type and mutant proteins localized to nucleoli (Fig. 3B, panels 9, 12, 15, 18), which were independently demarcated by fibrillarin staining (Weber et al. 1999 and data not shown).

TKO cells infected with retroviruses encoding the T-cell coreceptor CD8, wild-type ARF, or ARF Δ1–14 were analyzed for their ability to enter S phase. BrdU was added to the cultures 24 hours before fixation, and cells were stained and examined 48 and 96 hours after infection. Wild-type ARF inhibited almost all BrdU incorporation by 96 hours post-infection (Fig. 3B, panel 11), whereas ARF Δ1–14 was ineffective at preventing S phase entry (Fig. 3B, panels 14 and 17). Analysis of DNA content by flow cytometry revealed that a significant population of TKO MEFs underwent arrest in G1 phase following infection for 96 hours with the retrovirus encoding wild-type ARF (Fig. 3C). In contrast, TKO MEFs infected with viruses encoding control CD8 or ARF Δ1–14 remained distributed throughout the cell cycle. Like MEFs lacking p53 alone, a percentage of the TKO cells were polyploid, yielding a G2/M population with an 8N DNA content. Introduction of wild-type ARF, but not ARF Δ1–14, also reduced the fraction of 8N cells (Fig. 3C), consistent with the ability of ARF to restrict S phase entry. ARF may also sensitize TKO cells to apoptosis, albeit to a significantly lesser extent than p53 (data not shown), as indicated by an increased level of cells with DNA content less than 2N.

To determine whether the effects of ARF were persistent, infected TKO MEFs were cultured at low density and refed every three days. Emerging colonies were stained with Wright-Giemsa solution 12 days postinfection, revealing a significant reduction in colony formation in ARF-infected TKO MEFs but very little effect after introduction of ARF Δ1–14 as compared with CD8 (Fig. 3D). The few colonies that arose from TKO cells infected with wild-type ARF failed to express p19^{ARF} protein as determined by indirect immunofluorescence (data not shown), indicating that these were likely derived from uninfected cells. Although ARF Δ1–14 is severely impaired in its ability to arrest wild-type and TKO MEFs, it does retain some activity. We reproducibly saw a small effect on DNA replication (15% inhibition of BrdU incorporation) and a reduction in the number and size of colonies in TKO MEFs (Fig. 3D), which can likely be attributed to the retained functional ARF domains (Weber et al. 1999, 2000).

Several conclusions emerge from these experiments: (1) ARF can arrest cells lacking both p53 and Mdm2, but

Figure 2. ARF and DMP1 arrest mouse embryo fibroblasts (MEFs) lacking both *p53* and *Mdm2*. (A) Primary MEFs from mice with the indicated genotypes (*top*) were infected with retroviruses encoding CD8 (□) or p19^{ARF} (■). Cells labeled with 5-bromodeoxyuridine (BrdU) were analyzed by indirect immunofluorescence (green) and simultaneously scored for p19^{ARF} expression (red) using 3 coverslips and >100 cells per coverslip in 4 independent experiments. (B) Primary MEFs of indicated genotypes (*bottom*) were infected with retroviruses encoding CD8 (dark bars) or DMP1 (open bars). Cells labeled for 24 hours with BrdU were stained 96 hours postinfection (green) and scored simultaneously for DMP1 expression (red). Error bars indicate standard deviations. (C) Expression of p19^{ARF} protein in MEFs (3T3 strains, each at passage 9) of the indicated genotypes (*top*). Cells null for both *p53* and *Mdm2* (middle two lanes) were infected with retroviruses encoding either CD8 or DMP1 as indicated and assayed 96 hours later for ARF protein by direct immunoblotting. Equal quantities of protein (150 µg) were loaded per lane. (D) *p53/Mdm2*-null MEFs infected with retroviruses encoding CD8 (panels 1–3, 7–9) or DMP1 (panels 4–6, 10–12) were analyzed by immunofluorescence for BrdU incorporation and ARF induction 48 and 96 hours after infection.

not p53 alone. This implies that Mdm2 antagonizes some activity of ARF on targets other than p53; in the absence of Mdm2 this other activity of ARF is revealed. (2) In principle, the ability of overexpressed ARF to accumulate in the nucleolus might nonspecifically interfere with some aspect of nucleolar trafficking required for cell cycle progression. However, nucleolar localization of ARF is not sufficient to induce cell cycle arrest in the absence of both Mdm2 and p53, because ARF Δ 1–14 gains access to this compartment but is impaired in halting proliferation. (3) Conversely, the amino-terminal domain of p19^{ARF}, which binds to Mdm2, also appears to be important in ensuring the interaction of ARF with another target required for S phase entry. (4) Finally, although ARF arrests cells containing wild-type Mdm2 and p53 in both G1 and G2, its preferential effects at the G1/S boundary in TKO cells imply that the relevant ARF target protein is required only for G1 exit.

ARF functions are independent of p21^{Cip1} and RB

Proteins that can interfere with the G1/S transition include the CDK inhibitor p21^{Cip1}. Induction of p21^{Cip1} was evident after infection of TKO MEFs with a retrovirus-encoding wild-type p53 (Fig. 4A), although the infected cells rapidly underwent apoptosis rather than G1-phase arrest (data not shown). Introduction of ARF into TKO MEFs failed to induce p21^{Cip1} 96 hours after infection, in agreement with previous observations with *p53*-null cells (Kamijo et al. 1997). Moreover, induction of p21^{Cip1} is not obligatory for ARF-induced arrest in cells expressing wild-type p53, because MEFs derived from *p21^{Cip1}*-null or *p21^{Cip1}/p27^{Kip1}* double-null mice arrest on ARF overexpression (Groth et al. 2000).

Two p53-related proteins, p63 and p73, share significant homology with p53 within their N-terminal transactivation and central DNA-binding domains (Lohrum

Figure 3. p53-independent arrest requires the conserved ARF amino-terminal domain. (A) Triple knock out (TKO) mouse embryo fibroblasts (MEFs) infected with retroviruses encoding CD8, ARF, or ARF Δ1-14 were lysed 96 hours after infection, and separated proteins were blotted with affinity-purified polyclonal rabbit antibody to p19^{ARF}. (B) TKO MEFs infected with retroviruses encoding CD8 (panels 1–6), ARF (panels 7–12), or ARF Δ1-14 (panels 13–18) were labeled with 5-bromodeoxyuridine (BrdU) for 24 hours and analyzed 48 and 96 hours postinfection for incorporated BrdU (green) and p19^{ARF} expression (red). DNA was visualized with Hoechst yellow S769121 dye. (C) TKO MEFs (5×10^4) were infected with CD8, ARF, and ARF Δ1-14 retroviruses, stained with propidium iodide 96 hours after infection, and subjected to fluorescence-activated cell sorting to determine DNA content. (D) TKO MEFs were seeded at 300 cells per dish and infected with retroviruses encoding CD8, ARF, or ARF Δ1-14. Colonies were stained 12 days post-infection.

and Vousden 2000). Although Mdm2 can block transactivation by all three proteins, only p53 is targeted by Mdm2 for degradation (Zeng et al. 1999). Because p63 and p73 are capable of inducing p53-responsive genes such as p21^{Cip1} in response to various signals (Kaelin 1999), our findings also imply that ARF does not access p21^{Cip1} through the agency of these p53 family members.

Carnero et al. (2000) provided evidence that ARF might act independently of p53 in MEFs retaining Mdm2 function. Expression of ARF antisense constructs relieved the ability of endogenous p19^{ARF} to induce replicative senescence, whereas subsequent excision of the integrated antisense vector with Cre-recombinase restored the growth suppressive function of ARF even in the presence of a dominant-negative p53 mutant. Additionally, expression of SV40 large T antigen overcame the arrest induced by restored ARF function, suggesting that RB might be targeted by ARF through a p53-independent, but possibly Mdm2-dependent, pathway. Although these results disagree with several published reports that ARF acts upstream from p53 and does not depend on the RB pathway to induce arrest (Quelle et al. 1995; Pomerantz et al. 1998; Stott et al. 1998; Groth et al.

2000), the protocol for reversible antisense suppression is distinctly different from others' use of cells lacking the genes in question.

To explore a possible role for RB in our experimental system, we generated a population of TKO MEFs infected with a retrovirus expressing the green fluorescent protein (GFP) and encoding the human papilloma virus (HPV) E7 protein from an internal ribosomal entry sequence. Expression of E7 protein was documented by immunoblotting (Fig. 4B), and cells overexpressing E7 and GFP or GFP alone were infected with retroviral vectors encoding CD8 or ARF. Three days later, cells pulsed for 24 more hours with BrdU were fixed and analyzed for BrdU incorporation. To control for the effectiveness of E7 in inhibiting RB function, we also infected these cells with a retrovirus encoding p19^{INK4d}, a potent inhibitor of cyclin D-dependent kinases in which the ability to induce G1 phase arrest is RB-dependent (Hirai et al. 1995). Overexpression of E7 had no effect on BrdU incorporation in TKO MEFs but greatly increased the number of cells replicating DNA in the presence of p19^{INK4d} (Fig. 4C). Contrary to its ability to block p19^{INK4d}-induced arrest, E7 was unable to bypass the growth-inhibitory affects of p19^{ARF} (Fig. 4C). Loss of RB function thus ap-

Figure 4. ARF-induced arrest of triple knock out (TKO) cells does not depend on p21^{Cip1} and is not overridden by HPV E7. (A) TKO mouse embryo fibroblasts (MEFs) infected with retroviruses encoding CD8 or ARF (96 hours postinfection) or wild-type p53 (48 hours postinfection) were lysed, and separated proteins were blotted with antibodies to p53, p21^{Cip1}, and ARF. (B) TKO MEFs infected with retroviruses encoding E7 or CD8 were lysed 48 hours after infection, and separated proteins were blotted with antibody to E7. (C) TKO MEFs infected with MSCV-GFP-IRES or MSCV-GFP-IRES-E7 retroviruses were re-plated 48 hours later and infected with retroviruses encoding CD8, ARF, or p19^{IN4d}. Infected cells were analyzed 96 hours postinfection using an FITC filter to detect GFP-IRES-E7 infected cells (green), immunofluorescence to score 5-bromodeoxyuridine incorporation (red), and antibody to p19^{ARF} followed by Alexa 350 stain to visualize nucleolar ARF (blue). Alternatively, p19^{IN4d} was detected using a rabbit anti-p19^{IN4d} antibody followed by biotinylated anti-rabbit immunoglobulin and streptavidin Texas red to ensure >90% infection efficiency of TKO and E7-containing TKO cells.

pears to have no effect on the kinetics or properties of ARF arrest in the absence of both Mdm2 and p53. Moreover, the failure of E7 to override the ARF-induced G1 block implies that ARF prevents deregulated E2Fs from inducing S phase entry.

The ARF-Mdm2-p53 network

ARF responds to oncogenic signaling by neutralizing functions of Mdm2, thus allowing p53 to accumulate in a transcriptionally active form in the nucleoplasm. In cells that retain Mdm2 but lack p53 function, ARF loses its ability to rapidly halt cell cycle progression in the G1 and G2 phases of the division cycle, indicating that the

response of cells to ARF induction is normally p53-dependent. Although cells lacking p53 alone or both p53 and ARF are resistant to ARF-induced arrest, cells lacking p53 and Mdm2 or all three genes are surprisingly more sensitive. In the latter cell types, ARF-induced arrest occurs more slowly than in wild-type MEFs, and the cells accumulate in G1, but not G2. Therefore, when present, Mdm2 antagonizes the ability of ARF to induce cell cycle arrest through targets other than p53.

ARF might interfere with the shuttling of other proteins whose nuclear export is required for cell cycle progression, and these interactions are likely to depend on the conserved ARF amino-terminal domain (a.a. 1–14) required for Mdm2 binding. Although ARF binds to Mdm2 with high affinity (Weber et al. 2000), the amounts of Mdm2 expressed in resistant p53-null or ARF/p53 double-null MEFs are negligible when compared with the unusually high levels of endogenous p19^{ARF} protein or those achieved through ectopic ARF expression. It therefore seems unlikely that Mdm2 directly interacts with p19^{ARF} to inhibit its ability to access targets other than p53. Instead, the loss of Mdm2 seems to reset the cell's sensitivity to ARF through some other mechanism.

ARF steadily accumulates as wild-type MEFs are progressively passaged in culture and approach senescence (Zindy et al. 1998), whereas ARF-null MEFs maintain their proliferative capacity and appear to be immortal (Kamijo et al. 1997). Although the ability of ARF to induce arrest in cells lacking both p53 and Mdm2 raises the possibility that replicative limits might be reimposed by Mdm2 loss, these double-null cells, like those lacking p53 alone, do not senesce (McMasters et al. 1996). This may be due to a reduction in the p19^{ARF} level in Mdm2-deficient cells as compared with those in cells lacking only p53 (Fig. 2C).

Many lines of evidence now link ARF-Mdm2-p53 functions through a complex network rather than a simple linear pathway. It has long been appreciated that Mdm2 is a p53-responsive gene (Barak et al. 1993; Wu et al. 1993). Conversely, proliferating cells that lack p53 function alone synthesize relatively low basal levels of Mdm2 but, for still unexplained reasons, gratuitously express supra-physiologic levels of ARF (Quelle et al. 1995) that can again be suppressed by reintroduction of wild-type p53 (Kamijo et al. 1998; Stott et al. 1998). On the other hand, cells lacking both p53 and Mdm2 did not express high levels of p19^{ARF} (Fig. 2C,D). Considering that the one known biochemical function of Mdm2 is to facilitate protein degradation (Honda et al. 1997), we might imagine, for example, that Mdm2 could catalyze the destruction of an ARF negative regulator, such as Bmi-1 or Twist (Jacobs et al. 1999; Maestro et al. 1999). Whatever the mechanism, it seems evident that this regulatory network must depend on other proteins, the actions of which could in part compensate for the missing gene products in ways that we currently do not understand. Nonetheless, at least one ARF target, when regulated in an entirely Mdm2- and p53-independent manner, can be rate limiting for the G1/S transition.

Materials and methods

Interbreeding of mice and tumor analysis

Homozygous *Mdm2/p53*-null mice (129/Sv × C57BL/6J background) (Montes de Oca Luna et al. 1995) were intercrossed with *ARF*-null mice (129/Sv × C57BL/6J background) (Kamijo et al. 1997) to generate double-null (*p53/mdm2* and *p53/ARF*) and TKO mice. Genotypes were determined by polymerase chain reaction (PCR)-based assays as above. Moribund mice were killed and necropsied, and tumor tissues fixed in 10% formalin were embedded in paraffin for histopathological analyses (Kamijo et al. 1999).

Cell culture and virus infection

Primary MEFs were maintained in Dulbecco's-modified Eagle's medium with 10% fetal bovine serum, 2 mM glutamine, 0.1 mM non-essential amino acids, 55 μ M β -mercaptoethanol, and 10 μ g/ml gentamycin (GIBCO/BRL, Gaithersburg, MD). Virus production and infection of cells were performed using retroviral helper and vector plasmids (Roussel et al. 1995) provided by Charles Sawyers (University of California, Los Angeles). Vectors encoding p19^{ARF} and p19^{INK4a} Δ 1–14 were used as previously described (Weber et al. 2000). p19^{INK4a} cDNA was subcloned into pSR α -MSV-tkCD8 (Hirai et al. 1995), and cDNA for the papilloma virus E7 protein (Slebos et al. 1994) was subcloned into MSCV-IRES-GFP (Hawley et al. 1994).

Immunofluorescence

Primary MEFs from *ARF/p53*-null, *p53/Mdm2*-null, and *ARF/p53/Mdm2*-null backgrounds seeded onto glass coverslips (3×10^4 cells per slip) were infected with retroviruses encoding CD8, ARF, ARF Δ 1–14, or p19^{INK4a}. Conversely, *ARF/p53/Mdm2*-null MEFs (1×10^6) were infected with retroviruses encoding E7, replated 48 hours later (3×10^4 cells per glass coverslip), and infected with retroviruses encoding CD8, ARF, or p19^{INK4a}. BrdU (10 μ M) was added to the culture medium 24 hours before fixation. Cells were fixed in methanol/acetone (volume ratio, 1:1) and stained for 1 hr with affinity-purified rabbit anti-p19^{ARF} antibody (Quelle et al. 1995) or rabbit anti-p19^{INK4a} (Hirai et al. 1995), followed by 30-minute exposure to biotinylated anti-rabbit immunoglobulin and streptavidin-conjugated Texas red (both from Amersham, Arlington Heights, IL). Cells were treated for 10 minutes with 1.5 N HCl and stained for 1 hr with mouse monoclonal antibody to BrdU (Amersham), followed by fluorescein isothiocyanate (FITC)-conjugated anti-mouse immunoglobulin (Amersham). DNA was visualized with Hoechst yellow S769121 dye (Molecular Probes, Eugene, OR).

Long-term colony assay

ARF/p53/Mdm2-null primary mouse embryo fibroblasts were seeded (3×10^2) onto 100-mm diameter dishes and infected with CD8, ARF, or ARF Δ 1–14 retroviruses. Spent medium was replenished every three days, and emerging colonies were stained with Wright-Giemsa (Biochemical Sciences, Swedesboro, NJ) 12 days postinfection.

Immunoblotting

Frozen cell pellets were lysed in ice-cold Tween-20 lysis buffer (50 mM Hepes, pH 7.5, 150 mM NaCl, 1 mM ethylenediamine-tetraacetic acid, 2.5 mM ethyleneglycoltetraacetic acid, 0.1% Tween-20, 1 mM phenylmethylsulfonyl fluoride, 0.4 units/ml aprotinin, 10 μ g/ml pepstatin, and 10 μ g/ml leupeptin) and soni-

cated. Samples (200 μ g protein per lane) were separated by electrophoresis on denaturing polyacrylamide gels containing sodium dodecyl sulfate and were transferred to Immobilon polyvinylidene difluoride membranes (Millipore, Bedford, MA) preactivated in methanol. ARF, p53, p21, and E7 proteins were visualized by direct immunoblotting using rabbit polyclonal antibodies to the ARF C-terminus (Quelle et al. 1995) or to p53 (Ab421), p21, or E7 (all Santa Cruz Biotech.).

Acknowledgments

We thank Rose Mathew and JinLing Wang for assistance, Sam Lucas and Richard Ashmun for FACS analysis, and Frederique Zindy, Kazushi Inoue, and John Cleveland for helpful suggestions. This work was supported by National Institutes of Health grants CA-63230 (G.P.Z.) and CA-71907 (M.F.R.), Cancer Center Core Grant CA-21765, and the American Lebanese Syrian Associated Charities (ALSAC) of St. Jude Children's Research Hospital. C.J.S. is an investigator and J.D.W. is an associate of the Howard Hughes Medical Institute.

The publication costs of this article were defrayed in part by payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 USC section 1734 solely to indicate this fact.

References

- Barak, Y., Juven, T., Haffner, R., and Oren, M. 1993. Mdm2 expression is induced by wild-type p53 activity. *EMBO J.* **12**: 461–468.
- Bates, S., Phillips, A.C., Clarke, P.A., Stott, F., Peters, G., Ludwig, R.L., and Vousden, K.H. 1998. p14^{ARF} links the tumor suppressors pRB and p53. *Nature* **395**: 124–125.
- Carnero, A., Hudson, J.D., Price, C.M., and Beach D.H. 2000. p16^{INK4a} and p19^{ARF} act in overlapping pathways in cellular immortalization. *Nat. Cell Biol.* **2**: 148–155.
- De Stanchina, E., McCurrach, M.E., Zindy, F., Shieh, S.Y., Ferbeyre, G., Samuelson, A.V., Prives, C., Roussel, M.F., Sherr, C.J., and Lowe, S.W. 1998. E1A signaling to p53 involves the p19^{ARF} tumor suppressor. *Genes Dev.* **12**: 2434–2442.
- Donehower, L., Harvey, A.M., Slagle, B.L., McArthur, M.J., Montgomery, C.A. Jr., Butel, J.S., and Bradley, A. 1992. Mice deficient for p53 are developmentally normal but susceptible to spontaneous tumours. *Nature* **356**: 215–221.
- Eischen, C.M., Weber, J.D., Roussel, M.F., Sherr, C.J., and Cleveland, J.L. 1999. Disruption of the ARF-Mdm2-p53 tumor suppressor pathway in Myc-induced lymphomagenesis. *Genes Dev.* **13**: 2658–2669.
- Giacca, A.J. and Kastan, M.B. 1998. The complexity of p53 modulation: Emerging patterns from divergent signals. *Genes Dev.* **12**: 2973–2983.
- Groth, A., Weber, J.D., Willumsen, B.M., Sherr, C.J., and Rousse, M.F. 2000. Oncogenic Ras induces p19^{ARF} and growth arrest in mouse embryo fibroblasts lacking p21^{Cip1} and p27^{Kip1} without activating cyclin D-dependent kinases. *J. Biol. Chem.* (In press).
- Haupt, Y., Maya, R., Kazaz, A., and Oren, M. 1997. Mdm2 promotes the rapid degradation of p53. *Nature* **387**: 296–299.
- Hawley, R.G., Lieu, F.H., Fong, A.Z., and Hawley, T.S. 1994. Versatile retroviral vectors for potential use in gene therapy. *Gene Ther.* **1**: 136–138.
- Hirai, H., Roussel, M.F., Kato, J.Y., Ashmun, R.A., and Sherr, C.J. 1995. Novel INK4 proteins, p19 and p18, are specific inhibitors of the cyclin D-dependent kinases CDK4 and CDK6. *Mol. Cell. Biol.* **15**: 2672–2681.
- Honda, R., Tanaka, H., and Yasuda, H. 1997. Oncoprotein MDM2 is a ubiquitin ligase E3 for tumor suppressor p53.

- FEBS Lett.* **420**: 25–27.
- Honda, R. and Yasuda, H. 1999. Association of p19^{ARF} with Mdm2 inhibits ubiquitin ligase activity of MDM2 for tumor suppressor p53. *EMBO J.* **18**: 22–27.
- Inoue, K., Roussel, M.F., and Sherr, C.J. 1999. Induction of ARF tumor suppressor gene expression and cell cycle arrest by transcription factor DMP1. *Proc. Natl. Acad. Sci. U S A* **96**: 3993–3998.
- Jacks, T., Remington, L., Williams, B.O., Schmitt, E.M., Halachmi, S., Bronson, R.T., and Weinberg, R.A. 1994. Tumor spectrum analysis in p53-mutant mice. *Curr. Biol.* **4**: 1–7.
- Jacobs, J.J., Scheijen, B., Voncken, J.-W., Kieboom, K., Berns, A., and van Lohuizen, M. 1999. Bmi-1 collaborates with c-Myc in tumorigenesis by inhibiting c-Myc-induced apoptosis via INK4a/ARF. *Genes Dev.* **13**: 2678–2690.
- Jones, S.N., Roe, A.E., Donehower, L.A., and Bradley, A. 1995. Rescue of embryonic lethality in Mdm-2-deficient mice by absence of p53. *Nature* **378**: 206–208.
- Kaelin, W.G. Jr. 1999. The p53 gene family. *Oncogene* **18**: 7701–7705.
- Kamijo, T., Bodner, S., van de Kamp, E., Randle, D.H., and Sherr, C.J. 1999. Tumor spectrum in ARF-deficient mice. *Cancer Res.* **59**: 2217–2222.
- Kamijo, T., Weber, J.D., Zambetti, G., Zindy, F., Roussel, M.F., and Sherr, C.J. 1998. Functional and physical interactions of the ARF tumor suppressor with p53 and Mdm2. *Proc. Natl. Acad. Sci. U S A* **95**: 8292–8297.
- Kamijo, T., Zindy, F., Roussel, M.F., Quelle, D.E., Downing, J.R., Ashmun, R.A., Grosveld, G., and Sherr, C.J. 1997. Tumor suppression at the mouse *INK4a* locus mediated by the alternative reading frame product p19^{ARF}. *Cell* **91**: 649–659.
- Ko, L.J. and Prives, C. 1996. p53: Puzzle and paradigm. *Genes & Dev.* **10**: 1054–1072.
- Kubbutat, M.H., Jones, S.N., and Vousden, K.H. 1997. Regulation of p53 stability by Mdm2. *Nature* **387**: 299–303.
- Lohrum, M.A., Ashcroft, M., Kubbutat, M.H., and Vousden, K.H. 2000a. Contribution of two independent MDM2-binding domains in p14^{ARF} to p53 stabilization. *Curr. Biol.* **10**: 539–542.
- Lohrum, M.A., Ashcroft, M., Kubbutat, M.H.G., and Vousden, K.H. 2000b. Identification of a cryptic nucleolar-localization signal in MDM2. *Nat. Cell Biol.* **2**: 179–181.
- Lohrum, M.A.E. and Vousden, K.H. 2000. Regulation and function of the p53-related proteins: Same family, different rules. *Trends Cell Biol.* **10**: 197–202.
- Maestros, R., Dei Tos, A.P., Hamamori, Y., Krasnokutsky, S., Sartorelli, V., Kedes, L., Doglioni, C., Beach, D.H., and Hannon, G.J. 1999. Twist is a potential oncogene that inhibits apoptosis. *Genes Dev.* **13**: 2207–2217.
- McDonnell, T.J., Montes de Oca Luna, R., Cho, S., Amelse, L.L., Chavez-Reyes, A., and Lozano, G. 1999. Loss of one but not two *mdm2* null alleles alters the tumour spectrum in p53 null mice. *J. Pathol.* **188**: 322–328.
- McMasters, K.M., Montes de Oca Luna, R., Peña, J.R., and Lozano, G. 1996. *Mdm2* deletion does not alter growth characteristics of p53-deficient embryo fibroblasts. *Oncogene* **13**: 1731–1736.
- Midgley, C.A., Desterro, J.M., Saville, M.K., Howard, S., Sparks, A., Hay, R.T., and Lane, D.P. 2000. An N-terminal p14^{ARF} peptide blocks Mdm2-dependent ubiquitination in vitro and can activate p53 in vivo. *Oncogene* **19**: 2312–2323.
- Momand, J., Zambetti, G.P., Olson, D.C., George, D., and Levine, A.J. 1992. The mdm-2 oncogene product forms a complex with the p53 protein and inhibits p53-mediated transactivation. *Cell* **69**: 1237–1245.
- Montes de Oca Luna, R., Wagner, D.S., and Lozano, G. 1995. Rescue of early embryonic lethality in mdm2-deficient mice by deletion of p53. *Nature* **378**: 203–206.
- Oliner, J.D., Pietenpol, J.A., Thiagalingam, S., Gyuris, J., Kinzler, K.W., and Vogelstein, B. 1993. Oncoprotein MDM2 conceals the activation domain of tumour suppressor p53. *Nature* **362**: 857–860.
- Palmero, I., Pantoja, C., and Serrano, M. 1998. p19^{ARF} links the tumour suppressor p53 to ras. *Nature* **395**: 125–126.
- Pomerantz, J., Schreiber-Agus, N., Liégeois, N.J., Silverman, A., Alland, L., Chin, L., Potes, J., Chen, K., Orlow, I., Lee, H.-W., Cordon-Cardo, C., and DePinho, R. 1998. The *Ink4a* tumor suppressor gene product, p19^{ARF}, interacts with MDM2 and neutralizes MDM2's inhibition of p53. *Cell* **92**: 713–723.
- Quelle, D.E., Zindy, F., Ashmun, R.A., and Sherr, C.J. 1995. Alternative reading frames of the INK4a tumor suppressor gene encode two unrelated proteins capable of inducing cell cycle arrest. *Cell* **83**: 993–1000.
- Roth, J., Dobbstein, M., Freedman, D.A., Shenk, T., and Levine, A.J. 1998. Nucleo-cytoplasmic shuttling of the hdm2 oncoprotein regulates the levels of the p53 protein via a pathway used by the human immunodeficiency virus rev protein. *EMBO J.* **17**: 554–564.
- Roussel, M.F., Theodoras, A.M., Pagano, M., and Sherr, C.J. 1995. Rescue of defective mitogenic signaling by D-type cyclins. *Proc. Natl. Acad. Sci. U S A* **92**: 6837–6841.
- Schmitt, C.A., McCurrach, M.E., De Stanchina, E., Wallace-Brodeur, R.R., and Lowe, S.W. 1999. INK4a/ARF mutations accelerate lymphomagenesis and promote chemoresistance by disabling p53. *Genes Dev.* **13**: 2670–2677.
- Sherr, C.J. 1998. Tumor surveillance via the ARF-p53 pathway. *Genes Dev.* **12**: 2984–2991.
- Slebos, R.J., Lee, M.H., Plunkett, B.S., Kesis, T.D., Williams, B.O., Jacks, T., Hedrick, L., Kastan, M.B., and Cho, K.R. 1994. p53-dependent G1 arrest involves pRB-related proteins and is disrupted by the human papillomavirus 16 E7 oncoprotein. *Proc. Natl. Acad. Sci. U S A* **91**: 5320–5324.
- Stott, F.J., Bates, S., James, M.C., McConnell, B.B., Starborg, M., Brookes, S., Palmero, I., Ryan, K., Hara, E., Vousden, K.H., and Peters, G. 1998. The alternative product from the human *CDKN2A* locus, p14^{ARF}, participates in a regulatory feedback loop with p53 and MDM2. *EMBO J.* **17**: 5001–5014.
- Tao, W. and Levine, A.J. 1999. p19^{ARF} stabilizes p53 by blocking nucleo-cytoplasmic shuttling of Mdm2. *Proc. Natl. Acad. Sci. U S A* **96**: 6937–6941.
- Weber, J.D., Kuo, M.-L., Bothner, B., DiGiammarino, E.L., Kriwacki, R.W., Roussel, M.F., and Sherr, C.J. 2000. Cooperative signals governing ARF-Mdm2 interaction and nucleolar localization of the complex. *Mol. Cell Biol.* **20**: 2517–2528.
- Weber, J.D., Taylor, L.J., Roussel, M.F., Sherr, C.J., and Bar-Sagi, D. 1999. Nucleolar Arf sequesters Mdm2 and activates p53. *Nat. Cell Biol.* **1**: 20–26.
- Wu, X., Bayle, J.H., Olson, D., and Levine, A.J. 1993. The p53-mdm-2 autoregulatory feedback loop. *Genes Dev.* **7**: 1126–1132.
- Zeng, X., Chen, L., Jost, C.A., Maya, R., Keller, D., Wang, X., Kaelin, W.G. Jr., Oren, M., Chen, J., and Lu, H. 1999. MDM2 suppresses p73 function without promoting p73 degradation. *Mol. Cell Biol.* **19**: 3257–3266.
- Zhang, Y., Xiong, Y., and Yarbrough, W.G. 1998. ARF promotes MDM2 degradation and stabilizes p53: ARF-INK4a locus deletion impairs both the Rb and p53 tumor suppressor pathways. *Cell* **92**: 725–734.
- Zindy, F., Eischen, C.M., Randle, D., Kamijo, T., Cleveland, J.L., Sherr, C.J., and Roussel, M.F. 1998. Myc signaling via the ARF tumor suppressor regulates p53-dependent apoptosis and immortalization. *Genes Dev.* **12**: 2424–2433.

p53-independent functions of the p19^{ARF} tumor suppressor

Jason D. Weber, John R. Jeffers, Jerold E. Rehg, et al.

Genes Dev. 2000, **14**:

Access the most recent version at doi:[10.1101/gad.827300](https://doi.org/10.1101/gad.827300)

References

This article cites 48 articles, 22 of which can be accessed free at:
<http://genesdev.cshlp.org/content/14/18/2358.full.html#ref-list-1>

License

Email Alerting Service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).