Conversion of a gene-specific repressor to a regional silencer

Laura N. Rusché and Jasper Rine¹

Department of Molecular and Cell Biology, University of California, Berkeley, California 94720, USA

In *Saccharomyces cerevisiae*, gene silencing at the *HMR* and *HML* loci is normally dependent on Sir2p, Sir3p, and Sir4p, which are structural components of silenced chromatin. Sir2p is a NAD⁺-dependent histone deacetylase required for silencing. Silencing can be restored in cells lacking Sir proteins by a dominant mutation in *SUM1*, which normally acts as a mitotic repressor of meiotic genes. This study found that mutant Sum1-1p, but not wild-type Sum1p, associated directly with *HM* loci. The origin recognition complex (ORC) was required for Sum1-1p-mediated silencing, and mutations in *ORC* genes reduced association of Sum1-1p with the *HM* loci. Sum1-1p-mediated silencing also depended on *HST1*, a paralog of *SIR2*. Both Sum1-1p and wild-type Sum1p interacted with Hst1p in coimmunoprecipitation experiments. Therefore, the *SUM1-1* mutation did not change the affinity of Sum1p for Hst1p, but rather relocalized Sum1p to the *HM* loci. Sum1-1-Hst1p action led to hypoacetylation of the nucleosomes at *HM* loci. Thus, Sum1-1p and Hst1p could substitute for Sir proteins to achieve silencing through formation of a compositionally distinct type of heterochromatin.

[Key Words: Position effect; epigenetic; histone deacetylase; meiosis; chromatin immunoprecipitation]

Received December 13, 2000; revised version accepted February 20, 2001.

Transcriptional repression is as important as activation in determining the expression profile of a genome, yet it is less well understood. Repression mechanisms are often categorized as being either locus specific or regional. In yeast, locus-specific repression is mediated by regulatory sites, typically located in the vicinity of promoters, which are bound by site-specific DNA-binding proteins. These proteins recruit other protein complexes, or corepressors that mediate repression of adjacent genes.

In yeast, the two best-studied corepressors are the Ssn6-Tup1 and the Sin3-Rpd3 complexes. The Sin3-Rpd3 complex (Zhang et al. 1998; Dorland et al. 2000) is recruited to target loci by site-specific DNA-binding proteins, such as Ume6p (Kadosh and Struhl 1997). The binding of Rpd3p results in highly localized deacetylation of histones H3 and H4 (Kadosh and Struhl 1998; Rundlett et al. 1998). Rpd3p homologs in mammals, HDAC1 and HDAC2, are also in large complexes and are recruited to specific sites of repression (for review, see Ayer 1999; Ahlinger 2000). Like Sin3-Rpd3, the Ssn6-Tup1 complex is recruited to chromatin by site-specific DNA-binding proteins, such as $\alpha 1/\alpha 2$. Tup1p binds to the tails of histones H3 and H4 in vitro (Edmondson et al. 1996) and forms a scaffold along nucleosomes in vivo,

thereby stabilizing the positions of nucleosomes in an ordered, repressive conformation (Ducker and Simpson 2000). The Ssn6-Tup1 complex coimmunoprecipitates with Rpd3p and Hos2p, histone deacetylases (Watson et al. 2000), and may also inhibit transcription by direct interaction with the transcription machinery (Lee et al. 2000). For both the Ssn6-Tup1 and Sin3-Rpd3 repressor complexes, repression is a local phenomenon, restricted to individual genes adjacent to each target site (DeRisi et al. 1997).

Silencing has been distinguished operationally from repression by its gene nonspecific nature and its ability to block or reduce gene expression throughout a chromosomal domain, such as at telomeres or at the two *HM* loci of *Saccharomyces*. Silencing at the *HM* loci is mediated by regulatory sites known as the *E* and *I* silencers that flank both *HMR* and *HML*. These silencers provide binding sites for ORC, Rap1p, and Abf1p, which individually bind to a variety of other sites in the genome that are not subject to silencing. The silencer-binding proteins interact with Sir1p, Sir3p, and Sir4p (Moretti et al. 1994; Triolo and Sternglanz 1996), and it is presumed that a complex containing all four Sir proteins assembles at the silencer. Sir3p and Sir4p interact with histone H3 and H4 tails (Hecht et al. 1995), and this interaction is thought to allow them to spread throughout *HMR* and *HML*. Sir2p is presumed to spread with Sir3p and Sir4p, and its histone deacetylase activity results in hypoacetylation of nucleosomes throughout *HMR* and *HML* (Braunstein et al. 1993), thereby creating heterochromatin.

¹Corresponding author.

E-MAIL [jriner@uclink4.berkeley.edu](mailto:jrine@uclink4.berkeley.edu); FAX (510) 642-6420.

Article and publication are at www.genesdev.org/cgi/doi/10.1101/gad.873601.

Similar silencing mechanisms are used at other loci. Sir2p is also found at telomeres and the rDNA array (Hecht et al. 1996; Gotta et al. 1997), at which its deacetylase activity is required for silencing (Imai et al. 2000). Silencing at telomeres requires *SIR3* and *SIR4*, but silencing at rDNA does not. Whether other proteins serve the roles of Sir3p and Sir4p in rDNA silencing is unknown. The Sir2p family of NAD⁺-dependent histone deacetylases, including Hst1p–Hst4p, may also mediate other silencing or silencing-like phenomena (Brachmann et al. 1995).

This study describes a repressor protein, Sum1p, which bridges the gap between repression and regional silencing. *SUM1* encodes a sequence-specific DNA-binding protein that binds near the promoters of genes required at mid-sporulation (Xie et al. 1999). Binding of Sum1p to these genes results in their repression in mitotic cells, but, as with other repressors, does not repress the neighboring genes (Xie et al. 1999). The *SUM1* sequence does not suggest any function, except for a potential AT-hook domain that could mediate DNA binding.

SUM1 was discovered by virtue of the *SUM1-1* mutation, which restores transcriptional repression of *HMR* and *HML*, but not of *MAT*, in the absence of Sir proteins (Klar et al. 1985; Livi et al. 1990; Laurenson and Rine 1991). Repression is complete at *HMR*, allowing *MAT* α *sir* cells to mate, but it is only partial at *HML*, and *MAT* α *sir* cells do not mate. *SUM1-1* is a dominant mutation (Laurenson and Rine 1991) that creates a protein with an altered function, since neither a null mutation nor overexpression of the wild-type protein results in the *SUM1-1* phenotype (Chi and Shore 1996). The *SUM1-1* phenotype results from a single amino acid change near the C terminus of the protein (Chi and Shore 1996). Thus, a single mutation in a transcriptional regulator dramatically alters which genes it regulates.

To understand how *SUM1-1* achieves silencing in the absence of the Sir proteins, which have been considered essential structural components of heterochromatin, and to understand how a single mutation in a repressor can so dramatically alter its targets, we studied the mechanism of Sum1-1p-mediated silencing.

Results

Sum1-1p acted directly at HM loci

In principle, Sum1-1p could either directly repress transcription at *HMR* and *HML*, or it could alter the expression of another protein, which, in turn, leads to Sir-independent silencing. To determine whether Sum1-1p acts directly at *HM* loci, a chromatin immunoprecipitation assay was developed using an N-terminal myc-tagged allele of Sum1-1p expressed from the *SUM1* promoter at the *SUM1* locus. To assess the function of the tagged Sum1-1p, its ability to restore silencing at *HMR* was evaluated in *MAT* α *sir2* Δ cells. Loss of silencing leads to simultaneous expression of *a* and α information, thereby disrupting mating ability. Strains containing the

tagged and untagged versions of Sum1-1p both mated well (data not shown), indicating that *HMR* was silenced. In a more sensitive test of repression, both *SUM1-1* and *myc-SUM1-1* repressed transcription of *a1* mRNA to an undetectable level in *sir2* Δ cells (data not shown). Therefore, the tagged Sum1-1p was fully functional.

To determine whether Sum1-1p was associated with *HM* loci, myc-Sum1-1p and associated DNA were immunoprecipitated from crude extracts of formaldehyde cross-linked cells. This DNA was analyzed by simultaneous PCR amplification of the *HMR-E* silencer region and the promoter region of *SSC1*, whose transcription is not affected by the *SUM1-1* mutation (data not shown) and served as a negative control. To monitor whether the PCR reaction was sensitive to the amount of starting DNA, a second set of PCR reactions was performed on a twofold dilution of the starting material. Sum1-1p was preferentially associated with *HMR-E* relative to the *SSC1* negative control (Fig. 1A, cf. lanes 1,2 and 3,4). Neither *HMR* nor *SSC1* was precipitated from cells lacking the myc tag (lane 5) or in the absence of antibody (lane 6).

Chromatin silenced by Sir proteins is widely assumed to contain Sir proteins throughout the silenced domain, although they have only been shown to be at one site (Hecht et al. 1996). To determine whether Sum1-1-mediated silencing behaved similarly, the distribution of Sum1-1p across *HMR* was examined (Fig. 1B). In addition to *HMR-E*, Sum1-1p also associated with the *HMR-I* silencer (Fig. 1B, lanes 10–12; cf. ratios of *HMR* products with *SSC1* products in input [i] and immunoprecipitate [+]), and with two regions overlapping the coding sequence of the *a1* and *a2* genes (X-Ya, lanes 4–6; Ya-Z1, lanes 7–9). Therefore, Sum1-1p was clearly present at multiple sites across the *HMR* locus.

The size of the region repressed by Sir-mediated silencing is constrained by boundary elements (Bi et al. 1999; Donze et al. 1999), and the Sir proteins are thought to associate only with sequences between these boundaries. Similarly, Sum1-1p was not associated with a region just beyond a boundary element at the right side of *HMR-I* (Fig. 1C, lanes 1–3). Therefore, Sum1-1p appeared to associate with the *HMR* locus, much as the Sir proteins are thought to.

Sum1-1p mediates transcriptional repression at *HML* as well as *HMR*, but not at *MAT* (Livi et al. 1990). This phenotype predicted that Sum1-1p should also be found at *HML* but not at *MAT*. Sum1-1p associated with both silencers at *HML* (Fig. 1C, lanes 4–9), and was not associated with *MAT* (lanes 10–12). Therefore, Sum1-1p was localized to the genes that it silenced, ruling out indirect models in which Sum1-1p altered expression of some other gene(s) that, in turn, caused silencing.

SUM1-1 mutation relocated Sum1p

In principle, the *SUM1-1* mutation could restore silencing to *Sir*[−] strains by relocating the Sum1p repressor to *HM* loci. Alternatively, wild-type Sum1p may normally

The *SMK1* gene served as a positive control for immunoprecipitation because Sum1p binds to the promoter of *SMK1* in mobility shift assays (Xie et al. 1999). Consistent with this result, Sum1p associated with the *SMK1* promoter in chromatin immunoprecipitation assays (Fig. 2A, cf. lanes 1,2 and 3,4). Sum1p also associated with the *SPR3* promoter, as expected (Fig. 2B, lanes 4–6). In contrast, in the same samples, wild-type Sum1p was not associated with any of the silencers (lanes 7–18). Therefore, the *SUM1-1* mutation restored silencing to *sir* mutant strains by redirecting Sum1p to *HM* loci.

To determine whether the *SUM1-1* mutation reduced the association of the mutant protein with wild-type binding sites, or simply expanded the range of sites to which it bound, the association of Sum1-1p with *SMK1* and *SPR3* was compared with wild-type Sum1p's asso-

ciation. The amount of these loci immunoprecipitated with myc-Sum1-1p was three to sixfold less than was obtained with wild-type myc-Sum1p (Fig. 1C, lanes 16–21 for Sum1-1p and Fig. 2B, lanes 1–6 for Sum1p; cf. fraction of input *SMK1* or *SPR3* recovered). Therefore, Sum1-1p was only slightly associated with the promoter regions of *SMK1* and *SPR3*, indicating that the mutant protein lost most of its ability to associate with its original site of action. The *SUM1-1* mutation resulted in significant derepression of both *SMK1* and *SPR3* (Fig. 2C, cf. lanes 1 and 3), although these genes were still somewhat repressed compared with a *sum1Δ* strain (lane 2).

ORC was required for Sum1-1p-mediated repression of HMR

To identify elements required for the association of Sum1-1p with *HM* loci, sequences required for Sum1-1p-mediated silencing were identified. Plasmids containing the *HMRa* locus flanked by an intact *I* silencer and wild-type or mutant *E* silencers were transformed into *MATα* strains. Silencing mediated by Sir proteins (Fig. 3A, left) or Sum1-1p (right) was detected by a mating assay that led to growth. Sum1-1p could silence *HMR* containing a wild-type *E* silencer (Fig. 3A, row 2), but could not silence it when the *E* silencer was deleted (Fig. 3A, row 3). Therefore, sequences within the *E* silencer were required for Sum1-1p-mediated silencing.

To determine whether one or more of the three known protein binding sites were important for Sum1-1p silencing, a synthetic silencer, which resembles the wild-type silencer only in the presence and spacing of the ORC, Rap1p, and Abf1p-binding sites (McNally and Rine 1991), was tested. The synthetic silencer supported Sum1-1p-mediated silencing (Fig. 3A, row 4). Therefore, Sum1-1p-mediated silencing depended on one or more of the known binding sites, and not on other sequences within the natural *E* silencer. Mutation of both the Rap1p and Abf1p-binding sites together had no effect on Sum1-1p-mediated silencing (Fig. 3A, row 5). In contrast, mutation of the ORC-binding site eliminated Sum1-1p-mediated silencing (Fig. 3A, row 6). Therefore, the binding site for ORC, but not for Rap1p or Abf1p, was important for Sum1-1p-mediated silencing.

The requirement for an ORC-binding site for Sum1-1p-mediated silencing implied that ORC itself was required. *SUM1-1* suppressed *sir* silencing defects in the presence of wild-type ORC (Fig. 3B, first column), but silencing was completely abolished by the *orc5-1* (second column) or *orc2-1* (third column) temperature-sensitive mutations in cells grown at the permissive temperature.

The only known role of ORC in Sir-mediated silencing at *HM* loci is binding to Sir1p (Fox et al. 1997; Gardner et al. 1999). The N-terminal domain of Orc1p is required for this interaction (Triolo and Sternglanz 1996). Deletion of this domain disrupts Sir-mediated silencing but not the replication function of ORC (Bell et al. 1995). Similarly, the N-terminal domain of Orc1p was also required for Sum1-1p-mediated silencing. The *orc1Δ1-235* allele completely eliminated Sum1-1p-mediated silenc-

Figure 3. ORC was required for Sum1-1p-mediated silencing. (A) *MATα* haploids of the genotypes *SIR SUM1* (W303) and *sir2Δ SUM1-1* (JRY7152) were transformed with a plasmid vector (pRS316, line 1) or the vector containing *HMRa* (pJR759, line 2), *HMRa-eΔ* (pJR760, line 3), *HMRa*-synthetic silencer (pJR900, line 4), *HMRa*-no Rap1p/Abf1p binding sites (pJR891, line 5), or *HMRa*-synthetic silencer/no ORC binding site (pJR901, line 6). Representative transformants were tested for their mating ability by replica plating onto an *a* mating-type tester lawn (TD4) on minimal medium requiring plasmid retention for growth. Mating was indicated by growth of prototrophic diploids. (B) *MATα* haploids of the genotypes *SIR SUM1-1* (MC89), *sir2Δ SUM1-1* (JRY7152), *sir3Δ SUM1-1* (JRY7161), *sir4Δ SUM1-1* (JRY7166), *SIR orc5-1 SUM1-1* (JRY7149), *sir2Δ orc5-1 SUM1-1* (JRY7157), *sir3Δ orc5-1 SUM1-1* (JRY7164), *sir4Δ orc5-1 SUM1-1* (JRY7169), *SIR orc2-1 SUM1-1* (JRY7150), *sir2Δ orc2-1 SUM1-1* (JRY7159), *orc5-1* (JRY7146), and *orc2-1* (JRY7147) were tested for their mating ability as in A using mating tester JRY2726. (C) *MATα* haploids of the genotypes *SIR SUM1* (W303), *sir2Δ myc-SUM1-1* (JRY7153), *sir2Δ myc-SUM1-1 orc1Δ1-235* (*orc1ΔN*, JRY7176), and *sir2Δ myc-SUM1-1 ORC1* (Q236H, I237M) (*orc1***, JRY7177) were tested for their mating ability as in B.

ing (Fig. 3C). In fact, a Q236H I237M mutation, which has no effect on Sir-mediated silencing (Bell et al. 1995), significantly reduced Sum1-1p-mediated silencing (Fig. 3C). Thus, specific residues in the N-terminal portion of Orc1p were required for Sum1-1p-mediated silenc-

Figure 5. *HST1* was required for Sum1-1p-mediated silencing. (A) *MATα* haploids of the genotypes *SIR SUM1* (W303), *sir2Δ* (JRY4565), *sir3Δ* (JRY7160), *sir4Δ* (JRY7165), *SIR SUM1-1* (MC89), *sir2Δ SUM1-1* (JRY7152), *sir3Δ SUM1-1* (JRY7161), *sir4Δ SUM1-1* (JRY7166), *SIR SUM1-1 hst1Δ* (JRY7148), *sir2Δ SUM1-1 hst1Δ* (JRY7154), *sir3Δ SUM1-1 hst1Δ* (JRY7162), *sir4Δ SUM1-1 hst1Δ* (JRY7167), *SIR hst1Δ* (JRY7145), *sir2Δ hst1Δ* (JRY7156), *sir3Δ hst1Δ* (JRY7163), and *sir4Δ hst1Δ* (JRY7168) were tested for their mating ability as in Fig. 3B. (B) RNA from *MATα* haploids of the genotypes *SIR SUM1* (lane 1, W303), *sir2Δ* (lane 2, JRY4565), *sir2Δ SUM1-1* (lane 3, JRY7152), and *sir2Δ SUM1-1 hst1Δ* (lane 4, JRY7154) was analyzed for *a1* (top) or *ACT1* (bottom) mRNA.

promoter, was only partially functional, judged by its ability to restore silencing in a *sir2Δ SUM1-1* strain compared with wild-type Hst1p (data not shown). However, this HA-tagged Hst1p expressed from a CEN plasmid restored complete silencing (data not shown), indicating that the tagged Hst1p must still interact with any partner necessary for silencing function. Tagged Hst1p was immunoprecipitated from strains containing myc-tagged Sum1-1p, and the immunoprecipitates were evaluated for the presence of Sum1-1p (Fig. 6A). Sum1-1p coprecipitated with HA-tagged Hst1p (Fig. 6A, lane 5). Conversely, Hst1p coprecipitated with myc-tagged Sum1-1p (Fig. 6B, lane 5). Therefore, Sum1-1p and Hst1p were physically associated.

To determine whether the *SUM1-1* mutation changed the affinity of Sum1p for Hst1p, coimmunoprecipitation experiments were also conducted by use of a tagged ver-

sion of wild-type Sum1p. Sum1p coimmunoprecipitated with tagged Hst1p (Fig. 6A, lane 8), and Hst1p coimmunoprecipitated with tagged Sum1p (Fig. 6B, lane 8). In both cases, the amount of material precipitated was comparable with the amount precipitated with mutant Sum1-1p. Therefore, the change in function resulting from the *SUM1-1* mutation was probably not due to a change in affinity for the Hst1p corepressor.

Because Sum1-1p was bound to Hst1p and also associated with *HM* loci, it was likely that Hst1p was also associated with *HM* loci. Additionally, *HST1* was required with *SUM-1* for hypoacetylation of *HM* loci in *sir2Δ* strains, as described below, again suggesting that Hst1p was associated with *HM* loci. However, we were unable to preferentially precipitate any DNA known to associate with either Sum1-1p or Sum1p with antibodies against HA-tagged Hst1p (data not shown). Perhaps Hst1p does not cross-link well, or is too indirectly associated with the DNA to be detected by this assay. Alternatively, Hst1p may be only transiently associated with DNA.

The presence of Sum1-1p, but not Sum1p, at *HM* loci implied that Sum1-1p was responsible for bringing Hst1p to *HM* loci, and not the other way around. However, it was still possible that Hst1p itself contributed to the presence Sum1-1p at *HM* loci. In fact, Sum1-1p association with *HMR* was slightly reduced in an *hst1Δ* strain, although to a lesser extent than in *orc* mutants (Fig. 6C). Hst1p may stabilize the association of Sum1-1p with *HMR*, perhaps by providing another binding surface.

Sum1-1p-mediated repression resulted in histone deacetylation

Sir2 family members, including Hst1p, are NAD⁺-dependent histone deacetylases (Imai et al. 2000; Landry et al. 2000; Smith et al. 2000). Nucleosomes at silenced *HM* loci are hypoacetylated, particularly in response to Sir2p overexpression (Braunstein et al. 1993). The role of Hst1p in Sum1-1p-mediated repression (Fig. 5A,B) implied that histones H3 and H4 at *HM* loci would be hypoacetylated in a *SUM1-1* mutant. Therefore, chromatin immunoprecipitation assays using antibodies against acetylated histones were performed to test this hypothesis. As expected, all four silencers (Fig. 7, panels 1–4) and the *a1*–*a2* coding region (Fig. 7, panels 5,6) were under-represented in the immunoprecipitated material from the *SIR* strain (Fig. 7, lanes A), but not from the *sir2Δ* strain (Fig. 7, lanes B). In contrast, *MAT* and *ACT1* (Fig. 7, panels 7,8), nonsilenced loci, were nearly equally abundant in both samples. Therefore, the *HM* loci were underacetylated in the *SIR* strain, as expected. In a parallel experiment using a *sir2Δ SUM1-1* strain (Fig. 7, lanes C), all four silencers (Fig. 7, panels 1–4) and the *a1*–*a2* coding region (Fig. 7, panels 5,6) were hypoacetylated, whereas *MAT* and *ACT1* (Fig. 7, panels 7,8) were not. Therefore, Sum1-1p-mediated silencing resulted in hypoacetylation of the histone tails at both *HM* loci.

To determine whether this deacetylation was dependent on *HST1*, a *sir2Δ SUM1-1 hst1Δ* strain (Fig. 7, lanes

lanes 1,5,7), Sum1-1p (pJR2293, lanes 2,6), 3myc-Sum1p (pJR2292, lanes 3,8,10) or Sum1p (pJR2294, lanes 4,9). A total of 1/30 of input (lanes 1–4) or 1/15 of immunoprecipitated samples (lanes 5–10) was analyzed for the presence of Hst1–5HAp by immunoblotting. (C) Chromatin immunoprecipitations used *MATα* haploids of the genotypes *sir2Δ HST1 7myc-SUM1-1* (JRY7153), *sir2Δ hst1Δ 7myc-SUM1-1* (JRY7178), and *sir2Δ hst1Δ SUM1-1* (JRY7157). A total of 1/23,000 (lanes 1,3) or 1/46,000 (lanes 2,4) of the input DNA or 1/25 (lanes 5,7,9–11) or 1/50 (lanes 6,8) of the immunoprecipitated DNA was analyzed.

D) was examined. In contrast to the *sir2Δ SUM1-1* strain, all four silencers and the *a1–a2* coding region were acetylated to an extent comparable with the control loci, indicating that *HST1* was required for the Sum1-1p-mediated deacetylation of *HM* loci. This result was consistent with Hst1p being the responsible histone deacetylase. Thus, both Sir2p and Hst1p acted through a common mechanism to generate heterochromatin.

If nucleosome deacetylation were dependent on association of Sum1-1p and Hst1p with chromatin, then deacetylated nucleosomes, like Sum1-1p itself, should not extend beyond a boundary element at *HMR*. The chromatin distal to the boundary on the *I*-silencer side was equally acetylated in all strains (Fig. 7, panel 9). Therefore, within the resolution of the experiment, nucleosome deacetylation correlated with the presence of Sum1-1p.

Wild-type Sum1p also interacted with Hst1p (Fig. 6; Xie et al. 1999). Therefore, it was possible that nucleosomes at the promoters of genes repressed by wild-type Sum1p were also deacetylated. To test this hypothesis, the immunoprecipitates were examined for *SMK1* or *SPR3* promoter regions. If Sum1p–Hst1p-dependent deacetylation occurred, it would be greatest in *SUM1* strains (Fig. 7, lanes A,B), reduced in a mutant *SUM1-1* strain (Fig. 7, lanes C), and not seen when the Hst1p deacetylase is absent (Fig. 7, lanes D). However, *SMK1* and *SPR3* promoters (Fig. 7, panels 10,11) were equally well represented in all four strains, indicating that widespread histone deacetylation did not occur at these promoters.

Figure 6. Hst1p was in complexes with both Sum1-1p and wild-type Sum1p. (A) A *MATα* haploid of the genotype *sir2Δ sum1Δ hst1Δ* (JRY7151) was transformed with a plasmid expressing 3myc-Sum1-1p (pJR2291, lanes 1,2,5–7) or 3myc-Sum1p (pJR2292, lanes 3,4,8–10) and a second plasmid expressing Hst1–5HAp (pJR2289, lanes 1,3,5,7,8,10) or Hst1p (pJR2288, lanes 2,4,6,9). Transformed cells were grown in minimal medium requiring maintenance of both plasmids. A total of 1/30 of input (lanes 1–4) or 1/15 of immunoprecipitated samples (lanes 5–10) was analyzed for the presence of myc-Sum1-1p or myc-Sum1p by immunoblotting. Negative control lanes had samples prepared with an untagged version of Hst1p (lanes 6,9) or without antibody in the immunoprecipitation (lanes 7,10). (B) A *MATα* haploid of the genotype *sir2Δ sum1Δ hst1Δ* (JRY7151) was transformed with a plasmid expressing Hst1–5HAp (pJR2289, all lanes) and a second plasmid expressing 3myc-Sum1-1p (pJR2291,

SUM1-1 slow growth was dependent on HST1 and ORC

Consistent with previous reports (Chi and Shore 1996), we found that *SUM1-1* cells divided slowly, with a doubling time between 3 and 3.5 h at 23°, compared with wild-type cells, which doubled approximately every 2 h. This slow growth was not due to silencing of the *HM* loci, which are also silenced in wild-type strains, or to the loss of wild-type *SUM1* function, because *sum1Δ* strains doubled every 2 h. Sum1-1p may affect the growth rate by partially silencing one or more critical genes. Alternatively, Sum1-1p might interfere with DNA replication, through interaction with ORC. Interestingly, *HST1* and ORC, which are both required for Sum1-1p-mediated silencing at *HM* loci, were required for this slow growth. *SUM1-1 orc* and *SUM1-1 hst1* strains doubled every 2 to 2.5 h. Thus, Sum1-1p, Hst1p, and ORC may act together at other loci.

Discussion

The mechanism of Sum1-1p-mediated silencing

This study has established the mechanism of action of the enigmatic suppressor mutation, *SUM1-1*. The paradox of the *SUM1-1* mutation has been its ability to restore silencing in the absence of Sir proteins, which are key structural components of heterochromatin. We discovered that the *SUM1-1* mutation altered Sum1p in a manner that led to the formation of a compositionally distinct form of heterochromatin, in which Sum1-1p, Hst1p, and perhaps other proteins substitute for Sir pro-

Figure 7. Sum1-1p-mediated silencing was associated with histone deacetylation. Chromatin immunoprecipitations used *MATα* haploids of the genotypes *SIR SUM1* (lanes A, W303), *sir2Δ* (lanes B, JRY4565), *sir2Δ SUM1-1* (lanes C, JRY7152), and *sir2Δ SUM1-1 hst1Δ* (lanes D, JRY7154). DNA immunoprecipitated with one of two different anti-acetyl-histone H4 antibodies (AcH4, lanes 5–8,9–12) or anti-acetyl-histone H3 antibody (AcH3, lanes 13–16) was analyzed by simultaneous PCR amplification of the indicated region (top) and the *SSC1* promoter (bottom). A total of 1/25,000 of the input DNA (lanes 1–4) and 1/50 (lanes 5–8,17,18) or 1/250 (lanes 9–16) of the immunoprecipitated DNA was analyzed. Negative control lanes had samples prepared without antibody in the immunoprecipitation (lanes 17,18).

teins. Sum1-1p was associated with multiple sites within *HMR* and *HML* (Fig. 1B,C), and therefore, acted directly at those loci, where wild-type Sum1p does not act. Sum1-1p required *ORC* (Fig. 3) and *HST1* (Fig. 5) for silencing. *ORC* was required for association of Sum1-1p with *HM* loci (Fig. 4B,C). Hst1p was physically associated with Sum1-1p (Fig. 6) and was required for Sum1-1p-associated deacetylation of histones at *HM* loci (Fig. 7). These data inspired a model for how Sum1-1p causes silencing. In this model, Sum1-1p, in partnership with Hst1p and perhaps other proteins, associates with *ORC*

at silencers. The deacetylase activity of Hst1p then leads to hypoacetylation of nucleosomes at *HM* loci.

This study also explored how the *SUM1-1* mutation changes the action of Sum1p. The mutation altered the localization of the protein, thereby reducing repression of wild-type Sum1p targets and causing new repression at *HM* loci (Figs. 1 and 2). It did not change the affinity of Sum1p for Hst1p (Fig. 6), but did result in a new requirement for *ORC* (Fig. 3; data not shown). Perhaps the mutation, a threonine to isoleucine change in a locally hydrophobic region (Chi and Shore 1996), creates an especially hydrophobic sticky patch, analogous to the sickle cell anemia allele, and this hydrophobic patch creates an affinity for *ORC*. Alternatively, the mutation could reduce site-specific DNA binding, thereby permitting the protein to associate with other, lower-affinity sites. Thus, the *SUM1-1* mutation either created or exposed an affinity for *ORC*, which in turn directed the protein to new loci.

Although wild-type Sum1p binds DNA directly (Xie et al. 1999), several lines of evidence favored models in which *ORC* was required for Sum1-1p association with *HM* loci through a physical link between the two proteins. First, Sum1-1p did not associate with *HM* loci through the wild-type Sum1p-binding site, because Sum1p was not found at *HM* loci (Fig. 2B). Moreover, Sum1-1p had reduced association with promoters containing Sum1p-binding sites (Fig. 1C; M. Pierce and A. Vershon, pers. comm.). Second, *ORC* was required for Sum1-1p-mediated silencing (Fig. 3) and contributed to association of Sum1-1p with *HM* loci (Fig. 4B,C). Therefore, Sum1-1p was likely targeted to *HM* loci through *ORC*.

Sum1-1p-mediated silencing also required *HST1*. Hst1p was physically associated with Sum1-1p (Fig. 6A,B), which in turn was associated with *HM* loci (Fig. 1B,C). In addition, Hst1p, a histone deacetylase, together with Sum1-1p were required for hypoacetylation of *HM* loci in *sir2Δ* strains (Fig. 7), implying that Hst1p acted at, and therefore was associated with, *HM* loci. Hst1p did not recruit Sum1-1p to *HM* loci, because wild-type Sum1p was also physically associated with Hst1p (Fig. 6A,B) but was not found at *HM* loci (Fig. 2B). Therefore, Hst1p was brought to the *HM* loci by Sum1-1p and acted as a cosilencer through a mechanism similar to that of the Rpd3p corepressor.

The mechanism of wild-type Sum1p repression

Comparison of Sum1-1p-mediated silencing and Sum1p-mediated repression indicated that the *SUM1-1* mutation changed the way Sum1p functions. For example, *ORC* appeared not to play a role in repression by wild-type Sum1p because genes repressed by Sum1p were not derepressed in *orc* mutant strains (data not shown). However, these experiments required use of hypomorphic rather than null alleles of *ORC* genes. Therefore, the possibility of a role for *ORC* in Sum1p function could not be completely excluded.

Another difference between wild-type Sum1p-mediated repression and Sum1-1p-mediated silencing was the extent of hypoacetylation of nucleosomes at repressed loci. This study did not detect hypoacetylation of the promoter regions of *SMK1* and *SPR3*. Either Sum1p repression at these loci did not involve stable histone deacetylation, or the region of deacetylation was short or dispersed, and thus was not detected by this assay. In contrast, the hypoacetylation caused by *SUM1-1* was evident over a region spanning several kilobases.

The roles of ORC and Hst1p in other types of repression

The requirement for ORC in Sum1-1p-mediated silencing may reflect a fundamental role of ORC in multiple chromatin structures. In yeast, the N-terminal portion of Orc1p is required for Sir-mediated silencing but is dispensable for replication (Bell et al. 1995). Deletion of the Orc1p N-terminal domain also disrupted Sum1-1p-mediated silencing (Fig. 3C). This domain interacts with Sir1p (Triolo and Sternglanz 1996) and is critical in recruiting Sir1p to the silencer (Gardner et al. 1999). Similarly, in *Drosophila*, mutations in *ORC1* affect position and variegation, and the N terminus of Orc1p recruits HP1 to heterochromatin (Pak et al. 1997). Together, these results suggest a general role for ORC in heterochromatin and imply that the N-terminal portion of Orc1p has a special silencing function. Perhaps the silencing proteins recruited by Orc1p, including HP1, Sir1p, and Sum1-1p, share a common structural motif.

The role of *HST1* in Sum1-1p-mediated silencing has implications for the roles of *HST2-4* and other *SIR2* orthologs. Much as Hst1p, a histone deacetylase, was brought to *HM* loci by Sum1-1p, the Sir2p family of proteins may generally act as cosilencers recruited to chromosomal loci by adapter proteins analogous, but not necessarily homologous, to Sum1-1p. Once at these loci, they would act enzymatically to deacetylate histone tails, thereby altering chromatin structure. Given that the Sir2 family has five members in yeast, at least seven members in humans, and orthologs in essentially all genomes analyzed, recruitment of these proteins by repressors may be a common theme in gene silencing.

Other sites and types of repression

There are interesting parallels and differences between the mechanisms of Sir-mediated silencing and Sum1-1p-mediated silencing. Both require the *HMR-E* silencer. However, Sir-mediated silencing utilizes three silencer-bound proteins, whereas Sum1-1p-mediated silencing requires ORC but not the binding of Rap1p or Abf1p. In both cases, the silencer-bound proteins appear to act by recruiting silencing proteins, either the Sir proteins or Sum1-1p and Hst1p. These complexes, containing either the Sir2p or Hst1p deacetylase, then appear to associate with multiple sites throughout the *HM* loci, resulting in

the deacetylation of nucleosomes. The requirement for a deacetylase to act in the place of Sir2p during Sum1-1p-mediated silencing strengthens the link between hypoacetylation and repression. Whether Sum1-1p carries out the functions of both Sir3p and Sir4p or other as yet unknown analogous proteins associate with Sum1-1p and Hst1p remains to be determined. Therefore, for both Sum1-1p and Sir-mediated silencing, the requirement for silencer-binding proteins and the involvement of deacetylated nucleosomes were the same, but the proteins that create and ultimately constitute the heterochromatin differed.

Sir-mediated and Sum1-1p-mediated silencing represent two separate silencing routes that can be, but are not always, independent. The results of this and other studies indicate that, in some cases, Sum1-1p interacts with the Sir proteins. In a *SIR SUM1-1* strain, a synthetic silencer bearing a mutation of the ORC-binding site is silenced (Laurenson and Rine 1991). In contrast, in this study, the same silencer was not silenced in a *sir2Δ SUM1-1* strain (Fig. 3A). This lack of silencing was also observed in a *sir3Δ SUM1-1* strain (data not shown). Thus, in the original experiment, the importance of the ORC-binding site for Sum1-1p-mediated silencing was masked by the presence of Sir proteins. Similarly, *SIR2* is also required for *SUM1-1* to suppress the silencing defect of a strain bearing both *rap1-12* and a deletion of the ORC-binding site at *HMR* (Chi and Shore 1996). Therefore, it appears that when both Sir proteins and Sum1-1p are present, the two silencing routes can jointly silence a locus that neither could silence alone.

The interaction of Sum1-1p with Sir proteins and the requirement for ORC in Sum1-1p-mediated silencing explained the puzzling observation that *SUM1-1* does not suppress a *sir2Δ* defect in telomeric silencing but does enhance telomeric silencing in a *SIR* strain (Chi and Shore 1996). Like *HM* silencing, telomeric silencing requires Sir2p, Sir3p, and Sir4p. However, synthetic telomeres that can initiate silencing lack ORC-binding sites, consisting instead of polymers of Rap1p-binding sites. Because Rap1p contributed little if anything to Sum1-1p silencing at *HMR*, whereas ORC was of central importance, it would appear that in the absence of bound ORC at telomeres, Sum1-1p would have no way of associating with the telomeres to form heterochromatin in a *sir2Δ* mutant. In contrast, Sum1-1p could interact with Sir proteins at telomeres to enhance silencing.

The slow growth phenotype of *SUM1-1* strains and the dependence of this slow growth on ORC and *HST1* suggested that Sum1-1p-mediated silencing might occur at other loci. Genes located near ORC-binding sites are good candidates to be repressed by Sum1-1p, because the presence of ORC appears to be required for association of Sum1-1p with a locus. However, only a subset of the ~400 ORC-binding sites at origins of replication could be sites of Sum1-1p-mediated repression because the cells are viable. Therefore, some origins must have additional features that predispose them to be silenced.

Mechanistic links between repression and silencing

This study points to similarities in the mechanisms of promoter-local repression and long-range silencing. Sum1p is a repressor that acts directly at the *SMK1*, *SPR3*, and other promoters, and its effect does not spread to a nearby, divergently transcribed gene (Xie et al. 1999). On the other hand, Sum1-1p causes silencing at a distance through the formation of heterochromatin. Thus, two nearly identical proteins act through mechanisms that have sometimes been considered completely different. This narrow margin separating repression from silencing is perhaps the most striking finding of this study. Other clear similarities among repression and silencing mechanisms are emerging. For example, Ssn6-Tup1 associates with histone tails and forms a scaffold along nucleosomes (Edmondson et al. 1996; Ducker and Simpson 2000), much as the Sir proteins are thought to do (Hecht et al. 1996). Additionally, Rpd3-Sin3 repression, Ssn6-Tup1 repression, Sir-mediated silencing, and Sum1-1p-mediated silencing all use histone deacetylases recruited by DNA-binding proteins.

The chromosomal context in which the Sum1 protein finds itself may determine its range of action. For example, the presence or absence of nearby boundary elements could determine the extent of spreading. Perhaps genes like *SMK1* have boundary elements that limit the extent of deacetylation. Alternatively, elements like the *HMR-E* silencer may have features that promote the binding and subsequent movement or spreading of Sum1-1p and Hst1p.

Materials and methods*Plasmids and epitope tagging*

Plasmids pRS316, pRS412, and pRS416 (Sikorski and Hieter 1989) and pJR759, pJR760, pJR891, pJR900, and pJR901 (Laurenson and Rine 1991) were described previously. Table 1 lists plasmids generated for this study. *SUM1-1* and *SUM1* plasmids were derived from plasmids DMC283 and DMC326, respectively (Chi and Shore 1996). To generate myc-tagged alleles of *SUM1* or *SUM1-1*, the first three codons of *SUM1* were altered by site-specific mutagenesis to ATGCCTAGG, changing amino acids 2,3 from S,E to P,R, and creating an *AvrII* site at the 5' end of the ORF. A multimerized myc tag flanked by *AvrII* sites was ligated into this site. To generate pJR2291–pJR2294, the coding region was excised with *ApaI* and *SacI* and ligated into these sites of pRS412. *HST1* plasmids were derived from pLP316

(Brachmann et al. 1995). The *SacI* fragment containing *HST1* was ligated into the *SacI* site of pRS416 or pRS406. To generate an HA-tagged allele of *HST1*, the last three codons of *HST1* were altered by site-specific mutagenesis to CACCGGTAA, changing the last two amino acids from Q,Q to H,R, and creating an *AgeI* site at the 3' end of the ORF. A multimerized HA tag flanked by *AgeI* sites was ligated into this site.

Yeast strains and construction

Strains used in this study were all derived from W303-1a (Table 2). The *hst1Δ::KanMX* null allele was generated by one-step gene conversion (Rothstein 1983), completely deleting the *HST1* ORF. The *KanMX* gene was amplified from plasmid pFA6a-KanMX4 (Wach et al. 1994) by use of primers 5'-TACGAACACTTCTCTTTTGTGTTTGTGAGAAAAACGTACGCTGCAGGTCGAC-3' and 5'-GCGGTATACCTATTTTACTCCCCCTTCTGTGTTTCTTCATCGATGAATTCGAGCTCG-3', in which the underlined portion was complementary to the plasmid and the remainder was complementary to the yeast sequence flanking *HST1*. The PCR product was transformed into yeast by use of the modified LiOAC method (Schiestl and Gietz 1989) and geneticin-resistant colonies were selected. Gene disruption was confirmed by use of PCR analysis and DNA blotting. The *sum1Δ::LEU2* allele was created similarly, except the *LEU2* gene was amplified from plasmid pRS415 by use of primers 5'-ATCAAACGAAAGTTTCATACATAATTAACAAAATTCGTTTGTGTCGGGGGATTGTAAGTACGAGTGCACC-3' and 5'-TTTTTCATTTT TTTATCTATTCTCGAAACTGCCCAACGTACGGACCAGCCTGTGCGGTATTTACACACCG-3'. The *7myc-SUM1-1* and *3myc-SUM1* alleles were generated by transforming DNA containing the complete ORF with tag plus 5' and 3' flanking sequences (from plasmids pJR2295 and pJR2296) into a *sum1Δ::URA3* strain and selecting for 5-FOA-resistant colonies. The *orc5-1* and *orc2-1* alleles (Foss et al. 1993; Loo et al. 1995) and *LEU2::ORC1(Q236H,I237M)* and *LEU2::orc1Δ1-235* were described previously (Bell et al. 1995). The *sir1Δ::LEU2*, *sir2Δ::TRP1*, *sir2Δ::HIS3*, *sir3Δ::LEU2*, and *sir4Δ::HIS3* alleles were complete deletions of the ORFs generated by one-step gene conversion.

Mating assays

For mating assays, cells were patched on solid YPD medium (or minimal medium without uracil to retain plasmids) and grown 8–16 h before replica plating to mating-type tester lawns (JRY2726 unless otherwise indicated) on minimal medium. Plates were incubated at 30°C unless otherwise indicated and photographed after 2 or 3 d.

Immunoprecipitation techniques

For protein coimmunoprecipitation, 30 OD units cells were lysed by vortexing with glass beads in 100 μL of ice-cold Co-IP lysis buffer (50 mM Tris at pH 7.5, 5 mM EDTA, 250 mM NaCl, 0.1% NP40, 1.5 mM DTT, 0.1 mM TPCK, 1 mM PMSE, 5 μg/mL chymostatin, 2 μg/mL pepstatin A, 1 mM benzamide, and 1× complete protease inhibitors; Boehringer-Mannheim). The volume was then increased to 500 μL with Co-IP lysis buffer, and the lysate was drawn off the beads and then spun 10 min in a microfuge at 4°. The supernatant fraction was collected, and 100 μL was used in a final volume of 400 μL Co-IP lysis buffer for each immunoprecipitation reaction. The samples were incubated at 4°C with 5 μL of rabbit polyclonal IgG antibodies (Upstate Biotech) for 3 h and then 60 μL of Protein A agarose

Table 1. *Plasmids generated for this study*

Plasmid	Description
pJR2288	pRS416 <i>HST1</i>
pJR2289	pRS416 <i>HST1</i> -5HA
pJR2291	pRS412 3myc- <i>SUM1-1</i>
pJR2292	pRS412 3myc- <i>SUM1</i>
pJR2293	pRS412 <i>SUM1-1</i>
pJR2294	pRS412 <i>SUM1</i>
pJR2295	pRS415 7myc- <i>SUM1-1</i>
pJR2296	pRS415 3myc- <i>SUM1</i>

Table 2. *Strains used in this study*

Strain	Genotype	Source ^a
W303-1b	<i>MATα ade2-1 can1-100 his3-11 leu2-3, 112 trp1-1 ura3-1</i>	R. Rothstein
MC89	W303 <i>MATα SUM1-1</i>	D. Shore
TD4	<i>MATα can1 his4-519 leu2-3,112 trp1 ura3-52</i>	G. Fink
JRY2726	<i>MATα his4</i>	P. Schatz
JRY4565	W303 <i>MATα sir2Δ</i>	
JRY7145	W303 <i>MATα hst1Δ::KanMX</i>	
JRY7146	W303 <i>MATα orc5-1</i>	
JRY7147	W303 <i>MATα orc2-1</i>	
JRY7148	W303 <i>MATα SUM1-1 hst1Δ::KanMX</i>	
JRY7149	W303 <i>MATα SUM1-1 orc5-1</i>	
JRY7150	W303 <i>MATα SUM1-1 orc2-1</i>	
JRY7151	W303 <i>MATα sir2Δ::TRP1 hst1Δ::KanMX sum1Δ::LEU2</i>	
JRY7152	W303 <i>MATα sir2Δ::TRP1 SUM1-1</i>	
JRY7153	W303 <i>MATα sir2Δ::HIS3 7myc-SUM1-1</i>	
JRY7154	W303 <i>MATα sir2Δ::TRP1 SUM1-1 hst1Δ::KanMX</i>	
JRY7156	W303 <i>MATα sir2Δ::TRP1 hst1Δ::KanMX</i>	
JRY7157	W303 <i>MATα sir2Δ::TRP1 SUM1-1 orc5-1</i>	
JRY7158	W303 <i>MATα sir2Δ::HIS3 7myc-SUM1-1 orc5-1</i>	
JRY7159	W303 <i>MATα sirΔ::TRP1 SUM1-1 orc2-1</i>	
JRY7160	W303 <i>MATα sir3Δ::LEU2</i>	
JRY7161	W303 <i>MATα sir3Δ::LEU2 SUM1-1</i>	
JRY7162	W303 <i>MATα sir3Δ::LEU2 SUM1-1 hst1Δ::KanMX</i>	
JRY7163	W303 <i>MATα sir3Δ::LEU2 hst1Δ::KanMX</i>	
JRY7164	W303 <i>MATα sir3Δ::LEU2 SUM1-1 orc5-1</i>	
JRY7165	W303 <i>MATα sir4Δ::HIS3</i>	
JRY7166	W303 <i>MATα sir4Δ::HIS3 SUM1-1</i>	
JRY7167	W303 <i>MATα sir4Δ::HIS3 SUM1-1 hst1Δ::KanMX</i>	
JRY7168	W303 <i>MATα sir4Δ::HIS3 hst1Δ::KanMX</i>	
JRY7169	W303 <i>MATα sir4Δ::HIS3 SUM1-1 orc5-1</i>	
JRY7170	W303 <i>MATα sum1Δ::LEU2</i>	
JRY7172	W303 <i>MATα sir2Δ::HIS3 3myc-SUM1</i>	
JRY7173	W303 <i>MATα sir1Δ::LEU2 SUM1-1</i>	
JRY7174	W303 <i>MATα sir1Δ::LEU2 sir2Δ::TRP1 SUM1-1</i>	
JRY7176	W303 <i>MATα sir2Δ::HIS3 7myc-SUM1-1 orclΔ::TRP1 LEU2::orclΔ1-235</i>	
JRY7177	W303 <i>MATα sir2Δ::HIS3 7myc-SUM1-1 orclΔ::TRP1 LEU2::ORC1(Q236H,I237M)</i>	
JRY7178	W303 <i>MATα sir2Δ::HIS3 7myc-SUM1-1 hst1Δ::KanMX</i>	

^aUnless otherwise indicated, all strains were produced for this study.

beads for 1 h. The beads were washed twice in Co-IP lysis buffer, and proteins were eluted in SDS-PAGE sample buffer. Samples were electrophoretically fractionated on 7.5% polyacrylamide-SDS gels, transferred to membranes, and probed by use of mouse monoclonal antibodies (BAbCO).

For chromatin immunoprecipitations, cells were cross-linked with 1% formaldehyde for 20 min; 50 OD units cells were lysed by vortexing with glass beads in 100 μ L ice-cold ChIP lysis buffer (50 mM HEPES at pH 7.5, 1 mM EDTA, 140 mM NaCl, 1% Triton X-100, 0.1% sodium deoxycholate, 0.1 mM TPCK, 1 mM PMSF, 5 μ g/mL chymostatin, 2 μ g/mL pepstatin A, 1 mM benzamidine, and 1 $\times$ complete protease inhibitors; Boehringer Mannheim). The volume was then increased to 750 μ L with ChIP lysis buffer, and the lysate was drawn off the beads, sonicated four times for 10 sec each at setting 2 on a Branson Sonifier 450, and spun 10 min in a microfuge. The supernatant fraction was collected, and 150 μ L was used in a final volume of 400 μ L of ChIP lysis buffer for each immunoprecipitation reaction. Samples were first pre-cleared for 30 min with 30 μ L of a 50% slurry of Protein A agarose beads in 0.2 mg/mL salmon sperm DNA and 0.5 mg/mL BSA (Upstate Biotech). The samples were then incubated overnight with 4 μ L of antibody. Antibodies were 1 mg/mL rabbit polyclonal IgG anti-Myc tag (residues 409–

420), 2–10 mg/mL ChIP grade rabbit polyclonal IgG anti-acetyl-histone H4, 1 mg/mL rabbit polyclonal IgG anti-acetyl-histone H4, or 1 mg/mL rabbit polyclonal IgG anti-acetyl-histone H3 (Upstate Biotech). Samples were incubated with 60 μ L beads for 1 h, the beads were washed (wash 1: 150 mM NaCl, 20 mM Tris at pH 8.1, 2 mM EDTA, 1% Triton X-100, 0.1% SDS; wash 2: 500 mM NaCl, 20 mM Tris at pH 8.1, 2 mM EDTA, 1% Triton X-100, 0.1% SDS; wash 3: 10 mM Tris at pH 8.1, 1 mM EDTA, 0.25 M LiCl, 1% sodium deoxycholate, 1% NP40; washes 4 and 5: TE), and DNA was eluted in 500 μ L of 0.1 M NaHCO₃, 1% SDS. Cross-links were reversed by overnight incubation at 65°C, and the DNA was isolated by use of standard procedures.

PCR analysis of immunoprecipitated DNA

PCR reactions were performed using Amplitaq Gold enzyme (Perkin Elmer), with 3.5 mM MgCl₂, 0.2 mM dNTPs, and 0.5 μ M primers. After 13 min at 94°C to activate the enzyme, samples were cycled 28 times for 15 sec at 94°C, 20 sec at 55°C, and 2.5 min at 68°C. The oligonucleotides used are described in Table 3.

Table 3. Primers for PCR amplification of ChIP DNA

Region	Primer 1	Primer 2
<i>SSC1</i>	gcttcggcccggttcca	cagcaagcatcttgggtcg
<i>HMR-E</i>	ctaatacgcatttcttttcgtccac	taacaaaaaccaggagtacctgcgc
<i>HMR-I</i>	ccaacatttctgtatatggc g	cctaccacattatcaatccttgc
<i>HMR-X/Ya</i>	taccaacccatcgccg	tccgccatactacaaatcatc
<i>HMR-Ya/ZI</i>	gtggcattactccacttcaagtaag	caagagcaagacgatgggg
<i>3'HMR</i>	caacatgggttccaaagcac	gcagcttactcccaagagtgc
<i>HML-E</i>	ggatggatctagggttttatgcc	tttggcccccgaatcg
<i>HML-I</i>	ccagctgagtaactaactctcatgg	gctgttacggagatgcaag c
<i>MAT</i>	ggctatacgggacggatgac	gttcttagcttgtaccagaggaagc
<i>ACT1</i>	ctttttcttccacgtcctcttgc	tgggatgggtgcaagcg
<i>SMK1</i>	ggtgaccattgcgactgtgc	cataggctccttgcccagg
<i>SPR3</i>	gacggcgactgtccttggg	gcttcggcccggttcca

RNA blots

RNA was isolated from log-phase cells (Schmitt et al. 1990) and separated electrophoretically on formaldehyde agarose gels and blotted (Sambrook et al. 1989). DNA fragments for probes were generated by PCR amplification of genomic DNA (W303) by use of the primers below, and labeled with [α - ^{32} P]dCTP using the Multiprime Labeling System (Amersham). For *a1* probe, GGAT GATATTTGTAGTATGGCGG and CTGTTGCGGAAAGCT GAAAC were used. For *SMK1* probe, GCGACGAGGCGTG AGGGTAG and CATAGGCTCCCTTGCCCAGG were used. For *SPR3* probe, CCAAGGACAGTACGCCGTC and AATGT CAACAGCGCCCC were used. For *ACT1*, the GenePair primer set from Research Genetics was used.

Acknowledgments

We thank Andrew Vershon and Michael Pierce for stimulating discussion and communicating unpublished results, David Shore, Lorraine Pillus, and Stephen Bell for strains and plasmids, Charity Kirk for technical assistance, Paul Kaufmann, Ann Kirchmaier, and Peter Garber for critical reading of this manuscript, and members of the laboratory for comments and suggestions. This work was supported by a postdoctoral fellowship from the Damon Runyon-Walter Winchell Cancer Research Fund (L.N.R.) and a grant from the National Institutes of Health (GM31105, J.R.). Core support was provided by a National Institute of Environmental Health Sciences Mutagenesis Center Grant.

The publication costs of this article were defrayed in part by payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 USC section 1734 solely to indicate this fact.

References

- Ahringer, J. 2000. NuRD and SIN3 histone deacetylase complexes in development. *Trends Genet.* **16**: 351–356.
- Ayer, D. 1999. Histone deacetylases: Transcriptional repression with SINers and NuRDs. *Trends Cell. Biol.* **9**: 193–198.
- Bell, S., Mitchell, J., Leber, J., Kobayashi, R., and Stillman, B. 1995. The multidomain structure of Orc1p reveals similarity to regulators of DNA replication and transcriptional silencing. *Cell* **83**: 563–568.
- Bi, X., Braunstein, M., Shei, G., and Broach, J. 1999. The yeast HML I silencer defines a heterochromatin domain boundary

by directional establishment of silencing. *Proc. Natl. Acad. Sci.* **96**: 11934–11939.

- Brachmann, C., Sherman, J., Devine, S., Cameron, E., Pillus, L., and Boeke, J. 1995. The *SIR2* gene family, conserved from bacteria to humans, functions in silencing, cell cycle progression, and chromosome stability. *Genes & Dev.* **9**: 2888–2902.
- Braunstein, M., Rose, A., Holmes, S., Allis, C., and Broach, J. 1993. Transcriptional silencing in yeast is associated with reduced nucleosome acetylation. *Genes & Dev.* **7**: 592–604.
- Chi, M. and Shore, D. 1996. *SUM1-1*, a dominant suppressor of *SIR* mutations in *Saccharomyces cerevisiae*, increases transcriptional silencing at telomeres and *HM* mating-type loci and decreases chromosome stability. *Mol. Cell. Biol.* **16**: 4281–4294.
- DeRisi, J., Iyer, V., and Brown, P. 1997. Exploring the metabolic and genetic control of gene expression on a genomic scale. *Science* **278**: 680–686.
- Donze, D., Adams, C.R., Rine, J., and Kamakaka, R.T. 1999. The boundaries of the silenced *HMR* domain in *Saccharomyces cerevisiae*. *Genes & Dev.* **13**: 698–708.
- Dorland, S., Deegenars, M., and Stillman, D. 2000. Roles for the *Saccharomyces cerevisiae* *SDS3*, *CBK1* and *HYM1* genes in transcriptional repression by *SIN3*. *Genetics* **154**: 573–586.
- Ducker, C. and Simpson, R. 2000. The organized chromatin domain of the repressed yeast a cell-specific gene *STE6* contains two molecules of the corepressor Tup1p per nucleosome. *EMBO J.* **19**: 400–409.
- Edmondson, D., Smith, M., and Roth, S. 1996. Repression domain of the yeast global repressor Tup1 interacts directly with histones H3 and H4. *Genes & Dev.* **10**: 1247–1259.
- Foss, M., McNally, F., Laurensen, P., and Rine, J. 1993. Origin recognition complex (ORC) in transcriptional silencing and DNA replication in *S. cerevisiae*. *Science* **262**: 1838–1844.
- Fox, C.A., Ehrenhofer-Murray, A.E., Loo, S., and Rine, J. 1997. The origin recognition complex, *SIR1*, and the S phase requirement for silencing. *Science* **276**: 1547–1551.
- Gardner, K., Rine, J., and Fox, C. 1999. A region of the Sir1 protein dedicated to recognition of a silencer and required for interaction with the Orc1 protein in *Saccharomyces cerevisiae*. *Genetics* **151**: 31–44.
- Gotta, M., Strahl-Bolsinger, S., Renauld, H., Laroche, T., Kennedy, B., Grunstein, M., and Gasser, S. 1997. Localization of Sir2p: The nucleolus as a compartment for silent information regulators. *EMBO J.* **16**: 3243–3255.
- Hecht, A., Laroche, T., Strahl-Bolsinger, S., Gasser, S., and Grunstein, M. 1995. Histone H3 and H4 N-termini interact

- with SIR3 and SIR4 proteins: A molecular model for the formation of heterochromatin in yeast. *Cell* **80**: 583–592.
- Hecht, A., Strahl-Bolsinger, S., and Grunstein, M. 1996. Spreading of transcriptional repressor SIR3 from telomeric heterochromatin. *Nature* **383**: 92–96.
- Imai, S., Armstrong, C., Kaeberlein, M., and Guarente, L. 2000. Transcriptional silencing and longevity protein Sir2 is an NAD-dependent histone deacetylase. *Nature* **403**: 795–800.
- Kadosh, D. and Struhl, K. 1997. Repression by Ume6 involves recruitment of a complex containing Sin3 corepressor and Rpd3 histone deacetylase to target promoters. *Cell* **89**: 365–371.
- . 1998. Targeted recruitment of the Sin3-Rpd3 histone deacetylase complex generates a highly localized domain of repressed chromatin in vivo. *Mol. Cell. Biol.* **18**: 5121–5127.
- Klar, A., Kakar, S., Ivy, J., Hicks, J., Livi, G., and Miglio, L. 1985. *SUM1*, an apparent positive regulator of the cryptic mating-type loci in *Saccharomyces cerevisiae*. *Genetics* **111**: 745–758.
- Landry, J., Sutton, A., Tafrov, S., Heller, R., Stebbins, J., Pillus, L., and Sternglanz, R. 2000. The silencing protein SIR2 and its homologs are NAD-dependent protein deacetylases. *Proc. Natl. Acad. Sci.* **97**: 5807–5811.
- Laurenson, P. and Rine, J. 1991. *SUM1-1*: A suppressor of silencing defects in *Saccharomyces cerevisiae*. *Genetics* **129**: 685–696.
- Lee, M., Chatterjee, S., and Struhl, K. 2000. Genetic analysis of the role of Pol II holoenzyme components in repression by the Cyc8–Tup1 corepressor in yeast. *Genetics* **155**: 1535–1542.
- Livi, G., Hicks, J., and Klar, A. 1990. The *sum1-1* mutation affects silent mating-type gene transcription in *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* **10**: 409–412.
- Loo, S., Fox, C.A., Rine, J., Kobayashi, R., Stillman, B., and Bell, S. 1995. The origin recognition complex in silencing, cell cycle progression, and DNA replication. *Mol. Biol. Cell* **6**: 741–756.
- McNally, F. and Rine, J. 1991. A synthetic silencer mediates SIR-dependent functions in *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* **11**: 5648–5659.
- Moretti, P., Freeman, K., Coodly, L., and Shore, D. 1994. Evidence that a complex of SIR proteins interacts with the silencer and telomere-binding protein RAP1. *Genes & Dev.* **8**: 2257–2269.
- Pak, D., Pflumm, M., Chesnokov, I., Huang, D., Kellum, R., Marr, J., Romanowski, P., and Botchan, M. 1997. Association of the origin recognition complex with heterochromatin and HP1 in higher eukaryotes. *Cell* **91**: 311–323.
- Rothstein, R. 1983. One-step gene disruption in yeast. *Methods Enzymol.* **101**: 202–211.
- Rundlett, S., Carmen, A., Suka, N., Turner, B., and Grunstein, M. 1998. Transcriptional repression by UME6 involves deacetylation of lysine 5 of histone H4 by RPD3. *Nature* **392**: 831–835.
- Sambrook, J., Fritsch, E., and Maniatis, T. 1989. *Molecular cloning: A laboratory manual*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Schiestl, R. and Gietz, R. 1989. High efficiency transformation of intact yeast cells using single stranded nucleic acids as a carrier. *Curr. Genet.* **16**: 339–346.
- Schmitt, M., Brown, T., and Trumpower, B. 1990. A rapid and simple method for preparation of RNA from *Saccharomyces cerevisiae*. *Nucleic Acids Res.* **18**: 3091–3092.
- Sikorski, R. and Hieter, P. 1989. A system of shuttle vectors and yeast host strains designed for efficient manipulation of DNA in *Saccharomyces cerevisiae*. *Genetics* **122**: 19–27.
- Smith, J., Brachmann, C., Celic, I., Kenna, M., Muhammad, S., Starai, V., Avalos, J., Escalante-Semerena, J., Grubmeyer, C., Wolberger, C., et al. J. 2000. A phylogenetically conserved NAD⁺-dependent protein deacetylase activity in the Sir2 protein family. *Proc. Natl. Acad. Sci.* **97**: 6658–6663.
- Triolo, T. and Sternglanz, R. 1996. Role of interactions between the origin recognition complex and SIR1 in transcriptional silencing. *Nature* **381**: 251–253.
- Wach, A., Brachat, A., Pohlmann, R., and Philippsen, P. 1994. New heterologous modules for classical or PCR-based gene disruptions in *Saccharomyces cerevisiae*. *Yeast* **10**: 1793–1808.
- Watson, A., Edmondson, D., Bone, J., Mukai, Y., Yu, Y., Du, W., Stillman, D., and Roth, S. 2000. Ssn6–Tup1 interacts with class I histone deacetylases required for repression. *Genes & Dev.* **14**: 2737–2744.
- Xie, J., Pierce, M., Gailus-Durner, V., Wagner, M., Winter, E., and Vershon, A. 1999. Sum1 and Hst1 repress middle sporulation-specific gene expression during mitosis in *Saccharomyces cerevisiae*. *EMBO J.* **18**: 6448–6454.
- Zhang, Y., Sun, Z., Iratni, R., Erdjument-Bromage, H., Tempst, P., Hampsey, M., and Reinberg, D. 1998. SAP30, a novel protein conserved between human and yeast, is a component of a histone deacetylase complex. *Mol. Cell* **1**: 1021–1031.

Conversion of a gene-specific repressor to a regional silencer

Laura N. Rusché and Jasper Rine

Genes Dev. 2001, **15**:

Access the most recent version at doi:[10.1101/gad.873601](https://doi.org/10.1101/gad.873601)

References

This article cites 41 articles, 26 of which can be accessed free at:

<http://genesdev.cshlp.org/content/15/8/955.full.html#ref-list-1>

License

Email Alerting Service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).