RESEARCH COMMUNICATION

The nucleotides responsible for the direct physical contact between the chromatin insulator protein CTCF and the *H19* imprinting control region manifest parent of origin-specific long-distance insulation and methylation-free domains

Vinod Pant,¹ Piero Mariano,¹ Chandrasekhar Kanduri,¹ Anita Mattsson,¹ Victor Lobanenko,² Rainer Heuchel,³ and Rolf Ohlsson^{1,4}

¹Department of Development & Genetics, Evolution Biology Centre, Uppsala University, S-752 36 Uppsala, Sweden;

²Laboratory of Immunopathology, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, Maryland 20892-0760, USA; ³Ludwig Institute for Cancer Research, Biomedical Center, S-751 24 Uppsala, Sweden

The repression of the maternally inherited *Igf2* allele has been proposed to depend on a methylation-sensitive chromatin insulator organized by the 11 zinc finger protein CTCF at the *H19* imprinting control region (ICR). Here we document that point mutations of the nucleotides in physical contact with CTCF within the endogenous *H19* ICR lead to loss of CTCF binding and *Igf2* imprinting only when passaged through the female germline. This effect is accompanied by a significant loss of methylation protection of the maternally derived *H19* ICR. Because CTCF interacts with other imprinting control regions, it emerges as a central factor responsible for interpreting and propagating gamete-derived epigenetic marks and for organizing epigenetically controlled expression domains.

Received November 15, 2002; revised version accepted January 8, 2003.

The manifestation of genomic imprinting involves the translation of gametic marks into parent of origin-dependent gene expression patterns (Bartolomei and Tilghman 1997; Horsthemke et al. 1999). The neighboring *IGF2* and *H19* genes emerge as paradigms of genomic imprinting, because their expression is monoallelic from opposite parental alleles and governed by shared enhancers (Bartolomei and Tilghman 1997; Horsthemke et al.

1999). The repression of the maternal *IGF2* and paternal *H19* alleles depends on a differentially methylated imprinting control region (ICR) in the 5' region of the *H19* gene (Olek and Walter 1997; Kaffer et al. 2000). The proposal that the complex between the *H19* ICR and the 11 zinc finger protein CTCF organizes a CpG methylation-sensitive insulation of the maternal *Igf2* allele (Bell and Felsenfeld 2000; Hark et al. 2000; Kanduri et al. 2000a,b; Bell et al. 2001; Ohlsson et al. 2001) is supported by the observations that CTCF interacts with only the maternal *H19* ICR allele (Kanduri et al. 2000b) and that the chromatin insulator function is regulated by CpG methylation (Holmgren et al. 2001).

To directly prove this proposal, we mutated sequences within the *H19* ICR, which we have previously shown to be in direct physical contact with CTCF. Our results document that the CTCF-*H19* ICR complex is vital in the manifestation and propagation of gametic marks.

Results and Discussion

To directly demonstrate the function of the CTCF target sites within the *H19* ICR in association with the establishment and manifestation of the imprinting phenomenon, we changed the sequence GTGG to ATAT in three of the four CTCF target sites within the CGCG(T/G)GGTGGCAG-core motif. This sequence change deletes essential contact points for the CTCF while preserving the CpGs responsible for the methylation-sensitive portion of the CTCF target sites (Kanduri et al. 2000b). The remaining second CTCF target site was ignored, because it does not, in contrast to the other target sites, display any marked in vivo footprint or nuclease hypersensitivity (Kanduri et al. 2000a; Szabó et al. 2000). Following electroporation of the targeting construct into ES cells and exploiting the fact that the mutations introduced an *EcoRV* site in each of the three mutated target sites (Kanduri et al. 2000b), we were able to identify one cell clone, #142, which had the endogenous *H19* ICR replaced with all three mutated CTCF sites (Fig. 1A). The retained neomycin gene, which was flanked with *loxP* sites, was deleted by mating homozygous 142 mice with mice carrying a *Cre* transgene under the control of the β -actin promoter (Lewandoski and Martin 1997). Southern blot analysis confirmed the deletion of the neomycin gene in the offspring of these crosses to generate the 142* substrain (Fig. 1B).

To verify that the mutations affected the association between CTCF and the maternal allele of the *H19* ICR in vivo, we performed chromatin immunoprecipitation (ChIP) analyses of formaldehyde cross-linked DNA-protein complexes from E14.5 embryos, followed by PCR amplification. As accounted for above, *EcoRV* restriction of the amplified fragments encompassing CTCF target site #3 allowed the discrimination between the wild-type and mutated *H19* ICR alleles. Figure 2 shows that the wild-type maternal *H19* ICR allele was pulled down by the CTCF antibody in agreement with a previous report (Kanduri et al. 2000b). However, the mutated *H19* ICR allele could not be similarly detected upon either paternal or maternal inheritance. This difference did not depend on variation in quality between the formaldehyde-fixed protein-DNA complexes, because sequences included in the *Snrpn* imprinting box (Shemer et al.

[Keywords: Genomic imprinting; *Igf2*; chromatin insulator; CTCF; DNA methylation]

*Corresponding author.

E-MAIL Rolf.Ohlsson@ebc.uu.se; FAX 46-18-4712683.

Article and publication are at <http://www.genesdev.org/cgi/doi/10.1101/gad.254903>.

Figure 1. Schematic illustration of the knock-in strategy. (A) Recombination between targeting construct and endogenous H19 ICR as confirmed by Southern blot analysis. The recombination in the 142 clone was assessed by using a probe that is positioned 5' of the sequence covered in the targeting construct. The properly recombined insert generated a larger size fragment due to the extra sequence information provided by the neomycin gene. All three mutated CTCF target sites (cerise boxes) replaced the endogenous sequences in the 142 clone as determined by EcoRV (EV) digestion and Southern blot analysis. The white box identifies CTCF target site #2, which was not mutated. The identities of the fragments in the Southern blot image are depicted on the far right of the image. (B) Deletion of the neomycin gene by breeding the 142 strain with mice harboring a β -actin promoter-driven Cre recombinase gene to generate the 142* substrain. The mutated allele is now indistinguishable from the wild-type allele when digested with BamHI (B) and DraI (D). See the text for additional information.

2000), interacted with CTCF in both reciprocal crosses (Fig. 2). We conclude that point mutations of the H19 ICR CTCF target sites #1, #3, and #4 lead to complete failure of CTCF binding in vivo, confirming our initial anticipation that site #2 does not interact with CTCF in vivo.

We next examined whether or not the mutations af-

Figure 2. CTCF occupancy on wild-type and mutated H19 ICR alleles. Chromatin immunopurification analyses were performed on dispersed and formaldehyde cross-linked E14.5 embryos with the mutated H19 ICR allele inherited paternally or maternally. The PCR primers spanned CTCF target site #3, which displayed an EcoRV site only in the mutated H19 ICR allele, as indicated in the image. As an internal control, sequences of the *Snrpn* imprinting box were amplified from the same immunopurified DNA samples. In both instances, the sense primers were end-labeled with 32 P. See the text for additional information.

fected the imprinted state of the *Igf2* gene by crossing 142* mice with SD7 mice, which were derived from a congenic strain carrying the distal end of chromosome 7 of *Mus musculus spretus* on a *Mus musculus domesticus* background (Dean et al. 1998). A polymorphic *Bsa*AI site in the exon 4 of the *Igf2* locus allowed the discrimination of allelic expression patterns in the offspring of these two strains (Dean et al. 1998). Figure 3 shows reverse transcriptase PCR (RT-PCR) analyses of a large variety of tissue types documenting that neonatal offspring from 142* x SD7 crosses (in the order female to male) robustly expressed *Igf2* from both parental alleles. This loss of *Igf2* imprinting was accompanied by an 11% gain of weight in neonates when compared to neonates derived

Figure 3. RT-PCR analyses of *Igf2* and *H19* expression patterns in various organs dissected from different neonatal offspring (labeled #1 and #2) of reciprocal crosses between 142* and SD7 mice. These assays exploited species-specific *Bsa*AI and *Bgl*I polymorphisms in exon 4 of *Igf2* (Dean et al. 1998) and exon 5 of *H19* (Sasaki et al. 1995), respectively, as schematically indicated in the figure. The crosses are presented in the order female to male. See the text for additional information.

from the reciprocal SD7 x 142* crosses (in the order female to male), which display a normal *Igf2* imprinting status (Fig. 3; data not shown). We conclude that the contact points within CTCF sites #1, #3, and #4 in the H19 ICR are essential for maintaining the repressed status of the maternal *Igf2* allele in most cell types of both mesodermal and endodermal lineages of the developing conceptus. In each of these instances, *H19* imprinting appeared normal, although we noted a slight activation of the paternal H19 allele in offspring inheriting the mutated ICR allele paternally (Fig. 3).

Our strategy to keep the crucial CpGs intact in the mutated CTCF target sites allowed us to directly assess

if the methylation privilege status was lost as a consequence of the inability to interact with CTCF. CpG methylation analyses of the parental alleles, which were discriminated by using a polymorphic *Bst*XI site that restricts 142* *H19* ICR, but not the SD7 allele (Fig. 4A), was initially performed by restricting the DNA with the methylation-sensitive enzyme *Hha*I followed by Southern blot analysis. Although it is well-established that the maternal *H19* ICR allele is normally unmethylated (Olek and Walter 1997), Figure 4A shows that the mutation of the contact sequences for the CTCF sites resulted in significant de novo methylation, when maternally inherited. This observation was confirmed by bisulphite methylation analyses of the CTCF target site #3 of the mutated *H19* ICR allele in heart and liver. Figure 4B shows that the normally maintained unmethylated status at this CTCF target site when maternally inherited (Olek and Walter 1997; Liang et al. 2000) is practically lost when passed through the female germline. Interestingly, paternal transmission of the mutated *H19* ICR also led to partial methylation of the maternally inherited wild-type *H19* ICR allele in some tissues, such as heart and liver (Fig. 4A). Although this pattern of de novo methylation must be limited due to absence of expression of the maternal *Igf2* allele in the very same tissue specimens (Fig. 3) and shows variation among different conceptuses (data not shown), it suggests the existence of a *trans*-sensing mechanism. This deduction converges

with earlier proposals of *trans*-sensing within the *Igf2*/*H19* imprinting domain (LaSalle and Lalande 1996; Forne et al. 1997).

Our results document the dual feature of the CTCF–methylation link. In addition to the ability of CTCF to interpret the methylation status of its target sites within the *H19* ICR, CTCF also maintains a methylation-free domain at and around its target sites. Although it is likely that this latter feature involves protection by the CTCF–*H19* ICR complex towards the de novo methylation machinery, it is probable that many other neighboring protein–DNA complexes play a similar role. This deduction is supported by our observation that the mutated, maternal *H19* ICR allele remains methylation-free in the unfloxed 142 strain containing the transcriptionally active neomycin gene just 3' of the *H19* ICR (data not shown). Given that the de novo methylation of the mutated *H19* ICR in the floxed 142* strain is not complete, it is likely that additional factors interact with other linker regions between the positioned nucleosomes (Kanduri et al. 2002) to jointly maintain the methylation-free domain of the maternally inherited *H19* ICR allele with CTCF. One such candidate factor is Oct-1, which has been reported to interact with the *H19* ICR between CTCF target sites #2 and #3 to provide partial protection against de novo methylation (Hori et al. 2002).

This report documents that the CTCF target sites manifest not only the repressed status of the maternally inherited *Igf2* allele, but also the methylation-free domain of the maternal *H19* ICR allele in somatic cells. This latter issue is of particular interest with respect to the acquisition of the methylation mark during male germline development, because it implies that CTCF must be down-regulated and removed from the maternal *H19* ICR allele in adult testis prior to de novo methylation. Although CTCF is indeed down-regulated during differentiation of spermatogonia into spermatocytes (J. Whitehead, L. Liu, C. Kanduri, V. Pant, M. Lezcano, W.-Q. Yu, A. Kerjean, M. Parvinen, A. Paldi, E. Klenova, V. Lobanenko, and R. Ohlsson, unpubl.), this would seemingly be at odds with the vital role for CTCF in cell survival and development (G.N. Filippova, J. Whitehead, S. Fagerlie, S. Vatolin, K. Foley, D. Loukinov, E.M. Pugacheva, J.E. Ulmer, J.M. Moore, Y.J. Hu, E.M. Klenova, C. Kemp, S.J. Collins, P.E. Neiman, R. Ohlsson, and V.V. Lobanenko, unpubl.). We have earlier shown that a paralog of CTCF, termed BORIS for "Brother of Regulator of Imprinted Sites," is expressed exclusively in CTCF-negative spermatocytes in adult mouse testis (Loukinov et al. 2002). Because our preliminary observation shows that BORIS replaces CTCF on *H19* ICR target sites prior to imprint acquisition of the maternally derived *H19* ICR allele (J. Whitehead, L. Liu, C. Kanduri, V. Pant, M. Lezcano, W.-Q. Yu, A. Kerjean, M. Parvinen, A. Paldi, E. Klenova, V. Lobanenko, and R. Ohlsson, unpubl.), we speculate that the CTCF → BORIS switch is essential for the elimination of the methylation-protective properties of the CTCF–*H19* ICR complex in pachytene spermatocytes.

Figure 4. CpG methylation analyses. (A) Southern blot analysis of genomic DNA extracted from various organs dissected from neonatal offspring of reciprocal crosses between 142* and SD7 mice. The parental alleles were discriminated by exploiting a polymorphic *Bst*XI site, as indicated in A. (B) Bisulphite sequencing analyses of the mutated *H19* ICR allele between –2836 to –2479 bp from the *H19* transcriptional start site, covering the mutated CTCF target site #3. The wild-type and mutant *H19* ICR alleles were discriminated on the basis of the introduced point mutations within CTCF target site #3. The tissue type and the sex of the parent transmitting the mutant allele is indicated above each scheme. Filled boxes indicate methylated CpGs and no box indicates no methylation at the positions indicated above each row.

Materials and methods

Knock-in strategy

Three of the four CTCF target sites (Fig. 1) were mutated to change the core sequence from GTGG to ATAT, as has been described (Kanduri et al. 2000b). The targeting vector was assembled in the pGEMT vector, as indicated in Figure 1. The herpes simplex virus thymidine kinase expression cassette is followed by 4.3 kb of genomic 5'-sequence (long genomic arm), the PGK-neobpA expression cassette flanked by *loxP* sites ("floxed"), ~2.6 kb of genomic 3'-sequence (including the *H19* transcriptional start site) and the pGEMT vector backbone. The vector was linearised at the singular *NdeI* site, electroporated into R1 embryonic stem cells (Nagy et al. 1993) and colonies were selected with G418 and Gancyclovir. Homologous integration events were screened by Southern blotting (see below). The neomycin gene was deleted by mating homozygous mutant mice with a transgenic mouse strain carrying the *Cre*-recombinase gene under the control of the β -actin promoter (Lewandoski and Martin 1997).

Southern analysis for confirming recombination

Genomic DNA was isolated using Wizard genomic DNA isolation kit (Promega). DNA was digested with respective enzymes, analyzed by electrophoresis on a 1% agarose gel blotted onto a Hybond-N+ membrane (Amersham Pharmacia). A 1.3-kb PCR-generated fragment (–6750 to –5406 bp with respect to the transcription start site of the mouse *H19* gene) outside the targeting vector sequence was used to test for correct genomic integration of the targeting construct. To assess the presence of all three mutated CTCF target sites, DNA was digested with *DraI*, *BamHI*, and *EcoRV* and probed with an internal probe spanning the sequence –4296 to –1759 bp upstream of the *H19* transcription start site, covering all four CTCF sites in the *H19* ICR region. For screening of the recombinant clones, genomic DNA was digested with *DraI* and *BamHI* and probed with a 5' external probe. The presence of the neomycin gene allowed the discrimination of the mutated and wild-type alleles.

Chromatin immunopurification analyses

Dispersed embryonic cells were formaldehyde cross-linked (in 1% formaldehyde for 10 min at 37°C), and the DNA–protein complexes were immunopurified using anti-CTCF antibody and protein A 4 Fast Flow Sepharose beads (Pharmacia-Upjohn), as has been described (Kuo and Allis 1999; Kanduri et al. 2000b). To examine an association between *H19* ICR and CTCF, the immunopurified DNA was PCR amplified with primers spanning the mutated CTCF target site #3: sense primer 5'-CTCAGTGGTTCGATATATGGTTT-3'; antisense primer 5'-TGAGTCAAGTCTCTTGGTTC-3'. PCR conditions were 95°C for 3 min, 28× [94°C for 40 sec, 54°C for 40 sec, 72°C for 40 sec]. To compare the quality of the formaldehyde cross-linked samples, sequences included in the *Sn-rpn* imprinting box were amplified from the same samples, as follows: sense primer 5'-ATCCTGGATGCAAGAGCTGT-3'; antisense primer 5'-GCCGCACGTACAGTTACA-3'. PCR conditions were: 95°C for 3 min, 28× [94°C for 40 sec, 58°C for 45 sec, 72°C for 50 sec]. The amplified sequence is positioned in the first exon of *Sn-rpn* (nucleotides 3901–4162 in GenBank file AF130843). In both instances, the sense primers were end-labeled with ³²P- γ -ATP and polynucleotide kinase, as has been described (Kanduri et al. 2000b).

Allelic RT-PCR analyses

To determine the imprinting status of the *Igf2* and *H19* genes, homozygous 142* mice were mated with SD7 mice (a congenic strain carrying the distal end of chromosome 7 of *Mus musculus spretus* on a *Mus musculus domesticus* background). RNA was isolated from individual organs of 2-day-old neonatal or foetuses at E14.5 using TriPure DNA/RNA isolation reagent (Boehringer Mannheim). RNA was treated with RQ1 DNase I (Promega) to remove any contaminating genomic DNA. RT-PCR analyses were done using Qiagen one-step RT-PCR kit as per guidelines. Primers and PCR conditions were the same as previously published protocols for *Igf2* (Dean et al. 1998) and *H19* (Sasaki et al. 1995). The PCR product was purified and digested with *BsaAI* to determine the allelic origin of the *Igf2* transcripts and with *BglI* for the *H19* transcripts. Fragments were separated on 2% agarose gel and stained with ethidium bromide for visualization.

Methylation analyses

For Southern blot analysis, genomic DNA was isolated from tissues of 2-day-old neonatal mice. Fifteen micrograms of DNA were digested with

DraI, *BamHI*, and *BstXI* and with/without the methylation-sensitive restriction enzyme *HhaI*. *BstXI* has a unique site present only in the *Dom* allele but not in the SD7 allele at ~3537 bp of the *H19* transcription start site. Bisulfite treatment of DNA was carried out using an established protocol (Olek et al. 1996), with the following adaptations: DNA was restricted with an excess amount of *DraI* and *BamHI* to generate suitably small fragments containing the target sequence. One microgram of digested DNA was denatured with 0.3 M NaOH at 37°C for 15 min, then mixed with 2 volumes of 2% low-melting agarose dissolved in water. The mixture was pipetted into prechilled mineral oil to form the DNA–agarose beads. The prepared beads were then incubated with 1.2 mL 5.0 M NaHSO₃/20 mM Hydroquinone solution covered by mineral oil and incubated at 51°C for 6 h. Treated DNA beads were equilibrated with TE (1 mM EDTA, 10 mM Tris-HCl at pH 8.0), 6 × 15 min. Following desulphonation with 0.2 M NaOH, 2 × 15 min and equilibration with MilliQ water, 2 × 15 min at RT, the DNA beads were subjected to PCR amplification reactions, using the following published primers and conditions (Tremblay et al. 1997): Bhaa5t2, BHha5t, and BHha5t3 (the final PCR product covers bases –2836 to –2479 relative to the *H19* transcriptional start site). Amplified fragments were cloned into the pGEMT-Easy vector (Promega), and subsequently sequenced with the BigDye Terminator Cycle Sequencing Kit (Applied Biosystems).

Acknowledgments

We are grateful to Dr. Wolf Reik for generously providing both SD7 mice and sequence information of the *Mus musculus spretus* *H19* ICR, to Dr. Gail Martin for providing the *Cre*-transgenic mouse line, and to Dr. Ragnar Mattsson at the Lund University transgenic core facility for help with blastocyst injections. This work was supported by the Swedish Science Research Council (VR), the Swedish Cancer Research Foundation (CF), the Swedish Pediatric Cancer Foundation (BCF), and the Lundberg Foundation.

The publication costs of this article were defrayed in part by payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 USC section 1734 solely to indicate this fact.

References

- Bartolomei, M.S. and Tilghman, S.M. 1997. Genomic imprinting in mammals. *Annu. Rev. Genet.* **31**: 493–525.
- Bell, A.C. and Felsenfeld, G. 2000. Methylation of a CTCF-dependent boundary controls imprinted expression of the *Igf2* gene. *Nature* **405**: 482–485.
- Bell, A.C., West, A.G., and Felsenfeld, G. 2001. Insulators and boundaries: Versatile regulatory elements in the eukaryotic genome. *Science* **291**: 447–450.
- Dean, W., Bowden, L., Aitchison, A., Klose, J., Meneses, J., Reik, W., and Feil, R. 1998. Altered imprinted gene methylation and expression in completely ES cell-derived mouse fetuses: Association with aberrant phenotypes. *Development* **125**: 2273–2282.
- Forne, T., Oswald, J., Dean, W., Saam, J., Bailleul, B., Dandolo, L., Tilghman, S., Walter, J., and Reik, W. 1997. Loss of the maternal *H19* gene induces changes in *Igf2* methylation in both *cis* and *trans*. *Proc. Natl. Acad. Sci.* **94**: 10243–10248.
- Hark, A.T., Schoenherr, C.J., Katz, D.J., Ingram, R.S., Levorse, J.M., and Tilghman, S.M. 2000. CTCF mediates methylation-sensitive enhancer-blocking activity at the *H19/Igf2* locus. *Nature* **405**: 486–489.
- Holmgren, C., Kanduri, K., Dell, G., Ward, A., Mukhopadhyaya, R., Kanduri, M., Lobanov, V., and Ohlsson, R. 2001. CpG methylation regulates the *Igf2/H19* insulator. *Curr. Biol.* **11**: 1128–1130.
- Hori, N., Nakano, H., Takeuchi, T., Kato, H., Hamaguchi, S., Oshimura, M., and Sato, K. 2002. A dyad Oct-binding sequence functions as a maintenance sequence for the unmethylated state within the *H19/Igf2*-imprinted control region. *J. Biol. Chem.* **277**: 27960–27967.
- Horsthemke, B., Surani, M.A., James, T.C., and Ohlsson, R. 1999. The mechanisms of genomic imprinting. In *Genomic imprinting: An interdisciplinary approach* (ed. R. Ohlsson), pp. 91–118. Springer-Verlag, Berlin.
- Kaffer, C.R., Srivastava, M., Park, K.Y., Ives, E., Hsieh, S., Battle, J., Grinberg, A., Huang, S.P., and Pfeifer, K. 2000. A transcriptional insulator at the imprinted *H19/Igf2* locus. *Genes & Dev.* **14**: 1908–1919.

Pant et al.

- Kanduri, C., Holmgren, C., Franklin, G., Pilartz, M., Ullerås, E., Kanduri, M., Liu, L., Ginjala, V., Ullerås, E., Mattsson, R., et al. 2000a. The 5'-flank of the murine *H19* gene in an unusual chromatin conformation unidirectionally blocks enhancer-promoter communication. *Curr. Biol.* **10**: 449–457.
- Kanduri, C., Pant, V., Loukinov, D., Pugacheva, E., Qi, C.-F., Wolffe, A., Ohlsson, R., and Lobanenko, A. 2000b. Functional interaction of CTCF with the insulator upstream of the *H19* gene is parent of origin-specific and methylation-sensitive. *Curr. Biol.* **10**: 853–856.
- Kanduri, M., Kanduri, C., Mariano, P., Vostrov, A., Quitsche, W., Lobanenko, V., and Ohlsson, R. 2002. Multiple nucleosome positioning sites regulate the CTCF-mediated insulator function of the *H19* imprinting control region. *Mol. Cell. Biol.* **22**: 3339–3344.
- Kuo, M. and Allis, C. 1999. *In vivo* cross-linking and immunoprecipitation for studying dynamic Protein:DNA associations in a chromatin environment. *Methods* **19**: 425–433.
- LaSalle, J.M. and Lalande, M. 1996. Homologous association of oppositely imprinted chromosomal domains. *Science* **272**: 725–728.
- Lewandoski, M. and Martin, G. 1997. *Cre*-mediated chromosome loss in mice. *Nat. Genet.* **17**: 223–225.
- Liang, L., Kanduri, C., Pilartz, M., Svensson, K., Song, J., Wentzel, P., Eriksson, U., and Ohlsson, R. 2000. Dynamic readjustment of parental methylation patterns of the 5'-flank of the mouse *H19* gene during *in vitro* organogenesis. *Int. J. Dev. Biol.* **44**: 785–790.
- Loukinov, D., Pugacheva, E., Vatolin, S., Pack, S., Moon, H., Chernukhin, I., Mannan, P., Larsson, E., Kanduri, C., Vostrov, A., et al. 2002. BORIS, a novel male germline-specific protein associated with epigenetic reprogramming events, shares the same 11 Zn-finger domain with CTCF, the insulator protein involved in reading imprinting marks in the soma. *Proc. Natl. Acad. Sci.* **99**: 6806–6811.
- Nagy, A., Rossant, J., Nagy, R., Abramow-Newerly, W., and Roder, J. 1993. Derivation of completely culture-derived mice from early-passage embryonic stem cells. *Proc. Natl. Acad. Sci.* **90**: 8424–8428.
- Ohlsson, R., Renkawitz, R., and Lobanenko, V. 2001. CTCF is a uniquely versatile transcription regulator linked to epigenetics and disease. *Trends Genet.* **17**: 520–527.
- Olek, A. and Walter, J. 1997. The pre-implantation ontogeny of the *H19* methylation imprint. *Nat. Genet.* **17**: 275–276.
- Olek, A., Oswald, J., and Walter, J. 1996. A modified and improved method for bisulphite based cytosine methylation analysis. *Nucleic Acids Res.* **24**: 5064–5066.
- Sasaki, H., Ferguson-Smith, A., Shum, A., Barton, S., and Surani, M. 1995. Temporal and spatial regulation of *H19* imprinting in normal and uniparental mouse embryos. *Development* **121**: 4195–4202.
- Shemer, R., Hershko, A., Perk, J., Mostoslavsky, R., Tsubert, B.-Z., Cedar, H., Buiting, K. and Razin, A. 2000. The imprinting box of the Prader-Willi/Angelman syndrome domain. *Nat. Genet.* **26**: 440–443.
- Szabó, P., Tang, S.-H., Rentsendorj, A., Pfeifer, G., and Mann, J. 2000. Maternal-specific footprints at putative CTCF sites in the *H19* imprinting control region give evidence for insulator function. *Curr. Biol.* **10**: 607–610.
- Tremblay, K., Duran, K., and Bartolomei, M.S. 1997. A 5' 2-kilobase-pair region of the imprinted mouse *H19* gene exhibits exclusive paternal methylation throughout development. *Mol. Cell. Biol.* **17**: 4322–4329.

The nucleotides responsible for the direct physical contact between the chromatin insulator protein CTCF and the *H19* imprinting control region manifest parent of origin-specific long-distance insulation and methylation-free domains

Vinod Pant, Piero Mariano, Chandrasekhar Kanduri, et al.

Genes Dev. 2003, **17**:

Access the most recent version at doi:[10.1101/gad.254903](https://doi.org/10.1101/gad.254903)

References

This article cites 25 articles, 11 of which can be accessed free at:
<http://genesdev.cshlp.org/content/17/5/586.full.html#ref-list-1>

License

Email Alerting Service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).