RESEARCH COMMUNICATION

A novel cell-to-cell trafficking assay indicates that the KNOX homeodomain is necessary and sufficient for intercellular protein and mRNA trafficking

Jae-Yean Kim,^{1,2,3} Yeonggil Rim,¹ Jing Wang,² and David Jackson^{2,4}

¹Division of Applied Life Science (BK21 program), Environmental Biotechnology National Core Research Center, PMBBRC, Gyeongsang National University, Jinju 660-701, Korea; ²Cold Spring Harbor Laboratory, Cold Spring Harbor, New York 11724, USA

Cell-to-cell trafficking of regulatory proteins is a novel mechanism for communication during cell fate specification in plants. Although several developmental proteins traffic cell-to-cell, no signals that are both necessary and sufficient for this function in developmental proteins have been described. We developed a novel trafficking assay using trichome rescue in *Arabidopsis*. Fusion to KNOTTED1 (KN1) conferred gain-of-trafficking function to the cell-autonomous GLABROUS1 (GL1) protein. We show that the KNOX homeodomain (HD) is necessary and sufficient for intercellular trafficking, identifying a novel function for the HD as the minimal sequence required for trafficking of KN1 and its associated mRNA.

Received December 7, 2004; revised version accepted February 18, 2005.

Intercellular trafficking of regulatory proteins and mRNAs through plasmodesmal (PD) channels allows for the non-cell-autonomous regulation of plant development (Lucas et al. 1995; Perbal et al. 1996; Kim et al. 2001, 2003; Nakajima et al. 2001; Wada et al. 2002; Lee et al. 2003; Wu et al. 2003). However, the mechanisms by which macromolecules traffic cell-to-cell are largely unknown. In some cases transport may be nonselective (Wu et al. 2003), whereas in others the developmental regulation of protein or mRNA movement suggests a selective and regulated process (Kim et al. 2001, 2003; Nakajima et al. 2001; Lee et al. 2003; Gallagher et al. 2004; Sena et al. 2004).

If PD trafficking is regulated, one would expect to find specific signals that are both necessary and sufficient for movement in non-cell-autonomous proteins. Such signal sequence(s) should promote trafficking of a cell-autono-

mous reporter. However no such sequences have been described.

Studies of macromolecular trafficking were pioneered using viral movement proteins (MPs). These proteins traffic through plasmodesmata, increase the size exclusion limit (SEL), and facilitate spread of the viral RNA genome (Wolf et al. 1989; Citovsky et al. 1990; Waigmann et al. 1994). The discovery of a functionally related plant protein, CmPP16, suggests that MPs use an endogenous trafficking mechanism (Xoconostle-Cazares et al. 1999). Microinjection experiments suggested that most PD have an SEL of 1 kDa to a few kilodaltons (Wolf et al. 1989; Waigmann and Zambryski 1995). However, the intercellular trafficking of green fluorescent protein (GFP) suggests that in some cases the PD SEL is larger and is dynamically regulated according to the species, tissue, developmental stage, and environmental conditions (Oparka et al. 1999; Crawford and Zambryski 2000). GFP trafficking is thought to occur by a passive or nonselective diffusion-based mechanism. The developmental transcription factor protein LEAFY may also traffic by a similar nonselective mechanism, raising the question of whether targeting sequences are required for PD transport (Wu et al. 2003).

The maize KNOTTED1 (KN1) homeodomain protein was the first plant protein found to traffic cell-to-cell (Lucas et al. 1995; Kim et al. 2002). This property explains the noncell autonomy of dominant *KN1* alleles in maize leaf development (Hake and Freeling 1986), and trafficking may contribute to the normal function of *KN1* in stem cell maintenance (Kim et al. 2003).

To ask whether KN1 uses a selective trafficking pathway, a novel gain-of-function trafficking assay in *Arabidopsis* was developed, and used to identify a cis-signal for intercellular KN1 trafficking in vivo. Here we show that the homeodomain (HD) regions of KNOTTED1 and related KNOX proteins fit the definition of a trafficking signal domain, and can also promote trafficking of the *KN1* mRNA.

Results and Discussion

To determine whether KN1 contains specific trafficking signal domains, we designed a novel gain-of-function trafficking assay. Since KN1 and its *Arabidopsis* homologs are biologically active in the leaf when overexpressed, we developed the assay in the leaf. We also chose to use *Arabidopsis*, which is more amenable than maize to this kind of analysis. Our assay involves functional complementation of trichome development in *glabrous1* (*gl1*) mutants, mediated by the trafficking of GL1 fusion proteins. GL1, a member of the MYB transcription factor family, is expressed in the epidermis and is required for leaf trichome initiation (Oppenheimer et al. 1991). GL1 probably functions cell autonomously, because sectors lacking GL1 also lack trichomes (M. Hulskamp, pers. comm.). To test the hypothesis that GL1 is cell-autonomous, we expressed it using the *Arabidopsis* rubisco small subunit 2b (*RbcS2b*) promoter in a *gl1* mutant background. This promoter drives expression specifically in green cells (Kim et al. 2003), including mesophyll cells, which normally do not make trichomes, and epidermal guard cells, which are not relevant here as

[**Keywords:** KNOX homeodomain; trichome; trafficking signal; *Glabrous 1*; *knotted1*]

Corresponding authors.

³E-MAIL kimjy@nongae.gsnu.ac.kr; FAX 82-55-759-9363.

⁴E-MAIL jacksond@cshl.edu; FAX (516) 367-8369.

Article and publication are at <http://www.genesdev.org/cgi/doi/10.1101/gad.332805>.

Figure 1. Functional protein trafficking assay using trichome rescue reveals that KN1 fusion to GL1 provides a gain-of-trafficking function. (A) GL1 fusion constructs used for trichome rescue assay. Constructs include the mesophyll-specific *RbcS* promoter (*pRbcS*), GL1 or GL1-KN1 fusions, an internal ribosome entry sequence (IRES) upstream of the GUS coding sequence, and the nopaline synthase (*nos*) 3' terminator. (B) Leaf cross-section viewed in bright-field, with epidermal cells labeled "e". (C) GUS activity was imaged as green fluorescence in mesophyll cells (Immagen Green) in *gl1*/pRbcS::GL1-IRES-GUS plants. The *gl1* background control (D) shows no staining. (E–H) SEM images of wild-type seedling (E), *gl1* mutant seedling (F), *gl1* seedling carrying the control pRbcS::GL1-IRES-GUS transgene (no trichome rescue) (G), and a *gl1* seedling expressing GL1-KN1 showing trichome rescue (H). (I) Schematic of the functional trafficking assay; in the wild type, GL1 functions in an epidermal precursor cell to initiate trichome formation. GL1 expressed in mesophyll cells is cell-autonomous and cannot rescue trichomes in the *gl1* mutant. The KN1 fusion to GL1 can traffic into epidermal cells and rescue trichome formation in the *gl1* mutant. (Ep) Epidermal cells; (Me) mesophyll cells. Bars: (B–D), 50 μ m; (E–H) 100 μ m.

they are symplasmically isolated (Ding et al. 1997). To confirm that the transgene was expressed tissue-specifically, we included an internal ribosome entry sequence (IRES) followed by the β -glucuronidase (GUS) reporter gene (Fig. 1A). Expression of the cell-autonomous GUS reporter indicated that the bi-cistronic mRNA was expressed as expected in mesophyll cells and not in epidermal pavement cells (Fig. 1C).

As predicted, the mesophyll-specific expression of GL1 in *gl1* plants did not rescue trichome formation, indicating that GL1 is cell-autonomous (data from 329 independent transformed plants) (Fig. 1G). We next asked whether fusion of KN1 to GL1 could promote trafficking of the fusion protein. We fused GL1 at the N terminus of KN1, as previous studies showed that this arrangement preserved the biological function of KN1 (Kim et al. 2002). The GL1-KN1 fusion rescued trichome formation in 27% (84/316) of independent transgenic plants (Fig. 1H), suggesting that it could traffic from mesophyll to epidermal cells. The fact that rescue was not

observed in all lines may be due to variability of transgene expression (Peach and Velten 1991). The rescued plants had a mean of 10.2 trichomes per leaf pair, compared with ~60 per leaf pair for wild-type siblings. Trichome rescue was stably inherited into the T2 generation (data not shown).

To investigate whether KN1 contained specific sequences that conferred this cell-to-cell trafficking function, we made a series of deletions. GL1 fusions to the N-terminal region (KN1_{1–135}) or N-terminal/central region (KN1_{1–263}) gave no or negligible trichome rescue in >400 independent transgenic plants (two plants out of 159 transformed with the KN1_{1–263} construct made one trichome per leaf pair) (Fig. 2C). In contrast, a GL1 fusion to the C-terminal region (KN1_{256–359}) permitted trichome rescue in 48% of transgenic lines, with a mean of eight trichomes per leaf pair. These data indicate that the C-terminal region contains a sequence that was both necessary and sufficient for rescue. To confirm that the failure to rescue by the KN1 N-terminal/central region fusions was due to an inability to traffic, and not because these regions caused instability or mislocalization, we expressed them ubiquitously using the 35S promoter. These constructs were able to rescue; for example, 68% of the 35S::GL1-KN1_{1–263} transgenic seedlings showed trichome rescue ($n = 108$) (Fig. 2C), indicating that this fusion protein is biologically active when expressed in epidermal cells.

Figure 2. The KN1 homeodomain functions as a trafficking signal domain. (A) Schematic of domains in KN1: peptide antagonist (PA) (Kragler et al. 2000), MEINOX (Bellaoui et al. 2001), ELK (Kerstetter et al. 1994), homeodomain, and C-terminal region (C-pep). Red box indicates nuclear localization signal. (B) The predicted structure of the homeodomain region of KN1. Numbers in A and B indicate amino acids from the ATG. (C) Trichome rescue by different fusions to GL1. Trichome rescue rate (percent of T1 plants showing trichome rescue) is on the Y-axis. The number next to each bar is the average number of trichomes per leaf pair. Note that all constructs are fusions to GL1 driven by the *pRbcS* promoter, unless labeled "35S", in which case it is driven by the 35S promoter. (D,E) GFP fluorescence (D) and GUS activity (E) in rescued trichomes of *gl1*; pGL2::GUS plants containing the pRbcS::GFP-GL1-KN1_{256–359} transgene. Red arrow indicates GFP fluorescence in trichome nucleus. Bars: D,E, 150 μ m.

The C-terminal region of KN1 (KN1_{256–359}) that was necessary and sufficient for trafficking included the HD, and we attempted to further delimit trafficking activity. Deletions at the N terminus that removed any part of the HD, for example removal of the NLS region or of a part of the first helix, abolished trafficking activity, as in KN1_{273–359} or KN1_{295–359} (Fig. 2C). We also made deletions from the C terminus. We found that removal of the residues C-terminal to the HD (KN1_{327–359}) (C-pep in Fig. 2A) stimulated by ~1.6-fold the percentage of transgenic plants that showed trichome rescue (construct KN1_{256–326}) (Fig. 2C, cf. KN1_{256–359}). However any further deletion that encroached on the HD, for example removal of part of the third helix in KN1_{256–311}, led to a complete loss of trichome rescue. In summary, sequences within the HD at both the N and C termini were necessary for trichome rescue, and presumably therefore for trafficking. We made a minimal construct that contained only the HD and three amino acids at its N terminus (KN1_{261–326}), and this also gave trichome rescue (Fig. 2C). KN1_{273–326}, composed only of the three predicted HD helices, gave trichome rescue in only 6% of transgenic plants, and the degree of rescue was very low, with an average of only 2.2 trichomes per leaf pair. Consistent with the finding that both N- and C-terminal regions of the HD were necessary for trichome rescue, other constructs that expressed smaller regions of the HD were nonfunctional (e.g., KN1_{261–302}, KN1_{277–311}) (Fig. 2C).

Since the HD also contains the DNA binding activity, we asked whether trichome rescue was correlated with cell-to-cell trafficking of the KN1 fusion, rather than nonautonomous action of a downstream factor that might be expressed in response to DNA binding by the GL1–KN1 fusion. We used *GLABRA2* (*GL2*) as a reporter, since its expression in the trichome is dependent on the presence of *GL1* (Szymanski et al. 1998). *gl1* plants carrying a *pRbcS::GFP–GL1–KN1*_{256–359} construct, which encodes a fusion of GFP, GL1, and the C-terminal domain of KN1. As expected, expression of this construct allowed trichome rescue, and we also detected GFP fluorescence in nuclei of rescued trichomes, indicating that the GFP–GL1–KN1_{256–359} protein trafficked into the epidermal cells (Fig. 2D). Furthermore, we could detect activity of the *GL2* promoter as monitored by GUS activity in the rescued trichomes (Fig. 2E). These data strongly suggest that trichome rescue was due to trafficking of the GL1–KN1 fusion protein into the epidermis, rather than by movement of a nonautonomous downstream factor.

The HD is the defining feature of KNOX family members (Fig. 3A), so we investigated whether the trafficking activity was conserved in other plant HD proteins. The KNOX proteins can be divided into two classes based on sequence conservation and expression patterns (Kerstetter et al. 1994). We found that only the C-terminal regions of closely related class I KNOX proteins (*Arabidopsis* STM, KNAT1, or tomato LeT6) were functional in trichome rescue (Fig. 3B). The less closely related class I KNOX C-terminal regions from *Arabidopsis* proteins KNAT2 or KNAT6, or a class II HD (KNAT3) or a BEL-like protein (BELLRINGER) (Byrne et al. 2003) did not show any trichome rescue. Therefore trafficking appeared to be a property only of specific class I HDs closely related to KN1. The trichome rescue rate using

Figure 3. The HD trafficking signal is conserved in closely related class I KNOX homeodomain proteins, and promotes KN1 mRNA movement. (A) Phylogenetic tree of homeodomain sequences: LeT6 (accession no. T04317), BLR (AAP93641), BEL1 (AAM62510), STM (Q38874), KNAT1 (AAM03027), KNAT2 (P46640), KNAT3 (P48000), KNAT4 (P48001), KNAT5 (P48002), KNAT6 (NP_850951), KNAT7 (NP_564805). (B) Trichome rescue assays were performed using HD regions corresponding to KN1_{256–359}. Descriptions of the chart are as in Figure 2C. (C) RNA trafficking assay. Stop codons are shown as a red asterisk after the GL1 coding region. The shorter KN1 regions shown in constructs in lines 2, 4, and 7 correspond to KN1_{256–359}. The number of T1 plants showing trichome rescue/total number of T1 plants observed (rescue rate, percent) is shown on the right side of each construct. (3') 3' untranslated region (UTR) of KN1 or of nopaline synthase (see Materials and Methods).

these proteins was lower than with KN1, and we cannot at present explain these differences. However the proteins differ in sequences C-terminal to the homeodomain, and this region of KN1 had a negative effect on trafficking. They also differ in the HD itself, and future domain swap experiments should allow the effects of these regions to be distinguished. Our demonstration of the HD region as a minimal trafficking domain that is both necessary and sufficient agrees with earlier mutagenesis studies that found a mutation in the HD that blocked trafficking of microinjected or in vivo expressed proteins (Lucas et al. 1995; Kim et al. 2002). Our results conflict with studies that identified a motif in the N-terminal region of KN1 as a peptide antagonist of trafficking (Kragler et al. 2000), as that region appears not to be sufficient for trafficking in our assay.

Nonautonomy of KNOX mRNAs has also been reported (Lucas et al. 1995; Kim et al. 2001). To investigate whether trichome rescue was due to trafficking of the KN1 fusion proteins, or of their mRNAs, or both, we introduced a stop codon after the GL1 coding region in the pRbcS::GL1–KN1 or pRbcS::GL1–KN1_{256–359} constructs (Fig. 3C), so that they expressed only the GL1 protein from a GL1_{STOP}–KN1 fusion mRNA. Plants transformed with these constructs did not show trichome rescue (Fig. 3C), suggesting that the GL1–KN1 fusion protein was required and that the fusion mRNA itself was not sufficient. We also tested whether KN1 could facilitate the trafficking of its own mRNA, as suggested by microinjection studies (Lucas et al. 1995). As class I KNOX genes are normally not expressed

in the *Arabidopsis* leaf (Lincoln et al. 1994), we coexpressed the pRbcS::GL1_{STOP}-KN1 with GFP-KN1 or GFP-KN1₂₅₆₋₃₂₆ expressed from the 35S promoter. In these plants we observed trichome rescue, suggesting that GFP-KN1 or GFP-KN1₂₅₆₋₃₂₆ could facilitate trafficking of the GL1_{STOP}-KN1 mRNA (Fig. 3C). KN1 mRNA sequences were required for this rescue, because a control construct, pRbcS::GL1_{STOP}-GUS, did not rescue, even in the presence of GFP-KN1 or GFP-KN1₂₅₆₋₃₂₆ (Fig. 3C). This eliminates the possibility that rescue was due to GL1 protein movement by a nonspecific PD gating effect of GFP-KN1 overexpression, and suggests that the GL1_{STOP}-KN1 mRNA was specifically transported to the epidermis by GFP-KN1 or GFP-KN1₂₅₆₋₃₂₆.

Our results raise the question of whether trafficking and trichome rescue were due to a signal in the KN1 HD alone, or in the HD in combination with its mRNA. We showed here that KN1 mRNA trafficking is dependent on the presence of the KN1 protein or of the HD. Therefore the trafficking signal is certainly not solely contained within the KN1 mRNA. It remains to be seen whether the KN1 mRNA, in combination with the KN1 HD, is a necessary component of the signal for intercellular trafficking. However, our prior microinjection studies strongly suggest that at least the full-length KN1 protein can traffic without its mRNA (Lucas et al. 1995). Our results also imply a specific interaction between KN1 and its mRNA, because the control GL1-GUS fusion mRNA expressed together with KN1 did not rescue trichomes, and presumably did not traffic. The proposed interaction between KN1 and its mRNA could be mediated at the sequence level through binding of the HD to specific sequences in the KN1 mRNA, similar to the binding of CAUDAL mRNA by the BICOID HD protein (Dubnau and Struhl 1996).

KN1 and other class I KNOX proteins normally function in the shoot apical meristem (Reiser et al. 2000). We therefore investigated whether the trafficking signal region of KN1 was also necessary and sufficient in the shoot apical meristem (SAM). We used *WUSCHEL* (*WUS*) regulatory sequences to drive layer-specific expression. *WUS* is expressed in the L3 layer of the SAM (Mayer et al. 1998), and we found that 1.7 kb of the *WUS* 5' region (pWUS) drove expression of a cell-autonomous mGFP5-ER reporter in the SAM, specifically in the L3 layer (Fig. 4E,F). GFP-KN1 expression using the same promoter showed green fluorescence in all layers, indicating that GFP-KN1 could traffic from the L3 to the L2/L1 (Fig. 4G,H). This construct was also able to rescue the *shootmeristemless* mutation (Fig. 4B-D), indicating biological activity of the GFP-KN1 fusion protein in cells into which it was trafficking. The rescued plants were not completely normal, and this may be because pWUS also drove a low level of expression in developing leaf primordia (data not shown).

Because the SEL in the meristem is larger than in the leaf (Wu et al. 2003), we next used a GFP-YFP fusion as a reporter. Expression of GFP-YFP-KN1₁₋₂₆₃ using pWUS resulted in fluorescence restricted to the L3 layer, suggesting that, as in the leaf, the N-terminal/middle regions of KN1 were not sufficient for trafficking in the SAM (Fig. 4I,J). In contrast, the GFP-YFP-KN1₂₅₆₋₃₅₉ fusion could traffic from the L3 to the L2/L1 layers (Fig. 4K,L). Each pattern was confirmed in a minimum of four independent transgenic plants. Therefore the KN1 C-ter-

Figure 4. The KN1 trafficking signal domain functions in the meristem, and L3-specific expression of KN1 is sufficient to complement *shootmeristemless-11* (*stm-11*) mutants. (A) GFP fusion constructs used for protein trafficking assays in the SAM. Constructs include the L3-specific *WUS* promoter (pWUS), mGFP5-ER, GFP, or GFP-YFP fusions to KN1 or to KN1 deletions constructs, and the nopaline synthase terminator (nos 3') or the KN1 3'UTR (3'). The third construct contains KN1₁₋₂₆₃ and the fourth contains KN1₂₅₆₋₃₅₉. (B) *stm-11* mutants show fused cotyledons and no true leaves. pWUS::KN1 expression (C) rescued 14 plants out of 46 *stm-11* T1 homozygotes, and pWUS::GFP-KN1 expression (D) rescued four plants out of 10 *stm-11* T1 homozygotes. Thus pWUS-driven KN1 or GFP-KN1 expression can rescue the function of *STM*, which is normally expressed in all three layers of the SAM (Long et al. 1996). (E-L) Confocal microscope images of meristems expressing these GFP constructs; mGFP5-ER is restricted to the L3 (E,F). GFP-KN1₁₋₃₅₉ shows movement into L2 and L1 cells, seen as green nuclear fluorescence (G) or in grayscale with labeled nuclei indicated by arrows (H). GFP-YFP-KN1₁₋₂₆₃ is restricted to the L3 (I,J), and GFP-YFP-KN1₂₅₆₋₃₅₉ shows movement into L2 and L1 cells (K,L), indicating that the trafficking signal region identified in KN1₂₅₆₋₃₅₉ also functions in the meristem. Note that in the inverted grayscale images, the fluorescence of nuclei in the L1 and L2 layers is more pronounced. Labeled nuclei in the L1 and L2 layers are indicated by arrows in H,L. The outer surface of the L1 is marked by a blue line in E,I,K and by a black line in F,J,L. Bars: E-L, 50 μ m.

минаl trafficking signal identified using trichome rescue was also necessary and sufficient for trafficking in the SAM.

The trafficking signal that we report here is the first report of one that is both necessary and sufficient for selective intercellular trafficking in the context of a heterologous cell-autonomous reporter. During plant evolution, HD proteins may have gained a selective trafficking function to regulate cell fate through non-cell-autonomous signaling. The differential regulation of KNOX protein trafficking may provide a mechanism to differentiate homologous, potentially redundant KNOX gene functions when they are expressed in the same tissues. Indeed, closely related KNOX proteins have some degree of redundant function; for example, KNAT1 expression can partially complement the *stm* phenotype (Byrne et al. 2002), and KNAT1 may also regulate epidermal cell fate nonautonomously (Venglat et al. 2002).

Our functional trafficking assay could be used to determine whether diverse proteins use a selective trafficking pathway in vivo. Owing to the requirement for gain-

of-trafficking function, only proteins carrying a positive trafficking signal will promote trichome rescue. The LEAFY transcription factor was recently suggested to move by a nontargeted mechanism (Wu et al. 2003); however, it is possible that it contains redundant selective signals, and it will be interesting to test the ability of LEAFY and other plant proteins to promote trafficking in our system. Trafficking signals identified in these studies may be useful tools to selectively control the misexpression of other cell-autonomous proteins. However, our results do not imply that the KN1 HD can confer protein trafficking in any context, because its own trafficking is developmentally regulated (Kim et al. 2003). In conclusion, using a functional trafficking assay we found a novel role for the KNOX HD. In addition to its function as a DNA-binding domain, this domain also acts as an intercellular protein trafficking signal, and as a mediator for selective cell-to-cell mRNA trafficking.

Materials and methods

Trichome rescue assay

The *gll-1* mutant line originally isolated in Landsberg *erecta* (Ler) and introgressed to Wassilewskija (Ws) was transformed using *Agrobacteria*-containing constructs to express GL1 fusion proteins, according to Kim et al. (2003). Transgenic seedlings were screened on MS medium containing 50 µg/mL hygromycin. We scored trichome rescue from the first four leaves of 2-wk-old seedlings using a stereo microscope.

GUS staining, confocal and scanning electron microscopy

GUS staining using X-glucuronide was performed as described previously (Kim et al. 2003). The stained tissues were cleared in 70% ethanol for 3 h at 37°C, and photographed. Staining using the ImmagenGreen kit was performed according to the manufacturer (Molecular Probes). In brief, leaf cross-sections were incubated in 100 mM phosphate buffer (pH 7.5) containing 50 µM C₁₂FDGluU at 37°C for 2 h. Confocal and scanning electron microscopy were performed as described (Taguchi-Shiobara et al. 2001; Kim et al. 2003).

DNA constructs

The cloning of GFP-KN1 and *pRbcS* was described in Kim et al. (2002, 2003); the *pRbcS* was upstream of GFP-KN1 in pCambia1300. *pRbcS::GL1-KN1* was made by replacing GFP with GL1. The IRES sequence (Kim et al. 2002) was inserted in pCambia2301 to make the IRES-GUS construct; *pRbcS::GL1-KN1* was then inserted upstream to make *pRbcS::GL1-KN1-IRES-GUS*, and *pRbcS::GL1-IRES-GUS* was produced by deleting KN1 from this clone. To make deletion constructs of KN1 or KNOX proteins, specific regions were amplified by PCR and inserted to replace KN1 in *pRbcS::GL1-KN1*. For constructs used in the RNA trafficking assay, an in-frame stop codon was inserted after the GL1 coding region to produce *pRbcS::GL1_{STOP}-KN1* or *pRbcS::GL1_{STOP}-GUS*. The *pRbcS::GL1_{STOP}-KN1* construct carries the 3' untranslated region (UTR) of KN1, and *pRbcS::GL1_{STOP}-GUS* carries the 3'UTR of nopaline synthase. To engineer RNA trafficking constructs to overexpress KN1 or KN1₂₆₆₋₃₂₆, a HindIII fragment of 35S::GFP-KN1 (Kim et al. 2002) or of 35S::GFP-KN1₂₅₆₋₃₂₆ was end-filled by Klenow treatment and ligated into the PmlI site of *pRbcS::GL1_{STOP}-KN1* or the SmaI site of *pRbcS::GL1_{STOP}-GUS*. *pWUS* was produced by PCR using primers GGGGAATTCTGGATGGAATCCAAAAGTTTAT and GGGGGTACCCCGGTGTGTTTGATTCGACTTTTGT and subcloned upstream of "double GFP" reporters. The *pWUS* constructs carrying KN1 or its fragments (KN1₁₋₂₆₃, KN1₂₅₆₋₃₅₉) have the 3'UTR of KN1. All PCRs were carried out using high-fidelity DNA polymerase (Pfu, Stratagene) and were confirmed by sequencing.

In the primers listed below, f indicates forward, r denotes reverse, and numbers indicate locations of first or last amino acid of each amplified fragment. Lowercase letters include restriction enzyme sites added for cloning.

The following primers were used: KN1-1f, ggggagatctTATGGAGGAGATACCCAACA; KN1-135r, ggggggtaccTTACGCCGACACCTCCG

GTG; KN1-263r, ggggggtaccTTATGACAGTTCTTGCTTGAGCG; KN1-261f, ggggagatctGTCAAAGAAGAAGAAAGG; KN1-302r, ggggggtaccTTAAGCCAGTGCCACCTTCTG; KN1-277f, ggggagatctCCTTAGCTGGTGGGATCAG; KN1-311r, ggggggtaccTTAGAACCAGTTGTTGATCTGCT; KN1-295f, ggggagatctGAAGGTGGCACTGGCTGA; KN1-360r, ggggggtaccCATCCATCATCAGGTGGT; KNAT1-f, ggggagatctAACGATTTGAAGCAAGAAC; KNAT1-r, ggggggtaccTTATGGACCGAGACGATAAG; KNAT1-f, ggggagatctCAGTTCATTGAACTCGAGTT; KNAT1-r, ggggggtaccTTACTCGGTAAAGAATGTTTCA; KNAT1-f, ggagagatctAGTAGACATAAGAGAGGAGATATTA; KNAT1-r, ggggggtaccCTACGCCAACCCTCTCT; KNAT1-f, ggggagatctTAGTACTTTAAAGCTTGAGTTCTC; KNAT1-r, ggggggtaccTCATTCCTCGGTAAAGAAATG; STM-f, ggggagatctAGGGAGCCTCAAGCAAGA; STM-r, ggggggtaccTCAAAGCATGGTGGAGGAG; BLR-f, ggggagatctATTCTCTGATCATCTGCTCCT; BLR-r, ggggggtaccTCGAGTCTCCAGCATGTGAA; LeT6-f, ggggagatctGGGAAGCCTTAAGCAGGAGT; LeT6-r, gggggtaccTAGCAAGTACAACTATTTAATTCATAATTA.

Acknowledgments

We thank Myeong-Min Lee (Yonsei University) for *gll* and *gll* (GL2::GUS) mutant seeds, and M. David Marks (University of Minnesota) for GL1 cDNA. D.J. was supported by the National Science Foundation Integrative Plant Biology Grant 9727959 and Cold Spring Harbor Laboratory institutional funds. J.Y.K. was funded by a grant from KOSEF/MOST to the Environmental Biotechnology National Core Research Center (R15-2003-012-01003-0) and a grant from the BioGreen 21 Program, Rural Development Administration, Republic of Korea. This work was supported by Korea Research Foundation Grant KRF-2003-042-C00124.

References

- Bellaoui, M., Pidkowich, M.S., Samach, A., Kushalappa, K., Kohalmi, S.E., Modrusan, Z., Crosby, W.L., and Haughn, G.W. 2001. The *Arabidopsis* BELL1 and KNOX TALE homeodomain proteins interact through a domain conserved between plants and animals. *Plant Cell* **13**: 2455–2470.
- Byrne, M.E., Simorowski, J., and Martienssen, R.A. 2002. ASYMMETRIC LEAVES1 reveals knox gene redundancy in *Arabidopsis*. *Development* **129**: 1957–1965.
- Byrne, M.E., Groover, A.T., Fontana, J.R., and Martienssen, R.A. 2003. Phyllotactic pattern and stem cell fate are determined by the *Arabidopsis* homeobox gene BELLRINGER. *Development* **130**: 3941–3950.
- Citovsky, V., Knorr, D., Schuster, G., and Zambryski, P. 1990. The P30 movement protein of tobacco mosaic virus is a single-strand nucleic acid binding protein. *Cell* **60**: 637–647.
- Crawford, K.M. and Zambryski, P.C. 2000. Subcellular localization determines the availability of non-targeted proteins to plasmodesmatal transport. *Curr. Biol.* **10**: 1032–1040.
- Ding, B., Kwon, M.O., Hammond, R., and Owens, R. 1997. Cell-to-cell movement of potato spindle tuber viroid. *Plant J.* **12**: 931–936.
- Dubnau, J. and Struhl, G. 1996. RNA recognition and translational regulation by a homeodomain protein. *Nature* **379**: 694–699.
- Gallagher, K.L., Paquette, A.J., Nakajima, K., and Benfey, P.N. 2004. Mechanisms regulating SHORT-ROOT intercellular movement. *Curr. Biol.* **14**: 1847–1851.
- Hake, S. and Freeling, M. 1986. Analysis of genetic mosaics shows that the extraepidermal cell divisions in Knotted1 mutant maize plants are induced by adjacent mesophyll cells. *Nature* **320**: 621–623.
- Kerstetter, R., Vollbrecht, E., Lowe, B., Veit, B., Yamaguchi, J., and Hake, S. 1994. Sequence analysis and expression patterns divide the maize knotted1-like homeobox genes into two classes. *Plant Cell* **6**: 1877–1887.
- Kim, M., Canio, W., Kessler, S., and Sinha, N. 2001. Developmental changes due to long-distance movement of a homeobox fusion transcript in tomato. *Science* **293**: 287–289.
- Kim, J.Y., Yuan, Z., Cilia, M., Khalfan-Jagani, Z., and Jackson, D. 2002. Intercellular trafficking of a KNOTTED1 green fluorescent protein fusion in the leaf and shoot meristem of *Arabidopsis*. *Proc. Natl. Acad. Sci.* **99**: 4103–4108.
- Kim, J.Y., Yuan, Z., and Jackson, D. 2003. Developmental regulation and significance of KNOX protein trafficking in *Arabidopsis*. *Develop-*

- ment **130**: 4351–4362.
- Kragler, F., Monzer, J., Xoconostle-Cazares, B., and Lucas, W.J. 2000. Peptide antagonists of the plasmodesmal macromolecular trafficking pathway. *EMBO J.* **19**: 2856–2868.
- Lee, J.Y., Yoo, B.C., Rojas, M.R., Gomez-Ospina, N., Staehelin, L.A., and Lucas, W.J. 2003. Selective trafficking of non-cell-autonomous proteins mediated by NtNCAPP1. *Science* **299**: 392–396.
- Lincoln, C., Long, J., Yamaguchi, J., Serikawa, K., and Hake, S. 1994. A knotted1-like homeobox gene in *Arabidopsis* is expressed in the vegetative meristem and dramatically alters leaf morphology when overexpressed in transgenic plants. *Plant Cell* **6**: 1859–1876.
- Long, J.A., Moan, E.I., Medford, J.I., and Barton, M.K. 1996. A member of the KNOTTED class of homeodomain proteins encoded by the STM gene of *Arabidopsis*. *Nature* **379**: 66–69.
- Lucas, W.J., Bouche-Pillon, S., Jackson, D.P., Nguyen, L., Baker, L., Ding, B., and Hake, S. 1995. Selective trafficking of KNOTTED1 homeodomain protein and its mRNA through plasmodesmata. *Science* **270**: 1980–1983.
- Mayer, K.F., Schoof, H., Haecker, A., Lenhard, M., Jurgens, G., and Laux, T. 1998. Role of WUSCHEL in regulating stem cell fate in the *Arabidopsis* shoot meristem. *Cell* **95**: 805–815.
- Nakajima, K., Sena, G., Nawy, T., and Benfey, P.N. 2001. Intercellular movement of the putative transcription factor SHR in root patterning. *Nature* **413**: 307–311.
- Oparka, K.J., Roberts, A.G., Boevink, P., Santa Cruz, S., Roberts, I., Pradel, K.S., Imlau, A., Kotlizky, G., Sauer, N., and Epel, B. 1999. Simple, but not branched, plasmodesmata allow the nonspecific trafficking of proteins in developing tobacco leaves. *Cell* **97**: 743–754.
- Oppenheimer, D.G., Herman, P.L., Sivakumaran, S., Esch, J., and Marks, M.D. 1991. A myb gene required for leaf trichome differentiation in *Arabidopsis* is expressed in stipules. *Cell* **67**: 483–493.
- Peach, C. and Velten, J. 1991. Transgene expression variability (position effect) of CAT and GUS reporter genes driven by linked divergent T-DNA promoters. *Plant Mol. Biol.* **17**: 49–60.
- Perbal, M.C., Haughn, G., Saedler, H., and Schwarz-Sommer, Z. 1996. Non-cell-autonomous function of the Antirrhinum floral homeotic proteins DEFICIENS and GLOBOSA is exerted by their polar cell-to-cell trafficking. *Development* **122**: 3433–3441.
- Reiser, L., Sanchez-Baracaldo, P., and Hake, S. 2000. Knots in the family tree: Evolutionary relationships and functions of knox homeobox genes. *Plant Mol. Biol.* **42**: 151–166.
- Sena, G., Jung, J.W., and Benfey, P.N. 2004. A broad competence to respond to SHORT ROOT revealed by tissue-specific ectopic expression. *Development* **131**: 2817–2826.
- Szymanski, D.B., Jilk, R.A., Pollock, S.M., and Marks, M.D. 1998. Control of GL2 expression in *Arabidopsis* leaves and trichomes. *Development* **125**: 1161–1171.
- Taguchi-Shiobara, F., Yuan, Z., Hake, S., and Jackson, D. 2001. The fasciated ear2 gene encodes a leucine-rich repeat receptor-like protein that regulates shoot meristem proliferation in maize. *Genes & Dev.* **15**: 2755–2766.
- Venglat, S.P., Dumonceaux, T., Rozwadowski, K., Parnell, L., Babic, V., Keller, W., Martienssen, R., Selvaraj, G., and Datla, R. 2002. The homeobox gene BREVIPEDICELLUS is a key regulator of inflorescence architecture in *Arabidopsis*. *Proc. Natl. Acad. Sci.* **99**: 4730–4735.
- Wada, T., Kurata, T., Tominaga, R., Koshino-Kimura, Y., Tachibana, T., Goto, K., Marks, M.D., Shimura, Y., and Okada, K. 2002. Role of a positive regulator of root hair development, CAPRICE, in *Arabidopsis* root epidermal cell differentiation. *Development* **129**: 5409–5419.
- Waigmann, E. and Zambryski, P. 1995. Tobacco mosaic virus movement protein-mediated protein transport between trichome cells. *Plant Cell* **7**: 2069–2079.
- Waigmann, E., Lucas, W.J., Citovsky, V., and Zambryski, P. 1994. Direct functional assay for tobacco mosaic virus cell-to-cell movement protein and identification of a domain involved in increasing plasmodesmal permeability. *Proc. Natl. Acad. Sci.* **91**: 1433–1437.
- Wolf, S., Deom, C.M., and Beachy, R.N. 1989. Movement protein of tobacco mosaic virus modifies plasmodesmal size exclusion limit. *Science* **246**: 377–379.
- Wu, X., Dinnyen, J.D., Crawford, K.M., Rhee, Y., Citovsky, V., Zambryski, P., and Weigel, D. 2003. Modes of intercellular transcription factor movement in the *Arabidopsis* apex. *Development* **130**: 3735–3745.
- Xoconostle-Cazares, B., Xiang, Y., Ruiz-Medrano, R., Wang, H.L., Monzer, J., Yoo, B.C., McFarland, K.C., Franceschi, V.R., and Lucas, W.J. 1999. Plant paralog to viral movement protein that potentiates transport of mRNA into the phloem. *Science* **283**: 94–98.

A novel cell-to-cell trafficking assay indicates that the KNOX homeodomain is necessary and sufficient for intercellular protein and mRNA trafficking

Jae-Yean Kim, Yeonggil Rim, Jing Wang, et al.

Genes Dev. 2005, **19**:

Access the most recent version at doi:[10.1101/gad.332805](https://doi.org/10.1101/gad.332805)

References

This article cites 34 articles, 21 of which can be accessed free at:
<http://genesdev.cshlp.org/content/19/7/788.full.html#ref-list-1>

License

Email Alerting Service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).