RESEARCH COMMUNICATION

MtHAP2-1* is a key transcriptional regulator of symbiotic nodule development regulated by microRNA169 in *Medicago truncatula

Jean-Philippe Combier,¹ Florian Frugier,²
 Françoise de Billy,¹ Adnane Boualem,²
 Fikri El-Yahyaoui,¹ Sandra Moreau,¹
 Tatiana Vernié,¹ Thomas Ott,¹ Pascal Gamas,¹
 Martin Crespi,^{2,4} and Andreas Niebel^{1,3}

¹LIPM (Laboratoire des Interactions Plantes-Microorganismes) INRA-CNRS (Institut de National de la Recherche-Centre national de la recherche scientifique), 31326 Castanet-Tolosan, France; ²ISV (Institut des Sciences du Végétal), CNRS, 91198 Gif sur Yvette, France

In the model legume *Medicago truncatula*, we identified a new transcription factor of the CCAAT-binding family, *MtHAP2-1*, for which RNA interference (RNAi) and in situ hybridization experiments indicate a key role during nodule development, possibly by controlling nodule meristem function. We could also show that *MtHAP2-1* is regulated by microRNA169, whose overexpression leads to the same nodule developmental block as *MtHAP2-1* RNAi constructs. The complementary expression pattern of miR169 and *MtHAP2-1* and the phenotype of miR169-resistant *MtHAP2-1* nodules strongly suggest, in addition, that the miR169-mediated restriction of *MtHAP2-1* expression to the nodule meristematic zone is essential for the differentiation of nodule cells.

Supplemental material is available at <http://www.genesdev.org>.

Received July 24, 2006; revised version accepted September 25, 2006.

The symbiotic association between nitrogen-fixing bacteria, collectively known as rhizobia, and plants belonging to the legume family, results in the formation of specialized root organs called nodules, inside which differentiated rhizobia reduce atmospheric nitrogen to ammonium for the benefit of the plant. In the model legume *Medicago truncatula*, nodules are indeterminate; they originate from root cortical cells that are stimulated by rhizobia to re-enter mitosis, which leads to the formation of a persistent meristem, allowing the subsequent development of the nodule (Geurts et al. 2005; Stacey et al. 2006). To obtain insights into the largely unknown mechanisms regulating root nodule development, transcriptome studies were performed in *M. truncatula* to

search for transcription factors (TFs) implicated in early stages of nodulation. In this way we identified a gene, *MtHAP2-1*, whose expression is strongly up-regulated during nodule development (El Yahyaoui et al. 2004). This gene encodes an HAP2-type TF subunit, a component of the hetero-trimeric CCAAT-box-binding factor complex (CBF/NF-Y/HAP), comprising HAP2 (CBF-B, NF-YA), HAP3 (CBF-A, NF-YB), and HAP5 (CBF-C, NF-YC) subunits and characterized by its ability to bind the CCAAT motif present in many eukaryotic promoters (Maity and de Crombrughe 1998). Interestingly, several genes coding CBF subunits have been shown to play central roles in development. This is the case for *CBF-B* genes in the mouse (Bhattacharya et al. 2003), as well as for *LEAFY COTYLEDON* (*LEC1*) and *LEAFY COTYLEDON1-LIKE* (*L1L*), two *HAP3* genes shown to be central regulators of embryogenesis in *Arabidopsis thaliana* (Lottan et al. 1998; Kwong et al. 2003; Lee et al. 2003). Similarly, *HAP3* genes are involved in chloroplast biogenesis in rice (Miyoshi et al. 2003) and *HAP5* in flowering-time control in *Arabidopsis* (Ben-Naim et al. 2006). In contrast to yeast and animals (where only one or two genes encode each CBF subunit), plants possess significantly more complex gene families. In *A. thaliana* there are 10

Figure 1. *MtHAP2-1* expression is nodule specific and restricted to the meristematic-zone mRNA in situ hybridization of *MtHAP2-1* in nitrogen-fixing nodules. Sections are hybridized with an antisense *MtHAP2-1* probe (*a,b*) or a control sense *MtHAP2-1* probe (*c,d*). *a* and *c* are dark-field microscopy images, to visualize silver grains corresponding to *MtHAP2-1* mRNA localization. Nodule zones are indicated according to Vasse et al. (1990) (I, meristematic zone; II, infection zone; II-III, interzone; III, nitrogen fixation zone). *b* and *d* are color density maps performed using Image-Pro plus and Diatrack softwares. Bar, 100 µm.

[Keywords: Nodulation; microRNA; transcription factor; symbiosis; HAP2; *Medicago*]

Corresponding authors.

³E-MAIL aniebel@toulouse.inra.fr; FAX 33-5-61285061.

⁴E-MAIL martin.crespi@isv.cnrs-gif.fr; FAX 33-1-69823695.

Article is online at <http://www.genesdev.org/cgi/doi/10.1101/gad.402806>.

HAP2, 10 *HAP3*, and nine *HAP5* genes (Gusmaroli et al. 2002). Functionally specialized gene family members controlling specific developmental pathways thus appear to exist in plants, possibly because of their ability to bind to other TFs such as MADS-box TFs (Masiero et al. 2002) or nuclear regulators such as CONSTANS (Ben-Naim et al. 2006).

Results and Discussion

We therefore undertook a detailed characterization of the *MtHAP2-1* gene, which is strongly and specifically activated during symbiotic interactions (Supplementary Fig. 1). In situ hybridization showed that *MtHAP2-1* mRNA is most abundant in cells of the nodule meristematic zone, less abundant in adjacent cells of the nodule infection zone, and not detected in all other nodule tissues (Fig. 1). The symbiotic role of *MtHAP2-1* was then investigated by expressing *MtHAP2-1* RNA interference (RNAi) constructs in transgenic *M. truncatula* roots. This resulted in a reduction of *MtHAP2-1* expression (Fig. 4b, below) sufficient to significantly alter nodule development. Nodulation was delayed and nodule growth arrested at ~8–10 d post-inoculation (Fig. 2a,b). Inside the resulting spherical nodular structures, the absence of the clearly delimited zones found in wild-type nodules (I, meristem zone; II, Rhizobia infection zone; III, nitrogen-fixing zone; Fig. 2e,g) indicated an abnormal nodule developmental process. Growth arrest was often associated with the development of a closed endodermal layer surrounding the nodules (Fig. 2e). In addition, cells of the apical meristematic zone lost their typical polyhedral shape and accumulated numerous enlarged vacuoles (Fig. 2c,d). *MtHAP2-1* RNAi nodules were also unable to fix nitrogen as shown by the reduced development of plants grown in the absence of mineral nitrogen (Fig. 2h). This phenotype was confirmed by electron microscopy observations showing that in the infection zone of mature *MtHAP2-1*-RNAi nodules, bacteria were not released from infection threads (Supplementary Fig. 2). On the few occasions for which release could be observed, the corresponding rhizobia were arrested in their development and never differentiated into nitrogen-fixing type IV bacteroids (Vasse et al. 1990; data not shown). Taken together, these results suggest that *MtHAP2-1* is a novel symbiosis-specific TF playing a key role during nodule development by controlling meristem persistence and probably also symbiotic bacterial release.

MicroRNAs (miRNA) are small 21-nucleotide (nt) RNAs that play important regulatory roles in plants and metazoans by targeting mRNAs either for cleavage or translation inhibition (Jones-Rhoades et al. 2006). miRNAs have generally been shown to target genes controlling development, many of them encoding TFs (Li and Zhang 2005; Jones-Rhoades et al. 2006). Since previous studies had suggested

that HAP2 TFs are potential targets of miR169 in *A. thaliana* (Jones-Rhoades and Bartel 2004), we searched for miR169 complementary sequences in the *MtHAP2-1* gene, and found two such sequences in the 3' untranslated region (UTR) (Fig. 3a). An in silico approach then identified a *M. truncatula* gene susceptible to encode a miR169 precursor, called *MtmiR169-a*, and secondary structure predictions revealed the capacity of *MtmiR169-a* to form a stable stem-loop, as expected for a pre-miRNA (Fig. 3b). To investigate gene functionality we overexpressed the putative *MtmiR169-a* gene in transgenic *M. truncatula* roots under the control of a 35S

Figure 2. Altered nodulation phenotypes of *MtHAP2-1* RNAi and 35S:*MtmiR169-a* roots. (a,b) Nodule development on roots transformed with either an *MtHAP2-1* RNAi construct (a) or a control construct (empty vector) (b). Pictures were taken 46 d post-*Rhizobium* inoculation (d.p.i.). (c,d) Electron microscopy images of the meristematic zone of *MtHAP2-1* RNAi (c) and control (d) nodules. (V) Vacuoles. Cells of the meristematic region of *MtHAP2-1* RNAi nodules already start to lose meristematic characteristics at 10 d.p.i. (c), while they are still fully functional in control nodules at 27 d.p.i. (d). Bar, 5 μ m. (e–g) Light microscopic images of longitudinal sections of 20-d-old nodules, using an overlay of two fluorescence images. Note the absence of normal nodule zonation and the abnormal closed endodermis (E) in *MtHAP2-1* RNAi (e) and *MtmiR169-a*-overexpressing (f) nodules, compared with control (empty vector) (g) nodules, indicating nodule developmental arrest. Bar, 100 μ m. (h) Growth of *Rhizobium*-inoculated plants after 40 d in the absence of mineral nitrogen reveals a deficient nitrogen fixation phenotype for *MtHAP2-1* RNAi and 35S:*MtmiR169-a* plants compared with control plants.

Figure 3. A *miR169-a* precursor induces cleavage of *MtHAP2-1* mRNA. (a) Schematic representation of the *MtHAP2-1* gene with intron (line)–exon (boxes) structure and position of the two putative miR169 recognition sites (dotted lines) in the 3' UTR of the gene. Alignment of miR169 with the two miR169 recognition sites in *MtHAP2-1*; conserved nucleotides are indicated in bold on the *MtHAP2-1* sequence. Arrow shows predominant cleavage site revealed by 5' RACE-PCR. Underlined nucleotides are those changed by site-directed mutagenesis in the miR169-resistant *MtHAP2-1* mutant (miR169-R *MtHAP2-1*). (b) Most probable stem-loop structure of *MtmiR169-a*, indicating position of mature miR169 (line) and complementary miR169* (dotted line) sequences.

promoter. This led to an increased accumulation not only of the miR169 precursor (*pre-miR169-a*) (data not shown) but also of the mature miR169 (Fig. 4a), confirming that *MtmiR169-a* encodes microRNA 169. Evidence for miR169-mediated cleavage of *MtHAP2-1* mRNA was then obtained by 5' RACE-PCR experiments, which showed that cleavage takes place predominantly at the first miR169 recognition site (Fig. 3a). Overexpression of *MtmiR169-a* in transgenic roots led to a 3.5-fold reduction of *MtHAP2-1* transcript levels (Fig. 4b), confirming unambiguously that miR169 has a major impact on *MtHAP2-1* expression. Importantly, the nodulation phenotype of these roots was remarkably similar to that observed with an *MtHAP2-1* RNAi construct (Fig. 2e,f), indicating that miR169 regulation of *MtHAP2-1* is functioning in planta, and that the control of the precise level of *MtHAP2-1* mRNA is crucial for nodule development.

Real-time RT-PCR analysis showed that *MtmiR169-a* expression is undetectable in roots, expressed in young developing nodules, and maximal in 10-d-old functional nodules (Fig. 4c). This contrasts strikingly with *MtHAP2-1*, whose expression is also nodule specific but maximal in young developing nodules, and then decreases as the nodule becomes functional (Fig. 4c). A *pMtmiR169-a::GUS* fusion revealed that this miRNA is present in adjacent tissues to those expressing *MtHAP2-1* mRNA (Fig. 5a,b). Strikingly, *MtmiR169-a* transcript levels are highest in the infection zone (proximal to the meristem) (Fig. 5a,b), in which only lower levels of *MtHAP2-1* are detected (Fig. 1). Conversely, *MtmiR169-a* promoter activity could not be detected in the meristematic zone in which most *MtHAP2-1* mRNA is localized (Fig. 5b). These data are therefore consistent with a role of miR169 in restricting *MtHAP2-1* transcripts to the nodule meristematic zone.

To evaluate the functional relevance of this miR169-

mediated post-transcriptional regulation, a modified form of the *MtHAP2-1* gene, mutated in both miR169 recognition sites (miR169-resistant *MtHAP2-1*) (Fig. 3a) was expressed under the control of its own promoter, in the presence or absence of the p35S:*MtmiR169-a* construct. No difference was observed whether the p35S:*MtmiR169-a* construct was present or not (data not shown), suggesting that the mutated *MtHAP2-1* construct is indeed miR169 resistant. Nodules that developed on these roots showed no alterations in tissue differentiation or in nodule zonation (Fig. 5c) and seemed functional according to plant development in the absence of mineral nitrogen (data not shown). The miR169-resistant *MtHAP2-1* expressed under its own promoter can therefore restore nodulation, at least partly. This strongly suggests that the altered nodulation phenotype observed in p35S:*MtmiR169-a* roots is mainly mediated by *MtHAP2-1* and probably not by other putative miR169 targets, although a minor contribution cannot be excluded. The growth, of microRNA169-resistant nodules was, however, significantly reduced compared with control nodules (Fig. 5c,d). This result shows that when *MtHAP2-1* mRNA is not degraded by microRNA169 in the tissues surrounding the meristematic

Figure 4. miR169 cleaves *MtHAP2-1* in planta and *MtHAP2-1* and the *MtmiR169-a* are oppositely regulated during nodulation. (a) Northern blot analysis of mature 21-bp miR169 expression in transgenic roots transformed with either empty vector control (C) or the 35S:*MtmiR169-a* constructs (miR). (b) Real-time RT-PCR analysis of *MtHAP2-1* expression in *MtHAP2-1* RNAi and 35S:*MtmiR169-a* compared with empty vector controls. Results are expressed as a percentage of the expression in control roots. (c) Real-time RT-PCR analysis of the expression pattern of *MtmiR169-a* (black boxes) and *MtHAP2-1* (striped boxes) during nodule development. Relative expression levels are expressed in arbitrary units; the scale for *MtmiR169-a* is on the left while the scale for *MtHAP2-1* is on the right.

Figure 5. MtmiR169-a expression pattern in mature nodules and effects of a *MtHAP2-1* miR169-resistant construct on nodule development. (a,b) Expression analysis of *MtmiR169-a* in 14-d-old nodules revealed by promoter:GUS analysis. *b* is a magnification of the distal part of the nodule depicted in *a*, showing the expression of *MtmiR169-a* in the infection zone (II) and the absence of promoter activity in the meristematic region (I) of the nodule. Description of nodule zones is according to Vasse et al. (1990). Bar, 100 μ m. (c–d) Images of longitudinal sections through 20-d-old nodules are an overlay of fluorescence images acquired with two different wavelengths, showing nodule development on transgenic roots expressing either a miR169-resistant *MtHAP2-1* (miR169 R-HAP2-1) gene under the control of its own promoter (c), or an empty vector control (control) (d). Bar, 100 μ m.

zone, nodule elongation is affected, suggesting an inhibitory role of *MtHAP2-1* on cell differentiation and/or an indirect role on meristematic activity. This is coherent with the central role in cell proliferation and early development that has been shown for members of the CCAAT-box-binding family in other systems. Conditional CBF-B mutants in mouse cells, for example, strongly affect cell division and embryonic development (Bhattacharya et al. 2003), and *lec1* mutants are embryo lethal in *A. thaliana* (Lotan et al. 1998).

Our data show that *MtHAP2-1*, a novel symbiotically specialized CBF-B subunit, is essential for nodule meristematic persistence in *M. truncatula*, and that microRNA169 confers spatial and temporal accuracy to this developmental program. It is interesting to note that other TFs shown to specify shoot or root meristem development, such as CUC1-3 (Laufs et al. 2004), PHB, and PHV (Williams et al. 2005), or ICU4 (Ochando et al. 2006), are also the targets of microRNA mediated mRNA degradation. This additional level of regulation probably reflects the vital importance of meristems for plant development.

In conclusion, we propose that miR169-mediated regulation of *MtHAP2-1* expression leads to a critical spatial and temporal restriction of this TF to the nodule meristematic zone, thereby allowing correct tissue identity and the transition from meristematic to differentiated cells.

Materials and methods

Biological material

M. truncatula Gaertn cv. Jemalong genotype A17 was grown aeroponically as described in Journet et al. (2001), and inoculated with the *Sino-*

rhizobium meliloti strain RCR2011 pXLGD4 (GMI 6526) (Ardourel et al. 1994) as described previously (El Yahyaoui et al. 2004).

Expression analysis

mRNA in situ hybridizations were performed as described in Gamas et al. (1998). Image processing and analysis were performed using the Image-Pro plus software (Media Cybernetics) and the Diatrack software (Semasopht). For real-time RT-PCR experiments, RNA was isolated using the SV total RNA extraction kit (Promega), and quality checked using a Bioanalyser (Agilent). Three micrograms of total RNA were reverse-transcribed using the SuperScript II enzyme from Invitrogen. Real-time PCR was then performed on a LightCycler (Roche), using Syber-Green and specific primers. For *MtHAP2-1*, we used *MtHAP2-1R* and *MtHAP2-1F* (see Supplementary Table 1) while expression of the *pre-MtmiR169-a* was analyzed using miR169F and miR169R that amplify the stem-loop structure (249 base pairs [bp]) (Fig. 3b). Data were normalized using the *EF1- α* gene as described in El Yahyaoui et al. (2004). The specificity of primer pairs was confirmed by sequencing of PCR amplicons and the analysis of dissociation curves. Histochemical GUS staining was performed as described in Journet et al. (1994).

5' RACE-PCR experiment and Northern blot analysis of small RNAs

5' RACE-PCR was performed as described in Hirsch et al. (2006) using primers *MtHAP2-1-3'* (adapter), CBF_{rice}-miR1 (inner), and CBF_{rice}-miR2 (outer) (see Supplementary Table 1).

For Northern blots of small RNAs, extractions were performed using the TRIzol reagent (Invitrogen).

Twenty micrograms of each RNA were subjected to electrophoresis on a 17% polyacrylamide denaturing gel and electroblotted onto Hybond N⁺ membranes (GE Healthcare) using a Miniprotein II system (Bio-Rad). Blots were hybridized with miR169, miR166, or U6 probes as described in Hirsch et al. (2006).

Root transformation

Root transformation using *Agrobacterium rhizogenes* was performed as described in Boisson-Dernier et al. (2005). Composite plants were subsequently transferred to growth pouches supplemented with nitrogen-deprived medium as in Gallusci et al. (1991) and inoculated with *S. meliloti* strain RCR2011 pXLGD4 (Ardourel et al. 1994).

Constructions and vectors

All primers are listed in Supplementary Table 1.

For all root transformation experiments, we used a Pgreen (<http://www.pgreen.ac.uk>)-based vector with a modified polylinker, pPex (L. Sauviac, unpubl.).

For GUS fusion constructs, pPEX was modified by adding a GUS cassette from the pBS-GUS vector (Vernoud et al. 1999).

A 2.4-kb sequence upstream of the stem-loop structure of *MtmiR169-a* (Fig. 3b) was amplified using Pfx polymerase (Invitrogen) and primers *MtmiR169-5'* and *Pre-miR169-3'* and cloned into pPEX-GUS.

For RNAi experiments, the pPex vector was modified by introducing the intron from the pRNAi vector (Limpens et al. 2003) in addition to the cloning of the *DsRED* gene (under the control of the ubiquitin promoter) from the Predroot (Limpens et al. 2003) vector. A 298-bp region of the 3' UTR of the *MtHAP2-1* gene showing no significant homology with other known *MtHAP2* genes was amplified using primers CBF RNAi-5' and CBF RNAi-3' and cloned into the pPexRNAi vector.

For *MtmiR169-a* overexpression, a fragment located just outside of the stem-loop structure (Fig. 3b) was amplified with Pfx polymerase (Invitrogen) using miR169-5' and miR169-3' primers, and subsequently cloned into pPex.

For the miRNA-resistant version of *MtHAP2-1*, the gene including 2.3 kb of the upstream sequence was amplified with Pfx polymerase (Invitrogen) using the primers *ProMtHAP2-1-5'* and *MtHAP2-1-3'*, and then cloned into pPex. Mutagenesis of the two miR169 recognition sites was

performed by PCR amplification of this plasmid using Pfx polymerase (Invitrogen) and the following primers: first mutated miR169 recognition site (Fig. 3a): miR1muta-5' and miR1muta-3'; second miR169 recognition site (Fig. 3a): miR2muta-5' and miR2muta-3'. DpnI digestion was used to eliminate nonmutagenized plasmid prior to cloning.

Accession numbers: *MtHAP2-1*: MtC10582 (MENS database; <http://medicago.toulouse.inra.fr/Mt/EST>) and TC95981 (TIGR database; http://www.tigr.org/tigr-scripts/tgi/T_index.cgi?species=medicago).

Pre-MtmiR169-a: BAC AC148485.10 (NCBI databases; <http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=PubMed>).

Acknowledgments

We thank Laurent Sauviac (LIPM) for his kind gift of the pPex vector, René Geurts for his kind gift of the pRNAi and pRedroot vectors, Ton Timmers (LIPM) for fruitful discussions and advice concerning microscopy and imaging techniques, and Jean Marie Prosperi (INRA, Montpellier) for kindly providing seed stocks of *M. truncatula* A17. In addition, we are particularly grateful to David Barker, Jean Dénarié, Giles Oldroyd, and Hervé Vaucheret for critical reading of the manuscript. We thank Sakari Kaupinen (Exiqon, Denmark) for the kind gift of locked nucleic acid-modified oligonucleotides for detection of miR169. This work was supported by the European Community FP6 RIBOREG Project (LSHG-CT-2003503022) and FP6 Grain Legume Integrated Project (FP6-2002-FOOD-1-506223).

References

- Ardourel, M., Demont, N., Debelle, F., Maillet, F., de Billy, F., Prome, J.C., Denarie, J., and Truchet, G. 1994. *Rhizobium meliloti* lipooligosaccharide nodulation factors: Different structural requirements for bacterial entry into target root hair cells and induction of plant symbiotic developmental responses. *Plant Cell* **6**: 1357–1374.
- Ben-Naim, O., Eshed, R., Parnis, A., Teper-Bamnolker, P., Shalit, A., Coupland, G., Samach, A., and Lifschitz, E. 2006. The CCAAT binding factor can mediate interactions between CONSTANS-like proteins and DNA. *Plant J.* **46**: 462–476.
- Bhattacharya, A., Deng, J.M., Zhang, Z., Behringer, R., de Crombrughe, B., and Maity, S.N. 2003. The B subunit of the CCAAT box binding transcription factor complex (CBF/NF-Y) is essential for early mouse development and cell proliferation. *Cancer Res.* **63**: 8167–8172.
- Boisson-Dernier, A., Andriankaja, A., Chabaud, M., Niebel, A., Journet, E.P., Barker, D.G., and de Carvalho-Niebel, F. 2005. MtENOD11 gene activation during rhizobial infection and mycorrhizal arbuscule development requires a common AT-rich-containing regulatory sequence. *Mol. Plant Microbe Interact.* **18**: 1269–1276.
- El Yahyaoui, F., Kuster, H., Ben Amor, B., Hohnjec, N., Pühler, A., Becker, A., Gouzy, J., Vernié, T., Gough, C., Niebel, A., et al. 2004. Expression profiling in *Medicago truncatula* identifies more than 750 genes differentially expressed during nodulation, including many potential regulators of the symbiotic program. *Plant Physiol.* **136**: 3559–3576.
- Gallusci, P., Dedieu, A., Journet, E.P., Hugué, T., and Barker, D.G. 1991. Synchronous expression of leghaemoglobin genes in *Medicago truncatula* during nitrogen-fixing root nodule development and response to exogenously supplied nitrate. *Plant Mol. Biol.* **17**: 335–349.
- Gamas, P., de Billy, F., and Truchet, G. 1998. Symbiosis-specific expression of two *Medicago truncatula* nodulin genes, MtN1 and MtN13, encoding products homologous to plant defense proteins. *Mol. Plant Microbe Interact.* **11**: 393–403.
- Geurts, R., Fedorova, E., and Bisseling, T. 2005. Nod factor signaling genes and their function in the early stages of *Rhizobium* infection. *Curr. Opin. Plant Biol.* **8**: 346–352.
- Gusmaroli, G., Tonelli, C., and Mantovani, R. 2002. Regulation of novel members of the *Arabidopsis thaliana* CCAAT-binding nuclear factor Y subunits. *Gene* **283**: 41–48.
- Hirsch, J., Lefort, V., Vankersschaver, M., Boualem, A., Lucas, A., Thermes, C., d'Aubenton-Carafa, Y., and Crespi, M. 2006. Characterization of 43 non-protein-coding mRNA genes in *Arabidopsis*, including the MIR162a-derived transcripts. *Plant Physiol.* **140**: 1192–1204.
- Jones-Rhoades, M.W. and Bartel, D.P. 2004. Computational identification of plant microRNAs and their targets, including a stress-induced miRNA. *Mol. Cell* **14**: 787–799.
- Jones-Rhoades, M.W., Bartel, D.P., and Bartel, B. 2006. MicroRNAs and their regulatory roles in plants. *Annu. Rev. Plant Biol.* **57**: 19–53.
- Journet, E.P., Pichon, M., Dedieu, A., de Billy, F., Truchet, G., and Barker, D.G. 1994. *Rhizobium meliloti* Nod factors elicit cell-specific transcription of the ENOD12 gene in transgenic alfalfa. *Plant J.* **6**: 241–249.
- Journet, E.P., El-Gachtouli, N., Vernoud, V., de Billy, F., Pichon, M., Dedieu, A., Arnould, C., Morandi, D., Barker, D.G., and Gianinazzi-Pearson, V. 2001. *Medicago truncatula* ENOD11: A novel RPRP-encoding early nodulin gene expressed during mycorrhization in arbuscule-containing cells. *Mol. Plant Microbe Interact.* **14**: 737–748.
- Kwong, R.W., Bui, A.Q., Lee, H., Kwong, L.W., Fischer, R.L., Goldberg, R.B., and Harada, J.J. 2003. LEAFY COTYLEDON1-LIKE defines a class of regulators essential for embryo development. *Plant Cell* **15**: 5–18.
- Laufs, P., Peaucelle, A., Morin, H., and Traas, J. 2004. MicroRNA regulation of the CUC genes is required for boundary size control in *Arabidopsis* meristems. *Development* **131**: 4311–4322.
- Lee, H., Fischer, R.L., Goldberg, R.B., and Harada, J.J. 2003. *Arabidopsis* LEAFY COTYLEDON1 represents a functionally specialized subunit of the CCAAT binding transcription factor. *Proc. Natl. Acad. Sci.* **100**: 2152–2156.
- Li, X. and Zhang, Y.Z. 2005. Computational detection of microRNAs targeting transcription factor genes in *Arabidopsis thaliana*. *Comput. Biol. Chem.* **29**: 360–367.
- Limpens, E., Franken, C., Smit, P., Willemse, J., Bisseling, T., and Geurts, R. 2003. LysM domain receptor kinases regulating rhizobial Nod factor-induced infection. *Science* **302**: 630–633.
- Lotan, T., Ohto, M., Yee, K.M., West, M.A., Lo, R., Kwong, R.W., Yamagishi, K., Fischer, R.L., Goldberg, R.B., and Harada, J.J. 1998. *Arabidopsis* LEAFY COTYLEDON1 is sufficient to induce embryo development in vegetative cells. *Cell* **93**: 1195–1205.
- Maity, S.N. and de Crombrughe, B. 1998. Role of the CCAAT-binding protein CBF/NF-Y in transcription. *Trends Biochem. Sci.* **23**: 174–178.
- Masiero, S., Imbriano, C., Ravasio, F., Favaro, R., Pelucchi, N., Gorla, M.S., Mantovani, R., Colombo, L., and Kater, M.M. 2002. Ternary complex formation between MADS-box transcription factors and the histone fold protein NF-YB. *J. Biol. Chem.* **277**: 26429–26435.
- Miyoshi, K., Ito, Y., Serizawa, A., and Kurata, N. 2003. OsHAP3 genes regulate chloroplast biogenesis in rice. *Plant J.* **36**: 532–540.
- Ochando, I., Jover-Gil, S., Ripoll, J.J., Candela, H., Vera, A., Ponce, M.R., Martinez-Laborda, A., and Micol, J.L. 2006. Mutations in the microRNA complementarity site of the INCURVATA4 gene perturb meristem function and adaxialize lateral organs in *Arabidopsis*. *Plant Physiol.* **141**: 607–619.
- Stacey, G., Libault, M., Brechenmacher, L., Wan, J., and May, G.D. 2006. Genetics and functional genomics of legume nodulation. *Curr. Opin. Plant Biol.* **9**: 110–121.
- Vasse, J., de Billy, F., Camut, S., and Truchet, G. 1990. Correlation between ultrastructural differentiation of bacteroids and nitrogen fixation in alfalfa nodules. *J. Bacteriol.* **172**: 4295–4306.
- Vernoud, V., Journet, E.P., and Barker, D. 1999. MtENOD20, a nod factor inducible molecular marker for root cortical cell activation. *Mol. Plant Microbe Interact.* **12**: 604–614.
- Williams, L., Grigg, S.P., Xie, M., Christensen, S., and Fletcher, J.C. 2005. Regulation of *Arabidopsis* shoot apical meristem and lateral organ formation by microRNA miR166g and its AtHD-ZIP target genes. *Development* **132**: 3657–3668.

MtHAP2-1* is a key transcriptional regulator of symbiotic nodule development regulated by microRNA169 in *Medicago truncatula

Jean-Philippe Combier, Florian Frugier, Françoise de Billy, et al.

Genes Dev. 2006, **20**:

Access the most recent version at doi:[10.1101/gad.402806](https://doi.org/10.1101/gad.402806)

Supplemental Material

<http://genesdev.cshlp.org/content/suppl/2006/10/26/20.22.3084.DC1>

References

This article cites 28 articles, 11 of which can be accessed free at:
<http://genesdev.cshlp.org/content/20/22/3084.full.html#ref-list-1>

License

Email Alerting Service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).