RESEARCH COMMUNICATION

Maternal microRNAs are essential for mouse zygotic development

Fuchou Tang,^{1,4} Masahiro Kaneda,^{1,4}
 Dónal O'Carroll,² Petra Hajkova,¹
 Sheila C. Barton,¹ Y. Andrew Sun,³ Caroline Lee,¹
 Alexander Tarakhovskiy,² Kaiqin Lao,^{3,6} and
 M. Azim Surani^{1,5}

¹Wellcome Trust/Cancer Research UK Gurdon Institute of Cancer and Developmental Biology, University of Cambridge, Cambridge CB2 1QN, United Kingdom; ²The Laboratory for Lymphocyte Signaling and the Laboratory of Molecular Immunology, The Rockefeller University, New York, New York 10021, USA; ³Advanced Research Technology, Applied Biosystems, Foster City, California 94404, USA

MicroRNAs (miRNAs) have important roles in diverse cellular processes, but little is known about their identity and functions during early mammalian development. Here, we show the effects of the loss of maternal inheritance of miRNAs following specific deletion of Dicer from growing oocytes. The mutant mature oocytes were almost entirely depleted of all miRNAs, and they failed to progress through the first cell division, probably because of disorganized spindle formation. By comparing single-cell cDNA microarray profiles of control and mutant oocytes, our data are compatible with the notion that a large proportion of the maternal genes are directly or indirectly under the control of miRNAs, which demonstrates that the maternal miRNAs are essential for the earliest stages of mouse embryonic development.

Supplemental material is available at <http://www.genesdev.org>.

Received November 20, 2006; revised version accepted January 29, 2007.

MicroRNAs (miRNAs) are a large family of short non-coding RNAs (17–25 nucleotides) (He and Hannon 2004). A key function of miRNAs is to repress expression of their target genes through sequence complementation, which reduces the abundance of the target mRNAs and/or inhibits their translation (Bartel 2004; Bagga et al. 2005). MiRNA genes are first transcribed into miRNA primary transcripts by RNA polymerase II (Kim 2005). These primary transcripts are then processed into miRNA precursors by the Drosha/DGCR8 complex and transported from the nucleus to the cytoplasm. Finally, Dicer processes the miRNA precursors into mature miRNAs. From previous studies, Dicer seems to be critical for early mouse development since its loss of function is

embryonic lethal at embryonic day 7.5 (E7.5) (Bernstein et al. 2003).

In this study, we have examined the role of miRNAs in the mouse oocyte. The mature oocyte contains a number of molecules that are manufactured during oocyte maturation and utilized during early stages of development before activation of the embryonic genome (Dean 2002). It is likely that miRNAs would also be present in the oocyte, but no information is yet available in the mouse. The purpose of this study was to determine if there is significant inheritance of maternal miRNAs in mammalian zygotes, and to investigate if they play a critical role in early mammalian development. We have investigated how the loss of Dicer affects synthesis of miRNA during oocyte maturation and their impact on mRNA and early development.

Results and Discussion

First, we decided to investigate if there is significant biogenesis of miRNAs in developing oocytes, and their inheritance in the zygote. We therefore examined expression of miRNAs in single cells during oogenesis by a real-time PCR-based miRNA expression profiling method that we recently developed (C. Chen et al. 2005; Tang et al. 2006a,b). We compared the miRNA expression profiles of growing oocytes obtained from females 15–16 d after birth (postnatal days 15–16 [P15–P16]) and at P20–P21, and of mature oocytes from adult females. This analysis revealed dynamic changes in miRNA expression during oogenesis (Fig. 1; Supplementary Table S1).

Next, we compared expression of miRNA in the mature oocyte with miRNAs in the zygote, which showed essentially the same miRNA expression pattern in both these cells (Fig. 1; Supplementary Tables S1, S2). This observation indicates that the miRNAs we detected in the zygote are probably maternally inherited from oocytes and not transcribed in the zygote itself.

We then went on to determine if there are significant changes in miRNAs during early embryonic development. Indeed, we found that the miRNA expression profile of the zygote underwent dynamic changes during early embryonic development when examined through to the eight-cell stage (Fig. 1; Supplementary Table S2). Notably, we found that the total amount of miRNA is down-regulated by 60% between one- and two-cell-stage embryos ($P < 0.001$) (Fig. 2A; Supplementary Table S3), suggesting that a very significant proportion of the maternally inherited miRNAs present in the zygote is probably actively degraded during the first cell division. Some of these miRNAs were selectively lost by >95% (Supplementary Table S2). This is unexpected because there are no reports to show that miRNAs can be actively degraded *in vivo* under physiological conditions, although maternal mRNAs are globally degraded at this time (Hamatani et al. 2004). We confirmed our findings of the loss of maternal miRNAs by RNA *in situ* hybridization with LNA probes, which has been shown to be able to detect mature miRNA expression (Supplementary Fig. S2C,D; Supplementary Table S4; Kloosterman et al. 2006). Among the maternal miRNAs in the zygote, the most abundant are let-7 family miRNAs. They show dynamic regulation during oogenesis and early embry-

[**Keywords:** Maternal microRNAs; Dicer; oocyte; zygote]

⁴These authors contributed equally to this work.

Corresponding authors.

⁵E-MAIL as10021@mole.bio.cam.ac.uk; FAX 44-1223-334089.

⁶E-MAIL laokq@appliedbiosystems.com; FAX (650) 638-6343.

Article is online at <http://www.genesdev.org/cgi/doi/10.1101/gad.418707>.

Figure 1. Unsupervised hierarchical clustering heat map of miRNA expression profile of oocytes and early embryos. The cluster heat map was produced using expression levels (Ct value) of 214 miRNAs. Correlation coefficient was used as a similarity measure and a complete linkage method was used as a clustering method. Note that a higher Ct value means a lower expression level.

onic development (Supplementary Fig. S2A,B). The most abundant cluster is the miR-17-92 cluster. This miRNA cluster has been shown to be “oncomirs,” which is involved in cell proliferation (He et al. 2005; O’Donnell et al. 2005). Their abundance significantly increased during oogenesis and was inherited by the zygote and increased again after the two-cell embryo stage (Supplementary Fig. S3).

Further investigations showed that the total miRNA in a four-cell-stage embryo was ~2.2 times higher than the levels detected in a two-cell-stage embryo (Fig. 2A; Supplementary Table S3). This suggests that there is de novo expression of miRNAs between the two- and four-cell stages of development. Amongst the most significant miRNAs that are up-regulated in a four-cell-stage embryo are those from the miR-290 cluster, namely miR-290 to miR-295 (Houbaviy et al. 2003). Compared with the two-cell-stage embryos, these miRNAs were up-regulated by 15-fold and by 24-fold in four-cell- and eight-cell-stage embryos, respectively (Supplementary Fig. S4A,B; Supplementary Table S5). Thus, miRNAs from the miR-290 cluster are amongst the earliest to be expressed during early mouse embryonic development.

To determine if there are significant differences in miRNAs in individual blastomeres, we separated and compared the miRNA expression profiles of the two individual blastomeres from individual two-cell-stage embryos. We found that both blastomeres examined separately have essentially the same miRNA expression profile (Supplementary Fig. S5A; Supplementary Table S10). Furthermore, the total amount of miRNAs in the two blastomeres together was only ~40% of that detected in the zygote. This shows that the miRNA profiling method is reliable and confirms that a significant proportion of the miRNAs is degraded in embryos between the one- and two-cell stages. Similarly, the miRNA ex-

pression profiles of the four individual blastomeres from a four-cell-stage embryo are essentially the same (Supplementary Fig. S5B; Supplementary Table S10). These observations indicate that the individual blastomeres at two-cell and four-cell embryo stages have similar if not identical miRNA expression profiles.

Next, we asked if these maternally inherited miRNAs are functionally important for early development. To investigate this aspect, we used mice carrying the *Dicer* floxed conditional allele, where exon 23 of *Dicer* locus is flanked by two *loxP* loci (Yi et al. 2006). These mice with the floxed allele of *Dicer* were mated with *Zp3-Cre* transgenic mice, which express Cre recombinase under the control of Zona pellucida glycoprotein 3 promoter (*Zp3*) in the growing oocyte. We thus generated [*Dicer*^{-/*Flox*}, *Zp3-Cre*] animals. It is known that *Zp3* expression is detected only in growing oocytes from about P5, which allowed us to delete *Dicer* specifically from maturing oocytes (de Vries et al. 2000). We would therefore expect loss of *Dicer* function in growing oocytes, and this would in turn block biogenesis of miRNAs at this stage. To establish how the loss of *Dicer* from the growing oocytes affects miRNAs in oocytes, we examined the miRNA profile in the mutant oocytes. We found that most, if not all, miRNAs were essentially lost from the oocytes lacking *Dicer* (Supplementary Fig. S6; Supplementary Table S1). We also found that the loss of the *Dicer* allele from oocytes rendered the females infertile.

Figure 2. (A) Total amount of miRNA in mature oocytes and early embryos. The error bars were the standard deviations calculated from three independent samples. (B) Morphology of mutant embryos from *Dicer* knockout oocytes, compared with wild-type control embryos at E1.5. Embryos above the red line are mutant, while embryos below the red line are wild-type controls.

To determine the reason for infertility and when the effect manifests itself, we recovered E1.5 embryos from control and mutant oocytes following the deletion of *Dicer* from the growing oocytes that were fertilized by sperm from wild-type males (Fig. 2B). We found that the control females had healthy two-cell-stage embryos as expected, but all of the mutant oocytes failed to proceed through the first cell division, and about half of them were fragmented (three litters, $n = 28$). Thus, maternal *Dicer* and miRNAs in the oocyte are crucial for the earliest stages of embryonic development.

To determine more precisely the stage at which the loss of *Dicer* from oocytes starts to affect oogenesis and embryonic development, we examined the expression of some of the key genes by single-cell cDNA analysis at three different stages during oogenesis (Rajkovic et al. 2004). We found that expression of a number of genes, including *Oct4*, *Fragilis*, *Stella*, *C-mos*, *Bnc1*, *H2AX*, *H1foo*, *SCP3*, *Nobox*, *Gata4*, and *RFPL4*, was unaffected in growing oocytes at P15–P16 (Supplementary Fig. S1). The overall morphology of the mutant oocytes was also indistinguishable from that of control oocytes at this stage. Next, we checked expression of the same genes in mature ovulated oocytes, which we obtained from [*Dicer*^{-flox}, *Zp3-Cre*] females that were mated with vasectomized males. We again found that the *Dicer* mutant oocytes were morphologically indistinguishable from control oocytes in their appearance, maturity, and size. The overall numbers of ovulated mutant oocytes were also indistinguishable from those obtained from control females. However, we found that the *Dicer* mutant oocytes showed higher expression of *C-mos* and *H2AX* compared with the levels we found in control oocytes (Supplementary Fig. S1). In the E0.5 embryos, genes including *H1foo* and *SCP3* also showed higher expression compared with the control. Therefore, mature oocytes lacking in *Dicer*, and consequently miRNAs, already showed an effect on mRNAs.

To determine if the miRNAs are involved in maturation of the oocyte, we also checked the chromosome and spindle organization (Lefebvre et al. 2002; Terret et al. 2003). We found that compared with control oocytes, the *Dicer* mutant oocytes showed reduced and disorganized spindles, and the chromosomes were also not aligned

Figure 3. Spindle in mutant mature oocytes (D–F) compared with wild-type mature oocytes (A–C). The spindle was stained with rat monoclonal anti-tubulin (YL1/2) antibody (green), and the chromosome was stained with DAPI (blue).

Figure 4. Unsupervised hierarchical clustering heat map of wild-type and *Dicer* knockout single-oocyte cDNA microarray. All genes that are differentially expressed between wild-type (left six columns) and *Dicer* knockout oocytes (right six columns) are shown. Clustering is based on the log₂ of chemiluminescent intensities.

altogether properly (Fig. 3). This indicates that loss of *Dicer* and miRNAs affects the spindle organization of mature oocytes.

To obtain more comprehensive information on how miRNAs affect embryonic development at the whole-genome scale, we prepared single-cell cDNAs from control and *Dicer* mutant oocytes for microarray analysis (Kurimoto et al. 2006). We found that compared with control oocytes, *Dicer* knockout oocytes showed an increase in the levels of many genes that are probably important during early development. The cDNA levels of more than one-third of the genes expressed in oocytes increased (>1.5-fold) after the loss of *Dicer* and miRNAs (Fig. 4; Supplementary Table S7). We confirmed the microarray data by quantitative PCR on single-cell cDNAs (Supplementary Table S8). This analysis shows that miRNAs that are expressed during oogenesis profoundly shape the gene expression profile of the mature oocyte. Since we did not find significant enrichment of predicted target genes of miRNAs amongst the expression-increased genes in the *Dicer* knockout oocytes, our findings are in agreement with a similar study in zebrafish, where the microarray data of *Dicer* mutant and wild-type embryos were also compared (Giraldez et al. 2006).

To further understand the molecular basis for the role of maternal miRNAs on early development, we compared our microarray data with the predicted target genes of maternal miRNAs using a bioinformatics approach. We searched the sequence complement to the “seed” region of miRNAs in the 3′ untranslated region (UTR) of all the 30,677 genes with defined 3′UTRs represented on the ABI chip. We found that the genes that have no target sites in their 3′UTRs for the 101 maternal miRNAs have

4.2-fold more chance to be coexpressed in oocytes with these miRNAs ($P = 2.2\text{E-}16$) (Supplementary Tables S9, S11). This indicates that miRNAs expressed in the oocyte functionally shape the gene expression profile, which is in accordance with the reports that genes expressed in a particular tissue tend to avoid being the targets of coexpressed miRNAs in the same tissue (Farh et al. 2005; Stark et al. 2005; Sood et al. 2006). By combining the analysis of the single-cell miRNA profile with single-cell cDNA microarrays, we confirmed at the resolution of single cells that some of the genes expressed in the oocyte had a strong tendency to avoid being the target sites of coexpressed miRNAs.

As it is known that miRNAs also apparently have an effect on the expression of repetitive elements (Fukagawa et al. 2004; Kanellopoulou et al. 2005), we examined the expression levels of IAP (intracisternal A particle element; an endogenous retrovirus), LINE1 (long interspersed nuclear element 1; a nonretrovirus retrotransposon), and MTs (mouse transcript; a nonautonomous retrotransposon), which are the most abundant repetitive elements in the mouse oocyte. However, quantitative real-time RT-PCR analysis showed no significant differences in their expression between *Dicer* mutant and control oocytes (Fig. 5A). We also investigated the DNA methylation status of IAP and LINE1 by the

bisulphite genomic sequencing method and found no significant differences between *Dicer* mutant and control oocytes (Fig. 5B). Thus the phenotype of the *Dicer* mutant oocyte is due to the direct or indirect effects of the loss of miRNAs, not the derepression of repetitive elements.

In summary, we have generated a comprehensive miRNA expression profile for growing oocytes and embryos up to the eight-cell stage. Interestingly, unlike the observations in zebrafish and *Xenopus* (P.Y. Chen et al. 2005; Watanabe et al. 2005), we found abundant maternally inherited miRNAs in mouse zygotes, among which let-7 family miRNAs are the most abundant. It seems that sperm-borne miRNAs do not contribute significantly to miRNAs in the zygote (Amanai et al. 2006). Furthermore, in zebrafish it was shown that maternal miRNAs are dispensable for early embryonic development, although the maternal *Dicer* is important because it is necessary for zygotic synthesis and expression of miR-430 (Giraldez et al. 2005, 2006). In the mouse, while there is a significant global loss of maternal miRNAs between the one- and two-cell stages of development, de novo synthesis of miRNAs commences at the two-cell stage. This includes expression of miR-290 to miR-295, which are the first embryonic miRNAs to be detected. It is noteworthy that mir-290 to mir-295 are also specifically expressed in embryonic stem cells, which may suggest their association with pluripotency (Houbaviy et al. 2003). We cannot formally rule out the possibility that the *Dicer* mutant phenotype is not solely due to loss of miRNAs because *Dicer* might directly play some unknown roles in chromatin formation. Amongst the genes that were detected at higher levels in the oocytes following the loss of *Dicer*, some of these effects may be due to secondary consequences of miRNA depletion. Nevertheless, this study provides evidence that the maternal inheritance of miRNAs is crucial for early mammalian development. The detailed analysis of the oocyte miRNAs and their impact on mRNAs we present here may help to elucidate their precise roles in early mouse development in the future.

Materials and methods

Embryos and knockout mice

Embryos before implantation were recovered from F1 (C57BL/6 × CBA) females mated with F1 male mice (Nagy et al. 2003). Oocytes were isolated from F1 female mice. All the “mature oocytes” we mentioned in the text are ovulated mature oocytes.

The knockout mice carrying the *Dicer* floxed allele was described previously (Yi et al. 2006). Basically, exon 23 of the *Dicer* locus was floxed by two loxP loci (referred to as *Dicer*^{Flox}). The *Dicer*^{Flox/Flox} mice were mated with *Zp3-Cre* transgenic mice, which express Cre recombinase under the control of the Zona pellucida glycoprotein 3 promoter (de Vries et al. 2000). Then the [*Dicer*^{Flox/+}, *Zp3-Cre*] female mice were mated with *Dicer*^{Flox/Flox} male mice. From this mating, we obtained [*Dicer*^{-/Flox}, *Zp3-Cre*] mice, and following the deletion of the floxed allele in the oocyte, we generate oocytes that are the null mutants for *Dicer*. The control mice we used are littermates with the genotype of [*Dicer*^{+/Flox}, *Zp3-Cre*], *Dicer*^{+/Flox}, or *Dicer*^{-/Flox}.

The details of the microRNA expression profiling assay, RNA in situ hybridization by LNA probe, immunostaining, single-cell cDNA, real-time PCR, microarray gene expression procedures and analysis, and bisulphite genomic sequencing can be found in the Supplemental Material. The data of the single-oocyte cDNA microarray were deposited in GeneBank (<http://www.ncbi.nlm.nih.gov/geo>). The accession number is GSE6806.

Figure 5. (A) Transcript abundance of repetitive elements in control and *Dicer* mutant oocytes measured by quantitative real-time RT-PCR. The error bars represent standard deviations calculated from two independent samples. (B) DNA methylation status of repetitive elements in control and *Dicer* mutant oocytes measured by bisulphite genomic sequencing. Filled circles represent methylated CpG and open circles represent nonmethylated CpG. Horizontally aligned circles represent a single DNA molecule. The overall percentage of methylated CpGs is shown below each group.

Acknowledgments

We thank Eric Miska, Anne McLaren, Kenneth Livak, Naoki Miyoshi, Katsuhiko Hayashi, Maria Elena Torres Padilla, Jie Na, David Adams, and James Smith for their helpful discussions and generous suggestions. We also thank W.N. de Vries and B.B. Knowles for the *Zp3-Cre* transgenic mice. This work was supported by grants from the Wellcome Trust and BBSRC to M.A.S. M.K. is supported by the Japanese Society for the Promotion of the Science (JSPS).

References

- Amanai, M., Brahmajosyula, M., and Perry, A.C. 2006. A restricted role for sperm-borne microRNAs in mammalian fertilization. *Biol. Reprod.* **75**: 877–884.
- Bagga, S., Bracht, J., Hunter, S., Massirer, K., Holtz, J., Eachus, R., and Pasquinelli, A.E. 2005. Regulation by let-7 and lin-4 miRNAs results in target mRNA degradation. *Cell* **122**: 553–563.
- Bartel, D.P. 2004. MicroRNAs: Fenomics, biogenesis, mechanism, and function. *Cell* **116**: 281–297.
- Bernstein, E., Kim, S.Y., Carmell, M.A., Murchison, E.P., Alcorn, H., Li, M.Z., Mills, A.A., Elledge, S.J., Anderson, K.V., and Hannon, G.J. 2003. Dicer is essential for mouse development. *Nat. Genet.* **35**: 215–217.
- Chen, C., Ridzon, D.A., Broomer, A.J., Zhou, Z., Lee, D.H., Nguyen, J.T., Barbisin, M., Xu, N.L., Mahuvakar, V.R., Andersen, M.R., et al. 2005. Real-time quantification of microRNAs by stem-loop RT-PCR. *Nucleic Acids Res.* **33**: e179.
- Chen, P.Y., Mannings, H., Slanchev, K., Chien, M., Russo, J.J., Ju, J., Sheridan, R., John, B., Marks, D.S., Gaidatzis, D., et al. 2005. The developmental miRNA profiles of zebrafish as determined by small RNA cloning. *Genes & Dev.* **19**: 1288–1293.
- de Vries, W.N., Binns, L.T., Fancher, K.S., Dean, J., Moore, R., Kemler, R., and Knowles, B.B. 2000. Expression of Cre recombinase in mouse oocytes: A means to study maternal effect genes. *Genesis* **26**: 110–112.
- Dean, J. 2002. Oocyte-specific genes regulate follicle formation, fertility and early mouse development. *J. Reprod. Immunol.* **53**: 171–180.
- Farh, K.K., Grimson, A., Jan, C., Lewis, B.P., Johnston, W.K., Lim, L.P., Burge, C.B., and Bartel, D.P. 2005. The widespread impact of mammalian microRNAs on mRNA repression and evolution. *Science* **310**: 1817–1821.
- Fukagawa, T., Nogami, M., Yoshikawa, M., Ikeno, M., Okazaki, T., Takami, Y., Nakayama, T., and Oshimura, M. 2004. Dicer is essential for formation of the heterochromatin structure in vertebrate cells. *Nat. Cell Biol.* **6**: 784–791.
- Giraldez, A.J., Cinalli, R.M., Glasner, M.E., Enright, A.J., Thomson, J.M., Baskerville, S., Hammond, S.M., Bartel, D.P., and Schier, A.F. 2005. MicroRNAs regulate brain morphogenesis in zebrafish. *Science* **308**: 833–838.
- Giraldez, A.J., Mishima, Y., Rihel, J., Grocock, R.J., Van Dongen, S., Inoue, K., Enright, A.J., and Schier, A.F. 2006. Zebrafish MiR-430 promotes deadenylation and clearance of maternal mRNAs. *Science* **312**: 75–79.
- Hamatani, T., Carter, M.G., Sharov, A.A., and Ko, M.S. 2004. Dynamics of global gene expression changes during mouse preimplantation development. *Dev. Cell* **6**: 117–131.
- He, L. and Hannon, G.J. 2004. MicroRNAs: Small RNAs with a big role in gene regulation. *Nat. Rev. Genet.* **5**: 522–531.
- He, L., Thomson, J.M., Hemann, M.T., Hernandez-Monge, E., Mu, D., Goodson, S., Powers, S., Cordon-Cardo, C., Lowe, S.W., Hannon, G.J., et al. 2005. A microRNA polycistron as a potential human oncogene. *Nature* **435**: 828–833.
- Houbaviy, H.B., Murray, M.F., and Sharp, P.A. 2003. Embryonic stem cell-specific microRNAs. *Dev. Cell* **5**: 351–358.
- Kanellopoulou, C., Muljo, S.A., Kung, A.L., Ganesan, S., Drapkin, R., Jenuwein, T., Livingston, D.M., and Rajewsky, K. 2005. Dicer-deficient mouse embryonic stem cells are defective in differentiation and centromeric silencing. *Genes & Dev.* **19**: 489–501.
- Kim, V.N. 2005. MicroRNA biogenesis: Coordinated cropping and dicing. *Nat. Rev. Mol. Cell Biol.* **6**: 376–385.
- Kloosterman, W.P., Wienholds, E., de Bruijn, E., Kauppinen, S., and Plasterk, R.H. 2006. In situ detection of miRNAs in animal embryos using LNA-modified oligonucleotide probes. *Nat. Methods* **3**: 27–29.
- Kurimoto, K., Yabuta, Y., Ohinata, Y., Ono, Y., Uno, K.D., Yamada, R.G., Ueda, H.R., and Saitou, M. 2006. An improved single-cell cDNA amplification method for efficient high-density oligonucleotide microarray analysis. *Nucleic Acids Res.* **34**: e42.
- Lefebvre, C., Terret, M.E., Djiane, A., Rassini, P., Maro, B., and Verlhac, M.H. 2002. Meiotic spindle stability depends on MAPK-interacting and spindle-stabilizing protein (MISS), a new MAPK substrate. *J. Cell Biol.* **157**: 603–613.
- Nagy, A., Gertsenstein, M., Vintersten, K., and Behringer, R. 2003. *Manipulating the mouse embryo*, 3rd ed., pp. 194–200. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- O'Donnell, K.A., Wentzel, E.A., Zeller, K.I., Dang, C.V., and Mendell, J.T. 2005. c-Myc-regulated microRNAs modulate E2F1 expression. *Nature* **435**: 839–843.
- Rajkovic, A., Pangas, S.A., Ballow, D., Suzumori, N., and Matzuk, M.M. 2004. NOBOX deficiency disrupts early folliculogenesis and oocyte-specific gene expression. *Science* **305**: 1157–1159.
- Soed, P., Krek, A., Zavolan, M., Macino, G., and Rajewsky, N. 2006. Cell-type-specific signatures of microRNAs on target mRNA expression. *Proc. Natl. Acad. Sci.* **103**: 2746–2751.
- Stark, A., Brennecke, J., Bushati, N., Russell, R.B., and Cohen, S.M. 2005. Animal microRNAs confer robustness to gene expression and have a significant impact on 3'UTR evolution. *Cell* **123**: 1133–1146.
- Tang, F., Hajkova, P., Barton, S.C., Lao, K., and Surani, M.A. 2006a. MicroRNA expression profiling of single whole embryonic stem cells. *Nucleic Acids Res.* **34**: e9.
- Tang, F., Hajkova, P., Barton, S.C., O'Carroll, D., Lee, C., Lao, K., and Surani, M.A. 2006b. 220-plex microRNA expression profile of a single cell. *Nat. Protoc.* **1**: 1154–1159.
- Terret, M.E., Lefebvre, C., Djiane, A., Rassini, P., Moreau, J., Maro, B., and Verlhac, M.H. 2003. DOC1R: A MAP kinase substrate that control microtubule organization of metaphase II mouse oocytes. *Development* **130**: 5169–5177.
- Watanabe, T., Takeda, A., Mise, K., Okuno, T., Suzuki, T., Minami, N., and Imai, H. 2005. Stage-specific expression of microRNAs during *Xenopus* development. *FEBS Lett.* **579**: 318–324.
- Yi, R., O'Carroll, D., Pasolli, H.A., Zhang, Z., Dietrich, F.S., Tarakhovskiy, A., and Fuchs, E. 2006. Morphogenesis in skin is governed by discrete sets of differentially expressed microRNAs. *Nat. Genet.* **38**: 356–362.

Maternal microRNAs are essential for mouse zygotic development

Fuchou Tang, Masahiro Kaneda, Dónal O'Carroll, et al.

Genes Dev. 2007, **21**:

Access the most recent version at doi:[10.1101/gad.418707](https://doi.org/10.1101/gad.418707)

**Supplemental
Material**

<http://genesdev.cshlp.org/content/suppl/2007/03/09/21.6.644.DC1>

References

This article cites 29 articles, 9 of which can be accessed free at:
<http://genesdev.cshlp.org/content/21/6/644.full.html#ref-list-1>

License

Freely available online through the Genes & Development Open Access option.

**Email Alerting
Service**

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).

Focused on your science.