

## RESEARCH COMMUNICATION

# Aberrant chromatin at genes encoding stem cell regulators in human mixed-lineage leukemia

Matthew G. Guenther,<sup>1</sup> Lee N. Lawton,<sup>1</sup> Tatiana Rozovskaia,<sup>2</sup> Garrett M. Frampton,<sup>1,3</sup> Stuart S. Levine,<sup>1</sup> Thomas L. Volkert,<sup>1</sup> Carlo M. Croce,<sup>4</sup> Tatsuya Nakamura,<sup>4</sup> Eli Canaani,<sup>2</sup> and Richard A. Young<sup>1,3,5</sup>

<sup>1</sup>Whitehead Institute for Biomedical Research, Cambridge, Massachusetts 02142, USA; <sup>2</sup>Department of Molecular Cell Biology, Weizmann Institute of Science, Rehovot 76100, Israel; <sup>3</sup>Department of Biology, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA; <sup>4</sup>Department of Molecular Virology, Immunology and Medical Genetics and Comprehensive Cancer Center, Ohio State University, Columbus, Ohio 43210, USA

**Mixed-lineage leukemia (MLL) fusion proteins are potent inducers of leukemia, but how these proteins generate aberrant gene expression programs is poorly understood. Here we show that the MLL-AF4 fusion protein occupies developmental regulatory genes important for hematopoietic stem cell identity and self-renewal in human leukemia cells. These MLL-AF4-bound regions have grossly altered chromatin structure, with histone modifications catalyzed by trithorax group proteins and DOT1 extending across large domains. Our results define direct targets of the MLL fusion protein, reveal the global role of epigenetic misregulation in leukemia, and identify new targets for therapeutic intervention in cancer.**

Supplemental material is available at <http://www.genesdev.org>.

Received September 16, 2008; revised version accepted November 4, 2008.

Chromosomal translocations involving the mixed-lineage leukemia gene (*MLL*) are a frequent occurrence in human acute leukemias of both children and adults (Eguchi et al. 2005). In over half of all infant acute leukemias, the *MLL* protein fuses to one of >50 identified partner genes, resulting in a *MLL* fusion protein that acts as a potent oncogene (Krivtsov and Armstrong 2007). While extensive gene expression signatures have been determined for primary human leukemia samples (Armstrong et al. 2002; Yeoh et al. 2002; Ferrando et al. 2003; Ross et al. 2003; Rozovskaia et al. 2003; Haeflrich et al. 2005), the direct genomic targets of *MLL* fusion proteins remain unknown. This information is essential to deter-

mine how *MLL* fusion proteins impose oncogenic transcriptional programs and to identify targets for therapeutic intervention in human disease.

Distinct chromatin-modifying complexes and histone modifications are associated with distinct phases of transcription (Li et al. 2007). The trithorax group proteins, including *MLL*, catalyze histone H3-Lys-4 trimethyl (H3K4me3) modifications at the start sites of transcriptionally engaged genes (Ruthenburg et al. 2007). These H3K4me3-modified regions are largely constrained to the transcription start site regions of genes that are transcriptionally initiated, but not necessarily fully transcribed (Bernstein et al. 2006; Barski et al. 2007; Guenther et al. 2007). As a gene becomes fully transcribed, elongating RNA Polymerase II (Pol II) molecules proceed through gene coding regions along with associated elongation factors including DOT1, which catalyzes dimethylation of histone H3-Lys-79 (H3K79me2) (Li et al. 2007). Physical interactions between the most common *MLL* partner proteins and transcriptional elongation components suggest that defects in H3K4 and H3K79 methylation might be a key factor in *MLL* leukemogenesis (Erfurth et al. 2004; Milne et al. 2005b; Okada et al. 2005; Zeisig et al. 2005; Bitoun et al. 2007; Mueller et al. 2007), but the mechanism and extent of H3K79 methylation targeting throughout the genome is poorly understood in human cancer cells.

In order to define the portion of gene regulatory circuitry that is controlled directly by *MLL* fusion proteins in human leukemia, we determined the binding patterns of an *MLL* fusion protein and chromatin modifications across the entire human genome. We performed this mapping in leukemic cells harboring the *MLL*-AF4 fusion gene, because this rearrangement is the most common among *MLL* fusions and is associated with an extremely poor prognosis in infants and adults (Eguchi et al. 2005). Our results reveal that the *MLL*-AF4 oncogene produces gross defects in chromatin structure at a newly defined set of hematopoietic stem cell genes.

## Results and Discussion

### Identification of *MLL*-AF4-occupied regions of the genome in human leukemia cells

We used chromatin immunoprecipitation (ChIP) coupled to massively parallel sequencing (ChIP-seq) to determine how the *MLL*-AF4 fusion protein was distributed across the entire genome in human leukemia cells. This was performed using two acute lymphoblastic leukemia (ALL) cell lines (Fig. 1). The SEM cell line, which was derived from precursor B-cell ALL patient blast cells (Greil et al. 1994) harbors a t(4;11) chromosomal translocation and expresses *MLL*-AF4 fusion protein, endogenous *MLL* and endogenous AF4. The REH precursor B-cell ALL patient-derived cell line (Rosenfeld et al. 1977), which expresses only wild-type *MLL* and wild-type AF4, served as a control to identify regions bound by normal *MLL* and AF4, but not by the *MLL*-AF4 fusion protein. Because REH cells were derived from patients with B-cell ALL, these cells also served to control for general effects of B-cell-type leukemia.


We mapped protein-DNA interactions for the *MLL*-AF4 protein in SEM cells using antibodies to the C ter-

[Keywords: Chromatin; epigenetic; genome-wide; leukemia; stem cell; transcription]

<sup>5</sup>Corresponding author.

E-MAIL [young@wi.mit.edu](mailto:young@wi.mit.edu); FAX (617) 258-0367.

Article is online at <http://www.genesdev.org/cgi/doi/10.1101/gad.1741408>.




**Figure 1.** Mapping MLL-AF4 fusion protein-binding sites in human leukemia cells. (A) Schematic diagram of strategy for mapping MLL-AF4 fusion protein-binding sites. SEM precursor B acute leukemia cells express the MLL-AF4 fusion protein. REH precursor B acute leukemia cells express only endogenous AF4 and MLL1. The N terminus of MLL (blue) is recognized by ChIP antibody anti-MLL-N (blue) and immunoprecipitates both wild-type MLL and MLL-AF4 fusion protein in SEM cells. The C terminus of AF4 (red) is recognized by ChIP antibody anti-AF4-C and immunoprecipitates both wild-type AF4 and MLL-AF4 fusion protein in SEM cells. Wild-type AF4 and MLL-N are immunoprecipitated by anti-AF4-C and anti-MLL-N, respectively. (B) Binding of AF4 (red) and MLL-N (black) in SEM cells (top panels) and REH cells (bottom panels) as determined by ChIP-seq. Binding profiles are shown across an 800-kb portion of the genome surrounding the PROM1 gene (gene models shown in black below graph; a black arrow indicates transcription start sites). MLL-AF4 fusion protein binding is indicated by a red bar.

minus of AF4 (anti-AF4<sup>C</sup>) and the N terminus of MLL (anti-MLL<sup>N</sup>) (Fig. 1A). Since REH cells do not express MLL-AF4 fusion protein, the sites of MLL and AF4 occupancy that were highly similar in SEM and REH cells are most likely due to binding of normal MLL or AF4 proteins. We expected large numbers of regions in both SEM and REH to be occupied by either AF4 or MLL, since these proteins are widely involved in activated transcription by Pol II (Guenther et al. 2005; Milne et al. 2005a; Bitoun et al. 2007). Indeed, we found that MLL or AF4 localized near the transcription start sites of several

thousand genes in both SEM and REH cells (Supplemental Table S1). These bound regions tended to be relatively small, extended 1–3 kb into the 5' end of the transcribed portion of the gene (Supplemental Fig. S1), and were associated with transcriptionally active genes (>80% of transcripts called present by array-based methods in each cell type) (Supplemental Table S2). These data identified the set of genes occupied by MLL and/or AF4 in precursor B ALL cells and confirmed that these genes tend to be transcriptionally active.

We next examined the genome for evidence of binding




**Figure 2.** MLL-AF4 target genes are enriched for early developmental regulators. (A) Signals for AF4 (red) and MLL-N (black) reflecting binding of the presumptive MLL-AF4 fusion protein in SEM cells as determined by ChIP-seq. Binding profiles are shown across a 15- to 150-kb portion of the genome surrounding the *HOXA9*, *TWIST1*, *HMG2*, and *RUNX2* genes (gene models shown in black below graph; a black arrow indicates transcription start sites). Size of MLL-AF4-enriched region is indicated by top brackets. A detailed description of data analysis methods is provided in the Supplemental Material. (B) Composite AF4-C terminus (red) and MLL-N terminus (black) binding profiles for all MLL-AF4 target genes. The start site and direction of transcription of the average gene are indicated by an arrow. (C) ChIP-seq density heat map of AF4-C terminus (red) and MLL-N terminus (black) for all MLL-AF4 target genes. The genomic region from -5kb to +10kb relative to the transcription start site of each gene is shown. Gene order is determined by highest average MLL/AF4 read density from top to bottom. The start site and direction of transcription of the genes are indicated by an arrow. (D) Selected results of GSEA (<http://www.broad.mit.edu/gsea>) of MLL-AF4 target genes.

3405



Baylin 2007). Among these modifications is the methylation of histone H3 at Lys-79 (H3K79me2), which occurs at the 5' coding regions of genes that are experiencing productive transcriptional elongation (Steger et al. 2008), and is a critical checkpoint in transcriptional control (Peterlin and Price 2006; Saunders et al. 2006). Many common MLL partner proteins have been shown to interact with transcriptional elongation components, suggesting that H3K79 methylation might be a key factor in MLL leukemogenesis (Erfurth et al. 2004; Milne et al. 2005b; Okada et al. 2005; Zeisig et al. 2005; Bitoun et al. 2007; Mueller et al. 2007), but the mechanism and extent of H3K79me2 targeting throughout the genome is poorly understood in human cancer cells.




**Figure 4.** Mistargeting of chromatin modifications occur as epigenetic lesions at MLL-AF4 target regions. (A) Binding of H3K79me2 (green) and H3K4me3 (blue) in SEM cells as determined by ChIP-seq. Binding profiles are shown across an 800-kb portion of the genome surrounding the *PROM1* gene. Gene models shown in black below the graph; a black arrow indicates transcription start sites. MLL-AF4 fusion protein-binding regions are indicated by red bars. (B) Binding of H3K79me2 ChIPs (green) and H3K4me3 ChIPs (blue) in SEM cells as determined by ChIP-seq. Binding profiles are shown across 15- to 150-kb portions of the genome surrounding the *HOXA9*, *TWIST1*, *HMGA2*, and *RUNX2* genes. Gene models are shown in black (below graph). A black arrow indicates transcription start sites. MLL-AF4 fusion protein binding is indicated by a red bar. (C) H3K79me2 (green) and H3K4me3 (blue) binding profiles for all MLL-AF4 target genes in SEM cells. Genes are ordered as in Figure 2C. (D) Composite H3K79me2 ChIP enrichments (green) and H3K4me3 ChIP enrichments (blue) for all genes (left) and all MLL-AF4 target genes (right). The start site and direction of transcription of the average gene are indicated by an arrow.

We used ChIP-seq to determine how the H3K79me2 chromatin modification was distributed across the genome in SEM cells and to ascertain whether this modification was associated with all of the MLL-AF4 target regions (Fig. 4). We found ~8000 regions of H3K79me2 enrichment in MLL-AF4 leukemia cells or control cells, with the vast majority (95%) mapping to known transcripts (Supplemental Tables S14, S19). As expected, most genes (~95%) marked by H3K79me2 in MLL-AF4 leukemia and in control cells produced transcripts that were detectable by microarray-based methods (Supplemental Table S2), with peak enrichment occurring downstream from the transcription start sites (Fig. 4D; Supplemental Fig. S5). We next extracted the set of H3K79me2-enriched regions in SEM cells and compared them with genomic regions enriched for MLL-AF4 fusion protein. This analysis revealed that ~98% of MLL-AF4 targets were enriched for the H3K79me2 elongation mark. Strikingly, most of these H3K79me2 enrichments formed abnormal domains spanning 5–100 kb extending upstream of and/or downstream from target gene transcriptional start sites in a highly similar pattern to the MLL-AF4 fusion protein (Fig. 4A,B). In fact, there was a 92% overlap between the MLL-AF4 target regions and H3K79me2-enriched regions at the DNA base-pair level. Based on the induction of these chromatin modifications at MLL-AF4 target regions, we speculate that the MLL-AF4 fusion protein is directly involved in establishing these aberrant chromatin domains in MLL-linked cancer cells.

Since the elongation-linked H3K79me2 modification forms across large domains with MLL-AF4 (Fig. 4A–C), we asked whether regulators of the elongation checkpoint behaved similarly. Indeed, elongation proteins were present across aberrant H3K79me2 domains as determined by ChIP-chip and coimmunoprecipitation experiments. The ENL elongation factor (eleven-nineteen leukemia) bound across aberrant H3K79me2 domains at the *HOXA* and *MEIS1* loci (Supplemental Fig. S6; data not shown) and MLL-AF4 associated with ENL and the pTEFb elongation factor in SEM cells (Supplemental Fig. S7; data not shown). Together, these results indicate that the MLL-AF4 protein, the elongation-associated H3K79me2 modification, and additional elongation factors are mistargeted to regions of the genome encoding key developmental regulators.

We next carried out ChIP-seq experiments in MLL-AF4 leukemia cells using an antibody directed against the histone H3K4me3 modification. This modification is a mark of transcriptional initiation that can be deposited by the MLL complex near the start sites of genes in normal cells (Ruthenburg et al. 2007). Since MLL suffers a monoallelic deletion of its SET-containing the H3K4 methyltransferase domain in MLL-AF4 leukemia, we asked whether MLL-AF4 leukemia cells were able to deposit this histone modification normally across the genome. We found that H3K4me3 modification occurred normally at the start sites of most active genes (~90%) in MLL-AF4 leukemia cells (Supplemental Table S2). Strikingly, at many areas of H3K79me2 mistargeting, H3K4me3 was not only present, but also extended across broad domains of similar size (Fig. 4A–C). These chromatin modifications overlapped highly with the MLL-AF4 fusion and occurred at key leukemia and stem cell-associated genes including *HOXA7*, *HOXA9*, *PROM1*, and *HMGA2* (Fig. 4; Supplemental Table S2). This mislocalization of both H3K79me2 and H3K4me3 modifica-



**Figure 5.** Model of MLL-AF4-mediated gene activation in leukemia. Schematic diagram of transcriptional misregulation in MLL-AF4-induced leukemia. MLL-AF4 associates with elongation proteins (pTEFb, ENL, and DOT1) at sites of epigenetic lesions. Phosphorylated and elongating RNA Pol II shown in yellow. Blue (H3K4me3) and green (H3K79me2) circles represent histone modifications at aberrant chromatin domains. Genes annotated as bound by MLL-AF4 and playing a role in hematopoietic stem cell function are shown to the right and subdivided into functional categories.

tions in MLL-AF4-induced leukemia suggests that the MLL-AF4 fusion protein, a strong transcriptional activator, may be acting at target loci by directly coupling transcriptional initiation and elongation machinery.

#### *A model for MLL-AF4-mediated activation of stem cell-like transcriptional program in leukemia*

We describe here the first genome-wide assessment of chromatin modifications and oncogene binding in the most common form of human MLL-associated acute leukemia. The results indicate that MLL-AF4 fusion proteins selectively occupy regions of the genome that contain developmental regulators important for stem cell identity and self-renewal. Our results also show that abnormal patterns of chromatin modifications, including histone H3K79 and histone H3K4 hypermethylation, occur within large domains of MLL-AF4 occupancy. The observation that MLL-AF4 occupies regions of H3K79 hypermethylation and evidence that MLL-AF4 is physically associated with elongation factors including DOT1 strongly suggests that the fusion protein is responsible for generating large domains of H3K79 hypermethylation. The presence of these aberrant chromatin domains demonstrates that chromatin mistargeting to key regions across the genome is a feature of leukemogenesis.

The presence of aberrant chromatin domains in MLL-linked cancer cells suggests that an abnormal “epigenetic” state exists in these cells. Unlike “genetic lesions” that involve changes in genome sequence in disease cells, “epigenetic lesions” include changes to histone modification states, DNA methylation states, or distribution of chromatin-modifying enzymes (Esteller 2007; Feinberg 2007). By this definition, the aberrant chromatin domains associated with MLL-AF4 binding may thus be considered epigenetic lesions. While it has not been established that these epigenetic lesions cause disease, the H3K79 methylation within these domains likely contributes to oncogenesis since H3K79 methyltransferase activity is required for transformation in MLL-AF10-induced leukemia (Okada et al. 2005). The

mechanism of how MLL-AF4 and epigenetic lesions are targeted in the genome remains a central question in leukemia biology.

The binding of the MLL-AF4 fusion protein to a distinct set of developmental genes is of particular interest. Rather than associating with most cellular genes that are engaged in the act of transcription, the fusion protein imposes a more specialized gene expression program. Components of this program include genes not associated previously with MLL-AF4/H3K79me2 mistargeting in leukemia patient-derived cells. These include developmental transcription factors, chromatin regulators, and signaling proteins that are central to leukemia stem cell identity, hematopoietic stem cell identity, and self-renewal (Fig. 5). Our findings suggest that MLL-AF4 directly activates a partial hematopoietic stem cell-like transcriptional program found in leukemia stem cells (Krivtsov et al. 2006; Barabe et al. 2007) in concert with an underlying gross defect in chromatin structure.

#### **Materials and methods**

A detailed description of all experimental procedures and data analysis methods can be found in the Supplemental Material.

##### *Cells and cell culture*

Human SEM cells with the t(4;11) translocation and REH control cells were purchased from the American Type Culture Collection (ATCC). All cell lines were maintained in RPMI medium 1640 supplemented with 10% FBS. Cells were cross-linked with 1% formaldehyde as described in the Supplemental Material.

##### *ChIP-seq*

ChIP was combined with direct sequencing as described in detail (Supplemental Material). Briefly, DNA from  $1 \times 10^8$  cells was immunoprecipitated with epitope-specific antibody. Amplified DNA was gel purified and prepared for sequencing using Illumina’s Genomic DNA sample kit. Clustering and 26-cycle sequencing were performed using an Illumina Cluster station and 1G analyzer as per the manufacturer’s instructions. ChIP-seq reads were aligned to the human genome and analyzed as described in the Supplemental Material.

##### *Gene expression analysis*

Total RNA was isolated from  $5 \times 10^6$  REH or SEM cells by TRIzol extraction. One microgram of total RNA was labeled according to Affymetrix protocols and hybridized to Affymetrix HG-U133 2.0 plus arrays. The data were analyzed by using Affymetrix Gene Chip Operating Software using default settings. Additional gene expression data from human leukemia patient samples was collated from Ross et al. (2003). Gene expression data and analysis results are provided in the Supplemental Material.

##### *ChIP-seq, ChIP-chip, and microarray gene expression data and analysis*

Complete ChIP-seq, ChIP-chip, and microarray gene expression data, analysis methods, and results are provided in the Supplemental Material section and in the Gene Expression Omnibus (<http://www.ncbi.nlm.nih.gov/geo>) database under accession number GSE13313.

#### **Acknowledgments**

We thank Y. Zonis, T. Lee, A. Marson, F. Camargo, S. Gupta, and J. Love for help and discussions. We thank the National Cell Culture Center (NCCC) for REH and SEM cells. This work was supported by grants from the Israel Cancer Research Fund and the Wolfson Foundation (to E.C.), the US-Israel BSF (to E.C. and C.M.C.), NIH grant CA 128609 (to C.M.C. and T.N.), and NIH grant HG002668 (to R.Y.). L.N.L. was supported by individual NRSA fellowship from NHLBI (grant no. F32HL082448).

#### **References**

Armstrong, S.A., Staunton, J.E., Silverman, L.B., Pieters, R., den Boer, M.L., Minden, M.D., Sallan, S.E., Lander, E.S., Golub, T.R., and Kors-

- meyer, S.J. 2002. MLL translocations specify a distinct gene expression profile that distinguishes a unique leukemia. *Nat. Genet.* **30**: 41–47.
- Barabe, F., Kennedy, J.A., Hope, K.J., and Dick, J.E. 2007. Modeling the initiation and progression of human acute leukemia in mice. *Science* **316**: 600–604.
- Barski, A., Cuddapah, S., Cui, K., Roh, T.Y., Schones, D.E., Wang, Z., Wei, G., Chepelev, I., and Zhao, K. 2007. High-resolution profiling of histone methylations in the human genome. *Cell* **129**: 823–837.
- Bernstein, B.E., Mikkelsen, T.S., Xie, X., Kamal, M., Huebert, D.J., Cuff, J., Fry, B., Meissner, A., Wernig, M., Plath, K., et al. 2006. A bivalent chromatin structure marks key developmental genes in embryonic stem cells. *Cell* **125**: 315–326.
- Bitoun, E., Oliver, P.L., and Davies, K.E. 2007. The mixed-lineage leukemia fusion partner AF4 stimulates RNA polymerase II transcriptional elongation and mediates coordinated chromatin remodeling. *Hum. Mol. Genet.* **16**: 92–106.
- Cloos, P.A., Christensen, J., Agger, K., and Helin, K. 2008. Erasing the methyl mark: Histone demethylases at the center of cellular differentiation and disease. *Genes & Dev.* **22**: 1115–1140.
- Eguchi, M., Eguchi-Ishimae, M., and Greaves, M. 2005. Molecular pathogenesis of MLL-associated leukemias. *Int. J. Hematol.* **82**: 9–20.
- Erfurth, F., Hemenway, C.S., de Erkenez, A.C., and Domer, P.H. 2004. MLL fusion partners AF4 and AF9 interact at subnuclear foci. *Leukemia* **18**: 92–102.
- Esteller, M. 2007. Epigenetic gene silencing in cancer: The DNA hypermethylome. *Hum. Mol. Genet.* **16**: R50–R59. doi: 10.1093/hmg/ddm018.
- Feinberg, A.P. 2007. Phenotypic plasticity and the epigenetics of human disease. *Nature* **447**: 433–440.
- Ferrando, A.A., Armstrong, S.A., Neuber, D.S., Sallan, S.E., Silverman, L.B., Korsmeyer, S.J., and Look, A.T. 2003. Gene expression signatures in MLL-rearranged T-lineage and B-precursor acute leukemias: Dominance of HOX dysregulation. *Blood* **102**: 262–268.
- Greil, J., Gramatzki, M., Burger, R., Marschalek, R., Peltner, M., Trautmann, U., Hansen-Hagge, T.E., Bartram, C.R., Fey, G.H., Stehr, K., et al. 1994. The acute lymphoblastic leukaemia cell line SEM with t(4;11) chromosomal rearrangement is biphenotypic and responsive to interleukin-7. *Br. J. Haematol.* **86**: 275–283.
- Guenther, M.G., Jenner, R.G., Chevalier, B., Nakamura, T., Croce, C.M., Canaani, E., and Young, R.A. 2005. Global and Hox-specific roles for the MLL1 methyltransferase. *Proc. Natl. Acad. Sci.* **102**: 8603–8608.
- Guenther, M.G., Levine, S.S., Boyer, L.A., Jaenisch, R., and Young, R.A. 2007. A chromatin landmark and transcription initiation at most promoters in human cells. *Cell* **130**: 77–88.
- Haferlach, T., Kohlmann, A., Schnittger, S., Dugas, M., Hiddemann, W., Kern, W., and Schoch, C. 2005. Global approach to the diagnosis of leukemia using gene expression profiling. *Blood* **106**: 1189–1198.
- Jones, P.A. and Baylin, S.B. 2007. The epigenomics of cancer. *Cell* **128**: 683–692.
- Krivtsov, A.V. and Armstrong, S.A. 2007. MLL translocations, histone modifications and leukaemia stem-cell development. *Nat. Rev. Cancer* **7**: 823–833.
- Krivtsov, A.V., Twomey, D., Feng, Z., Stubbs, M.C., Wang, Y., Faber, J., Levine, M.E., Wang, J., Hahn, W.C., Gilliland, D.G., et al. 2006. Transformation from committed progenitor to leukaemia stem cell initiated by MLL-AF9. *Nature* **442**: 818–822.
- Li, B., Carey, M., and Workman, J.L. 2007. The role of chromatin during transcription. *Cell* **128**: 707–719.
- Milne, T.A., Dou, Y., Martin, M.E., Brock, H.W., Roeder, R.G., and Hess, J.L. 2005a. MLL associates specifically with a subset of transcriptionally active target genes. *Proc. Natl. Acad. Sci.* **102**: 14765–14770.
- Milne, T.A., Martin, M.E., Brock, H.W., Slany, R.K., and Hess, J.L. 2005b. Leukemogenic MLL fusion proteins bind across a broad region of the Hox a9 locus, promoting transcription and multiple histone modifications. *Cancer Res.* **65**: 11367–11374.
- Mueller, D., Bach, C., Zeisig, D., Garcia-Cuellar, M.P., Monroe, S., Sreekumar, A., Zhou, R., Nesvizhskii, A., Chinnaiyan, A., Hess, J.L., et al. 2007. A role for the MLL fusion partner ENL in transcriptional elongation and chromatin modification. *Blood* **110**: 4445–4454.
- O'Brien, C.A., Pollett, A., Gallinger, S., and Dick, J.E. 2007. A human colon cancer cell capable of initiating tumour growth in immunodeficient mice. *Nature* **445**: 106–110.
- Okada, Y., Feng, Q., Lin, Y., Jiang, Q., Li, Y., Coffield, V.M., Su, L., Xu, G., and Zhang, Y. 2005. hDOT1L links histone methylation to leukemogenesis. *Cell* **121**: 167–178.
- Pajer, P., Pecenkova, V., Karafiat, V., Kralova, J., Horejsi, Z., and Dvorak, M. 2003. The twist gene is a common target of retroviral integration and transcriptional deregulation in experimental nephroblastoma. *Oncogene* **22**: 665–673.
- Peterlin, B.M. and Price, D.H. 2006. Controlling the elongation phase of transcription with P-TEFb. *Mol. Cell* **23**: 297–305.
- Puisieux, A., Valsesia-Wittmann, S., and Ansseau, S. 2006. A twist for survival and cancer progression. *Br. J. Cancer* **94**: 13–17.
- Ramallo-Santos, M., Yoon, S., Matsuzaki, Y., Mulligan, R.C., and Melton, D.A. 2002. “Stemness”: Transcriptional profiling of embryonic and adult stem cells. *Science* **298**: 597–600.
- Rosenfeld, C., Goutner, A., Choquet, C., Venuat, A.M., Kayibanda, B., Pico, J.L., and Greaves, M.F. 1977. Phenotypic characterisation of a unique non-T, non-B acute lymphoblastic leukaemia cell line. *Nature* **267**: 841–843.
- Ross, M.E., Zhou, X., Song, G., Shurtleff, S.A., Girtman, K., Williams, W.K., Liu, H.C., Mahfouz, R., Raimondi, S.C., Lenny, N., et al. 2003. Classification of pediatric acute lymphoblastic leukemia by gene expression profiling. *Blood* **102**: 2951–2959.
- Rozovskaia, T., Feinstein, E., Mor, O., Foa, R., Blechman, J., Nakamura, T., Croce, C.M., Cimino, G., and Canaani, E. 2001. Upregulation of Meis1 and HoxA9 in acute lymphocytic leukemias with the t(4;11) abnormality. *Oncogene* **20**: 874–878.
- Rozovskaia, T., Ravid-Amir, O., Tillib, S., Getz, G., Feinstein, E., Agrawal, H., Nagler, A., Rappaport, E.F., Issaeva, I., Matsuo, Y., et al. 2003. Expression profiles of acute lymphoblastic and myeloblastic leukemias with ALL-1 rearrangements. *Proc. Natl. Acad. Sci.* **100**: 7853–7858.
- Ruthenburg, A.J., Allis, C.D., and Wysocka, J. 2007. Methylation of lysine 4 on histone H3: Intricacy of writing and reading a single epigenetic mark. *Mol. Cell* **25**: 15–30.
- Saunders, A., Core, L.J., and Lis, J.T. 2006. Breaking barriers to transcription elongation. *Nat. Rev. Mol. Cell Biol.* **7**: 557–567.
- Steger, D.J., Lefterova, M.I., Ying, L., Stonestrom, A.J., Schupp, M., Zhuo, D., Vakoc, A.L., Kim, J.E., Chen, J., Lazar, M.A., et al. 2008. DOT1L/KMT4 recruitment and H3K79 methylation are ubiquitously coupled with gene transcription in mammalian cells. *Mol. Cell Biol.* **28**: 2825–2839.
- Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A., Paulovich, A., Pomeroy, S.L., Golub, T.R., Lander, E.S., et al. 2005. Gene set enrichment analysis: A knowledge-based approach for interpreting genome-wide expression profiles. *Proc. Natl. Acad. Sci.* **102**: 15545–15550.
- Toren, A., Bielora, B., Jacob-Hirsch, J., Fisher, T., Kreiser, D., Moran, O., Zeligson, S., Givol, D., Yitzhaky, A., Itskovitz-Eldor, J., et al. 2005. CD133-positive hematopoietic stem cell “stemness” genes contain many genes mutated or abnormally expressed in leukemia. *Stem Cells* **23**: 1142–1153.
- Wagner, W., Ansorge, A., Wirkner, U., Eckstein, V., Schwager, C., Blake, J., Miesala, K., Selig, J., Saffrich, R., Ansorge, W., et al. 2004. Molecular evidence for stem cell function of the slow-dividing fraction among human hematopoietic progenitor cells by genome-wide analysis. *Blood* **104**: 675–686.
- Yang, J., Mani, S.A., Donaher, J.L., Ramaswamy, S., Itzykson, R.A., Come, C., Savagner, P., Gitelman, I., Richardson, A., and Weinberg, R.A. 2004. Twist, a master regulator of morphogenesis, plays an essential role in tumor metastasis. *Cell* **117**: 927–939.
- Yeoh, E.J., Ross, M.E., Shurtleff, S.A., Williams, W.K., Patel, D., Mahfouz, R., Behm, F.G., Raimondi, S.C., Relling, M.V., Patel, A., et al. 2002. Classification, subtype discovery, and prediction of outcome in pediatric acute lymphoblastic leukemia by gene expression profiling. *Cancer Cell* **1**: 133–143.
- Zeisig, D.T., Bittner, C.B., Zeisig, B.B., Garcia-Cuellar, M.P., Hess, J.L., and Slany, R.K. 2005. The eleven-nineteen-leukemia protein ENL connects nuclear MLL fusion partners with chromatin. *Oncogene* **24**: 5525–5532.



## Aberrant chromatin at genes encoding stem cell regulators in human mixed-lineage leukemia

Matthew G. Guenther, Lee N. Lawton, Tatiana Rozovskaia, et al.

*Genes Dev.* 2008, **22**:

Access the most recent version at doi:[10.1101/gad.1741408](https://doi.org/10.1101/gad.1741408)

---

### Supplemental Material

<http://genesdev.cshlp.org/content/suppl/2008/12/15/22.24.3403.DC1>

### References

This article cites 41 articles, 14 of which can be accessed free at:  
<http://genesdev.cshlp.org/content/22/24/3403.full.html#ref-list-1>

### License

### Email Alerting Service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).

---

Use CRISPRmod for targeted modulation of endogenous gene expression to validate siRNA data

horizon  
a PerkinElmer company