The SESA network links duplication of the yeast centrosome with the protein translation machinery

Bengü Sezen, Matthias Seedorf, and Elmar Schiebel¹

Zentrum für Molekulare Biologie der Universität Heidelberg (ZMBH), 69120 Heidelberg, Germany

The yeast spindle pole body (SPB), the functional equivalent of mammalian centrosome, duplicates in G1/S phase of the cell cycle and then becomes inserted into the nuclear envelope. Here we describe a link between SPB duplication and targeted translation control. When insertion of the newly formed SPB into the nuclear envelope fails, the SESA network comprising the GYF domain protein Smy2, the translation inhibitor Eap1, the mRNA-binding protein Scp160 and the Asc1 protein, specifically inhibits initiation of translation of *POM34* mRNA that encodes an integral membrane protein of the nuclear pore complex, while having no impact on other mRNAs. In response to SESA, *POM34* mRNA accumulates in the cytoplasm and is not targeted to the ER for cotranslational translocation of the protein. Reduced level of Pom34 is sufficient to restore viability of mutants with defects in SPB duplication. We suggest that the SESA network provides a mechanism by which cells can regulate the translation of specific mRNAs. This regulation is used to coordinate competing events in the nuclear envelope.

[**Keywords:** Regulation of translation; spindle pole body; nuclear pore complex; Smy2; Eap1; Scp160]

Supplemental material is available at <http://www.genesdev.org>.

Received January 28, 2009; revised version accepted May 6, 2009.

The yeast spindle pole body (SPB) is the functional equivalent of the mammalian centrosome (Jaspersen and Winey 2004; Jaspersen and Stearns 2008). Centrosomes and SPBs duplicate once per cell cycle, organize microtubules, and are associated with cell cycle regulators (Pereira and Schiebel 2001; Stegmeier and Amon 2004). In addition, both are connected with the nuclear envelope. The budding yeast SPB is embedded in the nuclear envelope in a similar way to the nuclear pore complexes (NPCs). This incorporation into the membrane differs from the centrosome of higher eukaryotes that is anchored to the nuclear envelope via a proteinaceous link (Jaspersen and Winey 2004; Tzur et al. 2006).

The yeast SPB duplicates in G1/S phase of the cell cycle by a template-based mechanism (Adams and Kilmartin 1999; Jaspersen and Winey 2004). An intermediate in this duplication process, the duplication plaque, develops on the cytoplasmic side of the nuclear envelope as an extension of the pre-existing SPB. The duplication plaque then becomes inserted into the nuclear envelope as a consequence of the action of the essential *NDC1*, *MPS2*, *BBP1*, and *NBP1* genes (Winey et al. 1991, 1993; Adams and Kilmartin 1999; Schramm et al. 2000; Araki et al. 2006). A defect in these genes leads to a failure in SPB duplication such that the insertion of the duplication

plaque into the nuclear envelope is either fully or partially compromised. The integral membrane protein Mps2 forms a tight complex with the cytoplasmic Bbp1. For SPB duplication the Mps2–Bbp1 complex cooperates with the Nbp1 protein, and the highly conserved integral membrane protein Ndc1 (Winey et al. 1993; Adams and Kilmartin 1999; Schramm et al. 2000; Araki et al. 2006; Mansfeld et al. 2006).

Ndc1 has an additional role in NPC biogenesis (Lau et al. 2004; Madrid et al. 2006; Mansfeld et al. 2006; Stavru et al. 2006). At the NPC, Ndc1 interacts with two other integral membrane proteins, Pom34 and Pom152 (Alber et al. 2007a,b; Onischenko et al. 2009). Ndc1, Pom34, and Pom152 have redundant functions in the biogenesis of NPCs (Madrid et al. 2006).

Curiously, *NDC1* shows genetic interactions with *EAP1* (Chial et al. 2000). In yeast, Eap1 and Caf20 are the two eIF4E-binding proteins that prevent formation of the eIF4E–eIF4G complex (Gingras et al. 1999), which is crucial for the recruitment of the 5' end of mRNAs to the 40S ribosomal subunit during initiation of translation (Gingras et al. 1999). Eap1 and Caf20 inhibit general initiation of translation in response to stress conditions such as cadmium and diamides in the growth medium or the occurrence of membrane stress (Deloche et al. 2004; Mascarenhas et al. 2008).

SMY2 was discovered as a high dosage suppressor of the conditional lethal *myo2-66* mutation (Lillie and Brown

¹Corresponding author.

E-MAIL e.schiebel@zmbh.uni-heidelberg.de; FAX 49-6221-545892.

Article is online at <http://www.genesdev.org/cgi/doi/10.1101/gad.524209>.

1994). Based on bioinformatics and in vitro studies with the truncated protein, it was concluded that the GYF domain of Smy2 interacts with proteins containing PPG repeats such as Eap1 and pre-mRNA splicing factors (Kofler et al. 2005; Georgiev et al. 2007). In addition, Smy2 has GYF domain-independent functions at COPII vesicles (Higashio et al. 2008). Thus, the molecular role of Smy2 is largely not understood.

We now describe a novel translational initiation control mechanism that is exerted by the Smy2–Eap1–Scp160–Asc1 network of proteins, named SESA, to regulate the initiation of translation of *POM34* mRNA, upon SPB duplication defects. The failure of the SPB to insert into the nuclear envelope triggers SESA to inhibit the translation initiation of *POM34* mRNA. The concomitant reduction in the level of Pom34 protein restores SPB duplication and so ensures survival of cells that encounter difficulties in inserting their SPB into the nuclear envelope.

Results

A genetic link between the *SMY2* gene and the SPB insertion machinery

MPS2 encodes an essential, single membrane-spanning protein that inserts the newly formed SPB into the nuclear envelope (Winey et al. 1991; de la Cruz Munoz-Centeno et al. 1999; Schramm et al. 2000; Araki et al. 2006). Attempts to germinate *mps2Δ* spores from an *mps2Δ/MPS2* parent frequently led to the appearance of slow-growing *mps2Δ* survivors indicating that a suppressor mutation can ensure survival of cells lacking *MPS2* (Fig. 1A; arrow). A genetic screen for the wild-type genes that suppress the lethal growth defect of *mps2Δ* cells identified *SMY2* as a high gene dosage suppressor (Fig. 1B, sector 4). Smy2 is an endoplasmic reticulum (ER)-associated, GYF domain protein whose function remains to be elucidated (Kofler et al. 2005; Higashio et al. 2008).

SMY2 functions through the translation inhibitor EAP1

The cellular function of *SMY2* is obscured by reports on its interaction with proteins involved in translation control, pre-mRNA splicing, and protein secretion (Georgiev et al. 2007; Kofler et al. 2005; Higashio et al. 2008). To unravel the function of *SMY2*, we performed a synthetic lethal screen to reveal genes that become essential in the absence of *SMY2* (Supplemental Fig. S1; Supplemental Material). The screen identified two groups of genes: one set that is involved in the regulation of protein translation, and the other in membrane function (Supplemental Fig. S1C). This data suggested that *SMY2* may act as a functional bridge that connects these two processes.

A link between Smy2 protein and control of protein translation was provided by the report that the GYF domain of Smy2 has the ability to interact with the PPG repeats of the translation inhibitor Eap1 (Kofler et al. 2005). To determine whether *SMY2* functions through *EAP1*, we asked whether *EAP1* becomes essential for viability of *mps2Δ 2μm-SMY2* cells. Loss of *EAP1*

Figure 1. Suppressor screen for the lethal phenotype of *MPS2* deletion. (A) Spore analysis of *MPS2/MPS2* and *mps2Δ/MPS2* cells. a, b, c, and d indicate cells that developed from spores of one complete tetrad. The cells marked by an arrow did not contain the *MPS2* gene. (B) *SMY2* is a multicopy suppressor of the lethality of *mps2Δ* cells. *mps2Δ* pRS316-*MPS2* cells were transformed with the indicated plasmids. Transformants were grown on SC-Leu or 5-FOA plates for 3 d at 23°C.

was lethal for the *mps2Δ 2μm-SMY2* mutant cells (Fig. 2A, sector 3). In addition, the PPG mutant *eap1*[G624A] did not allow growth of *mps2Δ* cells (Supplemental Fig. S2C, row 8) and the GYF domain mutant *smy2*[Y234A] was no longer able to suppress the essential function of *MPS2* (Supplemental Fig. S2A, sector 3). Thus, the interaction between the GYF element of Smy2 and the PPG domain of Eap1 is essential for the bypass of *MPS2*. In contrast, *CAF20* that encodes the second eIF4E inhibitor (Ibrahim et al. 2006), was not required for growth of *mps2Δ 2μm-SMY2* cells (Supplemental Table 1). This data suggests that *SMY2* functions through *EAP1*.

Coimmunoprecipitation experiments were performed to demonstrate that Smy2 interacts with Eap1 in vivo (Fig. 2B). The Eap1–Smy2 interaction was equally observed in wild-type and *mps2Δ 2μm-SMY2* cells (Fig. 2B), suggesting that the *mps2Δ* defect did not induce Smy2–Eap1 interaction. Moreover, the GYF domain of Smy2 was essential for the interaction, as Eap1 did not coimmunoprecipitate with the GYF domain-defective Smy2[Y234A] (Supplemental Fig. S2B). Together, this data suggests that Smy2 interacts in vivo with the translation inhibitor Eap1.

Eap1 allows bypass of *MPS2* function by inhibiting translation factor eIF4E

Eap1 inhibits cap-dependent translation by competing with the initiation factor eIF4G (a subunit of the eIF4F cap-binding complex) for binding to eIF4E (Sonnenberg and Gingras 1998). Because it was recently suggested that Eap1 may execute functions that are distinct from its role

Figure 2. *EAP1*, *SCP160*, and *ASC1* are essential components of the *SMY2* pathway. (A) Dependency of suppression on *EAP1*. *mps2Δ eap1Δ* pRS316-MPS2 cells were transformed with the listed plasmids and tested for growth on SC-Ade and 5-FOA plates. *eap1*[Y109A], *eap1*[L114A], and *eap1*[Y109A,L114A] code for Eap1 proteins with mutations that partially (*eap1*[Y109A], *eap1*[L114A]) or totally (*eap1*[Y109A,L114A]) disrupt binding to eIF4E (Fig. 2D; Ibrahimo et al. 2006). (B) Coimmunoprecipitation of Eap1, Scp160, and Asc1 by Smy2. Extracts from cells expressing *SMY2 EAP1*-9Myc or *SMY2*-6HA *EAP1*-9Myc were immunoprecipitated by anti-HA-coated magnetic beads and analyzed by immunoblotting with the indicated antibodies. (WCE) Whole-cell extract, 10% of the immunoprecipitation input; (IP) immunoprecipitation. (C) Coimmunoprecipitation of Eap1, eIF4E, and eIF4G by Smy2. Extracts from the yeast cells expressing *SMY2 EAP1*-9Myc or *SMY2*-6HA *EAP1*-9Myc were immunoprecipitated by anti-HA-coated magnetic beads and analyzed by immunoblotting with the indicated antibodies. Abbreviations as in B. (D) Coimmunoprecipitation of Smy2 and Eap1 and mutated Eap1 protein that fails to bind to eIF4E. Extracts from yeast cells with the indicated genotypes were immunoprecipitated by anti-HA-coated magnetic beads and analyzed by immunoblotting with anti-HA, anti-Myc, and anti-eIF4E antibodies. Abbreviations are as in B. (E,F) Dependency of suppression on *SCP160* and *ASC1*. *mps2Δ scp160Δ* pRS316-MPS2 (E) or *mps2Δ asc1Δ* pRS316-MPS2 cells (F) were transformed with the indicated plasmids and tested for growth on SC-Leu and 5-FOA at 23°C.

in the inhibition of translation (Chial et al. 2000), we asked whether *EAP1* regulates translation or plays some other function in *mps2Δ* cells. Coimmunoprecipitation experiments showed that eIF4E was associated with Smy2 and Eap1 in both wild-type and *mps2Δ* 2μm-SMY2 cells, whereas eIF4G was excluded from this association (Fig. 2C). The latter was expected as Eap1 competes with eIF4G for the same site in eIF4E.

The interaction between Eap1 and eIF4E is impaired by the *eap1*[Y109A, L114A] mutations (Fig. 2D). This impairment of Eap1/eIF4E association relieves the inhibitory influence of Eap1 upon translation initiation (Ibrahimo et al. 2006). In this *eap1*[Y109A, L114A] background, *SMY2* was no longer able to suppress the lethal phenotype of the *mps2Δ* mutation (Fig. 2A, sector 8), even though the Eap1[Y109A, L114A] protein still associated with Smy2 in coimmunoprecipitation assays (Fig. 2D). Thus, inhibition of translation factor eIF4E by Eap1 binding is crucial for the viability of *mps2Δ* cells.

SCP160, ASC1, and BFR1 become essential in cells lacking MPS2

The proteins that regulate and cooperate with Eap1 in response to membrane defects are largely unknown

(Deloche et al. 2004). With the *mps2Δ* 2μm-SMY2 cells in hand, we were able to test genes for function in conjunction with *EAP1* and *SMY2*. We analyzed about 40 genes with roles in translation control, the UPR pathway (Sidrauski and Walter 1997), and nuclear envelope function (Supplemental Table 1) for their requirement in the viability of *mps2Δ* 2μm-SMY2 cells. This analysis identified the genes coding for the polysome-associated, mRNA-binding protein Scp160 that has been shown to exist in a complex with Eap1 (Mendelsohn et al. 2003), the Scp160-interacting protein Asc1 (essential for ER localization of Scp160) (Baum et al. 2004), and the brefeldin A resistance protein Bfr1 (a component of polyribosome-associated mRNP complexes) (Lang et al. 2001), as being essential in the *mps2Δ* 2μm-SMY2 background (Figs. 2E,F, 3A; Supplemental Table 1).

Coimmunoprecipitation experiments validated this genetic approach. Eap1, Scp160, and Asc1 were found to be in complex with Smy2 (Fig. 2B), and in addition, Scp160, Smy2, and Eap1 cofractionated in part in sucrose gradients (Supplemental Fig. S3). These interactions were not mediated by ribosomes, since the ribosomal proteins Rps3 (40 S ribosomal subunit) and Rpl35 (60 S ribosomal subunit) were not detected in the Smy2 immunoprecipitate (Fig. 2B). Interestingly, coimmunoprecipitation of

Figure 3. *BFR1* is essential for viability of cells lacking *MPS2*. (A) Dependency of suppression on *BFR1*. *mps2Δ bfr1Δ* pRS316-MPS2 cells were transformed with the indicated plasmids and tested for growth on SC-Leu and 5-FOA plates at 23°C. (B) Coimmunoprecipitation of Bbp1 by Bfr1. Extracts from the yeast cells with the indicated genotypes were subjected to immunoprecipitation with anti-HA-coated magnetic beads and analyzed by immunoblotting with the indicated antibodies. Abbreviations as in Figure 2B.

Smy2 and Eap1 relied on the presence of the Scp160 protein but not on Asc1 (Supplemental Fig. S2D) and RNA, as RNaseA treatment only mildly affected the efficiency of coprecipitation (Fig. 4C). Thus, the Smy2–Eap1 interaction was stabilized by Scp160 but was not mediated by Asc1 and RNA.

Bfr1 coimmunoprecipitated with the Scp160 and Asc1 proteins, but failed to coimmunoprecipitate with Smy2 (Fig. 3B; Supplemental Fig. S4A), which establishes a distinct set of interactions from the Smy2–Eap1–Scp160–Asc1 association (Fig. 2B). In addition, we confirmed that the interaction of Bfr1 with Scp160 and Asc1 was RNA-dependent (Supplemental Fig. S4B), as has been reported for the Bfr1–Scp160 interaction (Lang et al. 2001). Thus, Bfr1 concurs with Scp160 and Asc1 in a manner that is distinct from the association of these proteins with Smy2. Based on the common genetic and biochemical interactions, we suggest that the gene products of *SMY2*, *EAP1*, *SCP160*, and *ASC1* closely cooperate in the suppression of *MPS2* deletion. This network of genes was named SESA. Bfr1 may interact through an RNA link with SESA network components.

Interestingly, two hybrid interactions between the SPB component Bbp1, which is in complex with Mps2, and the Bfr1 protein have been reported previously (Xue et al. 1996; Schramm et al. 2000). Coimmunoprecipitation experiments showed complexes containing Bfr1 and Bbp1

(Fig. 3B) and this interaction was independent of RNA (Supplemental Fig. S4B). Thus, Bfr1 and the SPB protein Bbp1 are present in common complexes. This Bbp1–Bfr1 interaction may provide the functional link to couple SPB duplication and the SESA network.

POM34 mRNA is in a complex with Smy2

In previous studies, Eap1 was identified as a global inhibitor of the initiation of protein translation in response to conditions that induce membrane stress. This effect of Eap1 could be measured in sucrose gradients by the increase in monosomes with concomitant decrease in

Figure 4. *POM34* mRNA binds to SESA components. (A,B) The indicated yeast cells were tested for growth on SC-Ura and 5-FOA plates at 23°C. All strains harbored initially the pRS316-MPS2 plasmid. (C) Coimmunoprecipitation of *POM34* mRNA together with Eap1, Scp160, and Asc1 by Smy2. Extracts from yeast cells were incubated with anti-HA-coated magnetic beads, with or without RNaseA treatment, and analyzed by immunoblotting with the indicated antibodies. RNA was isolated from the immunoprecipitates and analyzed by RT-PCR using primers specific to *POM34*, *POM152*, and *NDC1*. In this immunoprecipitation experiment the efficiency of Asc1 coimmunoprecipitation was reduced probably because of the longer incubation time due to RNaseA treatment (cf. Figs. 2 and 4C). Abbreviations are as in Figure 2B.

polysomes upon stress conditions (Deloche et al. 2004). Similar sucrose gradient analysis of the ratio between isolated polysomes and monosomes in wild-type and *mps2Δ* 2 μ m-SMY2 cells excluded the possibility of a global down-regulation of protein translation by Eap1 in *mps2Δ* 2 μ m-SMY2 cells (Supplemental Figs. S5 [cytoplasmic fraction], S6 [membrane-bound fraction]).

We therefore tested whether the interaction of the mRNA-binding protein Scp160 with Smy2 and Eap1 could restrict translation inhibition to a specific subset of mRNAs, which may bind to SESA through the known mRNA-binding protein Scp160 (Frey et al. 2001). Attempts to identify mRNAs that were clearly regulated by Scp160 on the level of translation were unsuccessful. We thus turned to a genetic approach to identify mRNA candidates that are bound to Scp160.

We reasoned that both translation inhibition of an mRNA by the SESA network and loss of the targeted gene could suppress the essential function of *MPS2*. A candidate approach was therefore adopted to screen for genes whose inactivation would suppress the lethality of *mps2Δ* cells. It has been previously reported that the SPB duplication defect of *ndc1-1* cells is suppressed by the loss of the *POM152* gene that encodes an integral membrane protein of the NPC (Chial et al. 1998). Thus, our suppression analysis was initially focused on NPC components. *mps2Δ* lethality was suppressed by deletion of *POM34*, *POM152*, the double deletion of *MLP1* and *MLP2*, and conditional lethal *NDC1* alleles (Fig. 4A). Ndc1 is an integral membrane protein that has overlapping functions with Pom152 and Pom34 in NPC biogenesis (Chial et al. 1998; Tcheperegine et al. 1999; Madrid et al. 2006). The myosin like Mlp1 and Mlp2 are associated with the nuclear side of the NPC and have diverse roles in the retention of nonspliced mRNAs, nuclear transport, and SPB duplication (Strambio-de-Castillia et al. 1999; Galy et al. 2004; Niepel et al. 2005). The *mps2Δ* suppression phenomenon was restricted to these NPC components. Deletion of other nucleoporins (e.g., *NUP133*, *NUP170*, *NUP188*) or other integral membrane protein components of the nuclear envelope (e.g., *UIP3*) did not overcome the lethality of *mps2Δ* (Fig. 4B; see Supplemental Table I for the full list of genes tested).

As outlined above, the SESA network may bind and down-regulate the translation of target mRNAs to suppress the essential requirement of *MPS2*. To test this notion, we immunoprecipitated Smy2 and then isolated the mRNAs that were precipitated in a complex with Smy2. RT-PCRs performed on these RNA samples revealed that *POM34* mRNA associated with the SESA proteins, whereas *POM152* and *NDC1* mRNAs did not (Fig. 4C). Coimmunoprecipitation experiments performed in conjunction with UV cross-linking did not change these outcomes (data not shown). Other mRNAs tested and also found not to associate with Smy2 included *MLP1*, *MLP2*, *SEC61*, *ADH1*, and *SIC1* (data not shown). We further established that the association of *POM34* mRNA with Smy2 and Eap1 was dependent on the presence of Scp160 (Supplemental Fig. S7A,B). In contrast, disruption of *SMY2* or *EAP1* did not affect the

association of *POM34* mRNA with Scp160 (Supplemental Fig. S7C). Taken together, these data indicate that *POM34* mRNA associates with the SESA network of proteins via an interaction with Scp160 protein.

SESA inhibits translation of *POM34* mRNA

The presence of the translation initiation inhibitor Eap1 in the SESA network and the essential requirement of the Eap1-eIF4E interaction for *mps2Δ* suppression (Fig. 2A,D), prompted us to test whether SESA activity controls the translation of *POM34* mRNA. A prediction of this hypothesis is that Pom34 protein levels would be lower in *mps2Δ* 2 μ m-SMY2 cells than in wild-type cells even though there would be no impact on *POM34* mRNA levels. To address this point, we first quantified the protein levels of NPC components in *mps2Δ* 2 μ m-SMY2 and wild-type cells. Our analyses showed that Ndc1 levels were the same in both cell types, whereas Pom152 showed a modest decrease in *mps2Δ* 2 μ m-SMY2 cells (Fig. 5A). Mlp1, Mlp2, Nup133, Nup170, and

Figure 5. Strongly reduced amount of Pom34 in *mps2Δ* 2 μ m-SMY2 cells. (A) Pom34 protein level is reduced in *mps2Δ* 2 μ m-SMY2 cells. Total cell extracts from yeast strains expressing *NDC1*-6HA, *POM152*-6HA, or *POM34*-6HA were analyzed by immunoblotting using anti-HA antibodies. Anti-Tub2 antibodies were used as loading control. The graphs underneath the immunoblots show the quantification of three independent experiments, normalized for the wild-type protein levels. Bars are standard deviations around the mean value. (B) *POM34* mRNA levels are similar in wild-type and *mps2Δ* 2 μ m-SMY2 cells. Total RNA extracts from wild-type (WT) and *mps2Δ* 2 μ m-SMY2 cells were analyzed by quantitative RT-PCR using primers specific to *POM34* mRNA. (C) *mps2Δ* 2 μ m-SMY2 cells do not show mRNA export defect. Wild-type and *mps2Δ* 2 μ m-SMY2 cells were fractionated into nuclear and cytoplasmic fractions and analyzed by immunoblotting with anti-Pgl1 and anti-Nop1 antibodies. RNA was isolated from the nuclear and cytoplasmic fractions and analyzed by RT-PCR using primers specific to *POM34*.

Nup188 levels were also same in wild-type and *mps2Δ* 2 μ m-SMY2 cells (data not shown). In contrast, the level of Pom34 protein was significantly lower in *mps2Δ* 2 μ m-SMY2 cells than in wild type controls (Fig. 5A) even though the stability of the Pom34 protein assessed by the cycloheximide treatment was similar in both cell types (Supplemental Fig. S8). The modest decrease of Pom152 in *mps2Δ* 2 μ m-SMY2 cells may arise from the strongly reduced level of Pom34 that interacts with Pom152 (Alber et al. 2007a,b).

The data described above suggest that Pom34 levels are decreased in *mps2Δ* 2 μ m-SMY2 cells either because of a reduction in the efficiency with which *POM34* mRNA is translated, or because nuclear export of this mRNA is impaired or, finally, that the stability of *POM34* mRNA is reduced. Quantitative RT-PCR established that total levels of *POM34* mRNA were not significantly affected in *mps2Δ* 2 μ m-SMY2 cells (Fig. 5B). Thus, we concluded that mRNA stability is probably not altered in *mps2Δ* 2 μ m-SMY2 cells. Furthermore, a comparison of the levels of *POM34* mRNA in the cytoplasm of *mps2Δ* 2 μ m-SMY2 and wild-type cells revealed no differences (Fig. 5C), indicating that the export of *POM34* mRNA was unperturbed by the *mps2Δ* 2 μ m-SMY2 genotype. We conclude that *POM34* is likely regulated on the level of translation.

Membrane localization of *POM34* mRNA requires initiation of translation

POM34 encodes an integral protein of the nuclear envelope (Miao et al. 2006). Its translation should therefore occur at the ER (Rapoport 1990). In order to pinpoint the step modulated by SESA control more precisely, we monitored the partitioning of *POM34* mRNA between cytosolic and membrane-bound fractions of yeast cell extracts alongside an assessment of the abundance of Scp160, Sec61, and Pgk1 proteins and ribosomal subunits.

In both wild-type and *mps2Δ* 2 μ m-SMY2 cells, cytoplasmic Pgk1 was detected in the soluble cytoplasmic fraction and the integral membrane protein Sec61 (Deshaies et al. 1991) was found almost exclusively in the membrane fraction (Fig. 6A). Scp160 was enriched in the membrane-bound fraction, whereas the ribosomal subunit Rpl35 was more abundant in the cytoplasm. The latter distributions were as expected, since most ribosomes are present in cytoplasm while Scp160 is enriched with ribosomes at the ER (Supplemental Fig. S9A; Frey et al. 2001). Together, the cellular distributions of Scp160, Sec61, Pgk1, and Rpl35 proteins demonstrate that the fractionation into membrane and cytoplasmic fractions was successful.

According to RT-PCR data, the mRNAs of *SEC61* (Fig. 6A), *NDC1*, and *POM152* (Supplemental Fig. S6), which all encode integral membrane proteins, were slightly enriched in the membrane-bound fraction in both wild-type and *mps2Δ* 2 μ m-SMY2 cells (Fig. 6A; Supplemental Fig. S6). In wild-type cells *POM34* mRNA distribution between the membrane fraction and the cytoplasm was similar to *SEC61* mRNA (Fig. 6A). Strikingly, however, in

Figure 6. *POM34* mRNA accumulates in the cytoplasm in *mps2Δ* 2 μ m-SMY2 cells. (A) Total cell extracts from wild-type and *mps2Δ* 2 μ m-SMY2 cells were fractionated into cytosolic and membrane-bound fractions. Fractions were analyzed by immunoblotting using the indicated antibodies. The lower molecular weight band (asterisk) is a cytoplasmic degradation product of Scp160 (Frey et al. 2001). *POM34* and *SEC61* mRNA levels were determined by quantitative RT-PCR from three independent experiments. The graph shows the ratio of cytosolic *POM34* mRNA levels to the membrane-bound *POM34* mRNA levels. The bars indicate standard deviation of the results from the mean value. (C) Cytosolic fraction; (M) membrane-bound fraction. (B) Subcellular localization of *POM34* mRNA in the translation initiation mutant *cdc33-1*. Cells were grown in YPAD at 23°C and were shifted for 2 h to 37°C. The cellular fractionation and analysis was performed as in A.

mps2Δ 2 μ m-SMY2 cells *POM34* mRNA was mostly found in the cytoplasm, and only very little in the membrane fraction (Fig. 6A). Thus, the cellular localization of *POM34* mRNA is significantly changed in *mps2Δ* 2 μ m-SMY2 cells.

Mislocalization of *POM34* mRNA may arise from an inhibition of initiation of protein translation, which is required to target the mRNA in association with the nascent Pom34 protein to the ER membrane. If correct, a general inhibition of translation should result in the same cellular mislocalization of *POM34* mRNA. To test this prediction, we analyzed *POM34* mRNA localization in the conditional lethal translation initiation mutant *cdc33-1*, in which the function of yeast eIF4E is impaired (Brenner et al. 1988). *POM34* mRNA had a predominately cytoplasmic distribution in *cdc33-1* cells incubated at the restrictive temperature (Fig. 6B). The extent of the mislocalization of *POM34* mRNA in *cdc33-1* cells was only slightly higher than in *mps2Δ* 2 μ m-SMY2 cells (Fig. 6A). This establishes that inhibition of translation initiation is indeed sufficient to mislocalize *POM34* mRNA to the cytoplasm.

We next asked whether translation elongation was also affected in *mps2Δ* 2 μ m-SMY2 cells. Separating the translating membrane-bound ribosomal subunits and polyosomes via a sucrose gradient fractionation was used as

experimental approach (Supplemental Fig. S6). Although less abundant in the membrane fraction, we observed no major difference in the relative *POM34* mRNA distribution between membrane-bound polysomes and monosomes in *mps2Δ* 2 μ m-*SMY2* cells with respect to wild-type cells (Supplemental Fig. S6C). Thus, our data do not suggest a defect in translation elongation for *POM34* mRNA in *mps2Δ* 2 μ m-*SMY2* cells. We also analyzed the cytoplasmic fraction of *POM34* mRNA in a sucrose gradient (Supplemental Fig. S5). This analysis showed that the cytoplasmic *POM34* mRNA, although more abundant in *mps2Δ* 2 μ m-*SMY2* cells than in wild-type cells (Fig. 6A), was predominately associated with the 40S, 60S, and 80S fractions of ribosomes in both cell types (Supplemental Fig. S5). A polysomal cytoplasmic fraction was not observed, which is consistent with the notion that translation elongation of *POM34* mRNA occurred at the ER. Thus, the main difference between wild type and *mps2Δ* 2 μ m-*SMY2* cells is the strongly reduced translation initiation efficiency of *POM34* mRNA in *mps2Δ* 2 μ m-*SMY2* cells. This defect then leads to the accumulation of *POM34* mRNA in the cytoplasm.

SPB duplication defects regulate Pom34 via the SESA network

Our data indicate that the SESA network inhibits translation of the *POM34* mRNA in *mps2Δ* 2 μ m-*SMY2* cells. However, overexpression of *SMY2* in otherwise wild-type cells was insufficient to down-regulate *Pom34* (Supplemental Fig. S9B), suggesting that a defect in *mps2Δ* cells induces the regulation of *POM34* mRNA. The implication is that defects in SPB duplication may activate SESA. In this respect it is important to note that the SESA

network component *EAP1* is essential for viability of the SPB duplication-defective *ndc1-1* cells that have normal NPCs (Chial et al. 2000). In addition, as was the case for *mps2Δ* (Fig. 1), the lethal growth defect of *ndc1-1* cells at 14°C was suppressed by the deletion of *POM34* (Fig. 7A), or high gene dosage of *SMY2* (Fig. 7B). Furthermore, *Pom34* levels were reduced in *ndc1-1* 2 μ m-*SMY2* cells (Fig. 7C). These similarities between the consequences of mutations in *NDC1* and *MPS2* support the notion that defects in SPB duplication down-regulate *POM34* mRNA translation. Consistently, growth of the SPB duplication-defective *bbp1-1*, *mps2-42*, and *ndc1-4* cells at the restrictive temperature was rescued by *pom34Δ* (Fig. 7A; Supplemental Fig. S10A). Although, high gene copy 2 μ m-*SMY2* did not suppress the growth defect of these cells (data not shown), this is explained by the failure of *SMY2* suppression at elevated temperatures (Fig. 7D).

To further demonstrate that SESA becomes important for survival of mutants defective in SPB duplication, we analyzed the consequence of SESA inactivation in *bbp1-1* and *ndc1-4* cells. Deletion of SESA network components enhanced the growth defect of *bbp1-1* or *ndc1-4* cells grown at elevated growth temperatures (Fig. 7E; Supplemental Fig. S10B), suggesting that SESA network is important under conditions where *SMY2* is not overexpressed.

Down-regulation of POM34 restores SPB duplication of mps2 cells

SPB duplication fails in the absence of *BBP1*, *NDC1*, and *MPS2* function. In these mutant cells the duplication plaque is not inserted into the nuclear envelope. The nuclear side of the SPB with the component *Spc110* is therefore not assembled, whereas the *Spc42* becomes

Figure 7. Genetic interactions between SESA and *POM34* with genes involved in SPB duplication. (A) Deletion of *POM34* suppresses *ndc1-1*, *bbp1-1*, and *mps2-42*. The listed yeast cells were tested for growth on YPAD plates at indicated temperatures. (B) *SMY2* is a multicopy suppressor of the cold-sensitive phenotype of *ndc1-1*. *ndc1-1* cells transformed with pRS315-*NDC1* or pRS425-*SMY2* were tested for growth on YPAD plates at 14°C and 30°C. (C) *Pom34* protein level is reduced in *ndc1-1* 2 μ m-*SMY2* cells. Total cell extracts from yeast strains expressing *POM34*-6HA were analyzed by immunoblotting using anti-HA antibodies. Anti-Tub2 antibodies were used as loading control. The graph shows the quantification of three independent experiments, normalized for the wild-type protein levels. Bars are standard deviations around the mean value. (D) *mps2Δ* 2 μ m-*SMY2* and *mps2Δ pom34Δ* cells are temperature sensitive for growth. The listed yeast cells were tested for growth on YPAD plates at indicated temperatures. (E) Deletion of SESA components enhances *bbp1-1* growth defects. The listed yeast cells were tested for growth on YPAD plates at the indicated temperatures.

incorporated into the defective SPB (37°C: note *mps2-42* cells with only one Spc110-GFP signal but two Spc42-eqFP611 signals) (Fig. 8B; Schramm et al. 2000). However, deletion of *POM34* restored SPB duplication of *mps2-42* cells at 37°C indicating the importance of Pom34 levels in SPB duplication mutants (Fig. 8B; *mps2-42 pom34Δ* cells at 37°C with two colocating Spc110-GFP and Spc42-eqFP611 signals). Furthermore, analysis of *mps2Δ* 2 μ m-*SMY2* and *mps2Δ pom34Δ* cells at 23°C showed that the SPB duplicated in G1/S with similar kinetics as in wild-type cells (data not shown), and that the newly formed SPB carried the markers Ndc1-GFP, Bbp1-GFP, Nbp1-GFP, and Spc110-GFP ($n = 150$ cells) (Supplemental Figs. S11, S12). Thus, deletion or reduced translation of *POM34* suppresses SPB duplication defects.

Discussion

The initiation is the rate-limiting step in mRNA translation. Deregulating initiation by overexpression of the

CAP-binding protein eIF4E leads to malignant transformation and therefore, not surprisingly, eIF4E is elevated in many human cancers (De Benedetti and Rhoads 1990; Lazaris-Karatzas et al. 1990). In addition, TOR signaling and stress situations including membrane defects inhibit global initiation of translation by regulating binding of proteins (4E-BPs) to the initiation factor eIF4E (Cosentino et al. 2000; Deloche et al. 2004; Matsuo et al. 2005; Ibrahim et al. 2006).

In this study, we unraveled an unexpected link between the SPB duplication pathway and regulation of translation initiation of the *POM34* mRNA. In response to SPB duplication defects, the SESA network, comprising of the known mRNA-binding protein Scp160 (Frey et al. 2001; Li et al. 2003, 2004; Baum et al. 2004), the ribosome-associated Asc1 (Baum et al. 2004; Gerbasi et al. 2004), the translation inhibitor Eap1 (Cosentino et al. 2000), and the protein Smy2 (Kofler et al. 2005), was identified as being responsible for the translation control of *POM34* mRNA. We demonstrate that SESA inhibits translation of *POM34* mRNA by binding of the 4E-BP Eap1 to the conserved translation initiation factor eIF4E. By showing that Smy2, Eap1, Scp160, and Asc1 physically and functionally interact, we provide a first understanding of how Eap1 is regulated on the molecular level. This regulation of *POM34* mRNA by SESA is essential to ensure survival of cells with defects in SPB duplication.

SESA binds to *POM34* mRNA

Based on the presence of the eIF4E inhibitor Eap1 in the SESA network, we expected to see a general inhibition of translation initiation in *mps2Δ* cells as is the case in cells exposed to membrane stress conditions (Deloche et al. 2004). Surprisingly, however, in *mps2Δ* 2 μ m-*SMY2* cells translation of proteins was not inhibited on a global scale (Supplemental Figs. S5, S6) indicating that the SESA network regulates only a subset of mRNAs.

The mRNA-binding protein Scp160 could target mRNAs to SESA regulation (Frey et al. 2001; Baum et al. 2004; Li et al. 2004). A microarray analyses of mRNAs released from affinity isolated Scp160-containing complexes identified a limited set of mRNAs that bind to Scp160 (Li et al. 2003). We tested the two most prominent mRNAs identified by Li et al. (2003), the *DHH1* and *YOR338w* transcripts, but did not find an enrichment of these mRNAs in anti-Smy2 immunoprecipitates nor any importance of the genes for survival of *mps2Δ* 2 μ m-*SMY2* cells (B Sezen, unpubl.). Furthermore, a recent study identified mainly mRNAs coding for proteins of the cell wall, plasma membrane, ER and nucleolus in association with Scp160 (Hogan et al. 2008).

To identify mRNAs that are regulated by SESA in the context of *MPS2* function, we turned to a genetic approach, which identified the *POM34* mRNA (Fig. 4) as being associated with SESA components. In addition, Pom34 levels were down-regulated in *mps2Δ* cells that require SESA for viability (Fig. 5A). Together, this strongly supports the notion that *POM34* mRNA is regulated by SESA.

Figure 8. SPBs are inserted into the nuclear envelope in *mps2-42 pom34Δ* cells. (A,B) Analysis of wild-type, *mps2-42*, and *mps2-42 pom34Δ* cells with *SPC110*-GFP *SPC42*-eqFP611 by fluorescence and phase contrast (DIC) microscopy at 23°C (A) and 37°C (B) (2 h). Note that in B Spc110-GFP is only associated with one of the two Spc42-eqFP611-marked SPBs of *mps2-42* cells. This is the typical phenotype of cells with a defect in duplication plaque insertion (Schramm et al. 2000; Jaspersen and Winey 2004). Bars, 5 μ m. (C) Shown is a cartoon of the SPB with the localization of Spc42 and Spc110 relative to the nuclear envelope (NE) (Adams and Kilmartin 1999).

SESA inhibits initiation of translation of *POM34* mRNA

We suggest that SESA inhibits initiation of translation of *POM34* mRNA (Fig. 9, step 4). This model is supported by a number of findings. First, Eap1 blocks the crucial binding of eIF4G to eIF4E (Fig. 2C), which is normally an essential step in translation initiation (Gingras et al. 1999). Second, the Eap1[Y109A, L114A] mutations, which impair the Eap1–eIF4E interaction (Fig. 2D; Ibrahim et al. 2006), also abrogate the function of SESA in *mps2Δ* cells (Fig. 2A) indicating that Eap1–eIF4E binding is essential for the down-regulation of Pom34. Third, although the *POM34* mRNA levels were unchanged in *mps2Δ* cells (Fig. 5B), Pom34 protein was strongly reduced in comparison with wild-type cells (Fig. 5A). A detailed analysis excluded *POM34* mRNA nuclear export defects and altered Pom34 protein stability as factors that could decrease Pom34 protein levels in *mps2Δ* cells (Fig. 5C; Supplemental Fig. S8). However, analysis of the distribution of *POM34* mRNA in *mps2Δ* cells clearly showed a shift toward the cytoplasmic fraction that has not initiated translation (Fig. 6A). A similar cytoplasmic shift of *POM34* mRNA was observed in cells defective in the translation initiation factor eIF4E (Fig. 6B; Brenner et al. 1988), supporting the idea that SESA inhibits initiation of translation. Fourth, conditions that inhibited *POM34* translation did not affect translation of other mRNAs (*NDC1*, *POM152*, *MLP1*, *MLP2*, *SEC61*, *ADH1*, and *SIC1*). This implies that SESA is a specific inhibitor of translation of a subset of mRNAs. Smy2, Scp160, and Asc1 in the SESA network may confer Eap1 translation inhibition specificity toward a subset of mRNAs.

The Scp160 protein with its conserved KH RNA-binding domains (Frey et al. 2001) was deduced as the factor that binds the *POM34* mRNA in the SESA network (Fig. 4C; Supplemental Fig. S7). Scp160 shows genetic and biochemical interactions with the Asc1 protein, which is a core component of the 40S ribosomal subunit and as

such binds Scp160 to ribosomes (Baum et al. 2004; Nilsson et al. 2004). However, also a cytoplasmic pool of Asc1 exists (Brodersen and Nissen 2005) that may direct the SESA–*POM34* mRNA complex to ribosomes. It is important to note that Asc1 functions as G-protein subunit coupled to glucose responsiveness in yeast (Zeller et al. 2007), and that the G protein α subunit Gpa1 transmits a signal through Scp160 (Guo et al. 2003). This raises the exciting possibility of a cross-talk between the SESA network and external stimuli.

Regulation of *POM34* mRNA in response to SPB duplication defects

Mutants of *NDC1* and *MPS2* that failed to insert the SPB duplication plaque into the nuclear envelope showed genetic interactions with SESA components and, in addition, showed mislocalization of *POM34* mRNA and reduced levels of Pom34 in comparison with wild-type cells. These dependencies suggest a mechanism by which SESA becomes active in response to defects in SPB duplication (Figs. 1, 7; Chial et al. 2000; Schramm et al. 2000). This model is consistent with the activation of mammalian 4E-BPs by external stimuli (Gingras et al. 1999) or of yeast Eap1 by membrane defects (Deloche et al. 2004). In addition, elevated gene levels of *SMY2*, may arise due to chromosome missegregation in mutants with SPB duplication defects, which are genetically unstable (Winey et al. 1991, 1993; Schramm et al. 2000; Araki et al. 2006). Increased gene dosage of *SMY2* then likely makes the SESA network more sensitive to defects (Fig. 1).

The SESA network may be linked to the SPB by the Bfr1 protein and the interacting SPB component Bbp1 (Xue et al. 1996). Consistently, *BFR1* was found to be essential in cells lacking *MPS2* and the Bfr1 protein showed co-immunoprecipitation with Bbp1, Scp160, and Asc1 (Fig. 3). However, in contrast to SESA components, RNA mediated the interactions of Bfr1. Binding of Bfr1 to SESA-associated *POM34* mRNA could activate the ability of Eap1 to inhibit the translation of this mRNA.

Why is the NPC component Pom34 important for SPB duplication? Pom34, Ndc1, and Pom152 belong to a group of functionally redundant and interacting integral membrane proteins. They are part of the membrane ring of the NPC and as such important for NPC biogenesis (Fig. 9, step 1; Wozniak et al. 1994; Chial et al. 1998; Madrid et al. 2006; Miao et al. 2006; Alber et al. 2007a,b; Onischenko et al. 2009). A functional link between SPB insertion and NPCs, which are frequently observed near duplicating SPBs, was already proposed (Adams and Kilmartin 1999; Jaspersen and Winey 2004). Curiously, deletion of either *POM34* or *POM152* or mutations in *ndc1* suppress the essential function of *MPS2* (Fig. 4A). Based on this observation we suggest that the Pom34–Pom152–Ndc1 complex inhibits SPB duplication (Fig. 9, step 2). The Pom34–Pom152–Ndc1 complex either directly binds to SPBs and inhibits its duplication or SPBs and NPCs compete for components of a common nuclear envelope insertion machinery. The dual function of Ndc1 in SPB

Figure 9. Model for the function of SESA network. (Step 1) Pom34, Pom152, and Ndc1 form a complex that functions in NPC biogenesis (Chial et al. 1998; Madrid et al. 2006; Alber et al. 2007a,b; Onischenko et al. 2009). (Step 3) Ndc1 has a dual role and together with Mps2, Bbp1, and Nbp1 it also functions in SPB duplication (Winey et al. 1993; Araki et al. 2006). (Step 2) Deletion of *POM34* or *POM152* or mutations in *ndc1* rescue SPB duplication defects (Fig. 4; Chial et al. 1998), suggesting an inhibitory role of the Pom34–Pom152–Ndc1 complex in SPB duplication. In response to SPB duplication defects, SESA down-regulates translation of *POM34* (Figs. 4–7). (Step 4) This in turn rescues the defect and allows for SPB duplication (Fig. 8; Supplemental Fig. S12).

duplication (Fig. 9, step 3) and NPC biogenesis (Fig. 9, step 1) supports the second competition model (Chial et al. 1998; Madrid et al. 2006). Defects in SPB duplication down-regulate expression of *POM34* via SESA (Fig. 9, step 4). Reduced Pom34 levels then relieve the inhibitory function of the Pom34–Pom152–Ndc1 complex. Thus, depending on the conditions, the SESA network may promote either SPB duplication or NPC biogenesis.

This study unraveled a novel mechanism by which cells can regulate the translation of specific mRNAs. In contrast to the general translation inhibition in response to, for example, membrane stress (Deloche et al. 2004), regulation by the SESA pathway enables the cell to modify the proteome in a very specific way at the level of translation. Thus, in respect of specificity SESA regulation is similar to the translation control of mRNAs containing a cytoplasmic polyadenylation element (CPE) by Maskin or 4E-T (4E-BP) in *Xenopus laevis*. In this case the CPE-binding protein (CPEB) acts as the specific RNA-binding protein (Stebbins-Boaz et al. 1999; Minshall et al. 2007). Another such example is the translational repression of *oskar* mRNA in *Drosophila* by Cup (a 4E-BP) where Bruno acts as the mRNA-binding protein (Nakamura et al. 2004; Chekulaeva et al. 2006; for reviews, see Richter and Sonenberg 2005; Sonenberg and Hinnebusch 2009).

Materials and methods

Strain constructions and growth conditions

Gene deletions and epitope tagging of genes at their endogenous loci were performed using PCR-based methods (Janke et al. 2004). The strains and plasmids used in this study are listed in Supplemental Table 2. All yeast strains were derivatives of S228c with the exception of *ndc1-4* and *ndc1-39*, which were derived from W303 and were compared with the corresponding wild type.

Typically cells were grown in yeast extract peptone glucose medium (YPD) at 23°C. For analysis of temperature-sensitive mutants they were shifted for 2 h to 37°C before analysis.

Construction of CDC33, SMY2, and EAP1 mutants

SMY2 and *EAP1* with regulatory and coding regions were cloned into the *LEU2*-based yeast shuttle vector pRS425 and pRS315, respectively (Sikorski and Hieter 1989). Mutations in *SMY2* and *EAP1* were introduced by PCR-directed mutagenesis and confirmed by DNA sequencing. *cdc33-1* with regulatory and coding regions were cloned into the *LEU2*-based yeast integration vector pRS305 (Sikorski and Hieter 1989) after amplification by PCR from the genomic DNA of a *cdc33-1* strain (Brenner et al. 1988) and then inserted into our S288c strain background.

Antibodies and immunoblotting

Yeast extracts were prepared using alkaline lysis and TCA precipitation (Janke et al. 2004). To detect proteins by immunoblotting procedures, blocked membranes (Protein, Schleicher & Schuell) were incubated for 2 h at 20°C or overnight at 4°C with antibodies diluted in blocking buffer (PBS, 0.2% Tween 20, 5% dry milk powder) followed by peroxidase-conjugated secondary antibodies [Sigma] and detection with ECL [Roche

Molecular Biochemicals). Anti-Scp160 (GST-Scp160), anti-Asc1 (NVIRVWQVMTAN-COOH), anti-Rps3 (VALISKRRKLVDACONH2), anti-Rpl35 (CPIRKYAIKV-COOH), anti-Sec61, anti-Clb2 (GST-Clb2¹⁻²⁷¹), anti-Pds1 (GST-Pds1¹⁻¹⁷³), and anti-Tub2 antibodies (yeast β -tubulin, GST-Tub2⁴³⁶⁻⁴⁵⁷) were prepared in rabbits or sheep against purified recombinant proteins or peptides (Frey et al. 2001; Pereira and Schiebel 2003; Baum et al. 2004). Monoclonal mouse anti-Myc (9E10) and anti-HA (12CA5) antibodies were from Roche Molecular Biochemicals. Anti-Pgk1 and anti-Nop1 antibodies were gifts from M. Knop. Anti-eIF4E and anti-eIF4G antibodies were gifts from M. Ashe.

Coimmunoprecipitation

Logarithmically growing cells (3×10^8) were disrupted with glass beads in 300- μ L immunoprecipitation buffer (50 mM triethanolamine, 150 mM KCl, 5 mM EDTA, 5 mM EGTA) containing a protease inhibitor cocktail (Boehringer Mannheim), phenylmethylsulfonyl fluoride (PMSF) and benzamidine. Lysates were incubated with 1% Triton X-100 for 10 min at 4°C. Extracts were cleared by centrifugation (6000 rpm for 10 min at 4°C). After removing an aliquot that served as the input control, the resulting extract was incubated with monoclonal anti-HA antibody, 12CA5, coated magnetic beads (Dyna) for 2 h at 4°C. Beads were washed three times with immunoprecipitation buffer containing 0.1% Triton X-100.

mRNA coimmunoprecipitation

Protein–mRNA coimmunoprecipitations were performed as described (Munchow et al. 1999; Bohl et al. 2000; Long et al. 2000). In brief, 3×10^8 logarithmically growing cells were disrupted with glass beads in 200- μ L breakage buffer BB (50 mM HEPES-KOH at pH 7.3, 50 mM potassium acetate, 2 mM magnesium acetate, 1% Triton X-100) containing a protease inhibitor cocktail (Boehringer Mannheim) and 0.5% BSA. Extracts were cleared by centrifugation (6000 rpm for 10 min). After removing an aliquot that served as input control, the resulting extract was incubated at 4°C with monoclonal anti-HA antibody, 12CA5, coated magnetic beads (Dyna). Beads were washed three times with BB lacking BSA. Pellets were extracted with phenol-chloroform, ethanol precipitated, resuspended in RQ1 DNase buffer, and treated with RQ1 DNase (Promega). The remaining RNA was extracted, precipitated, and resuspended in water. RT-PCR was performed with 1 μ L RNA as a template using the Qiagen RT-PCR kit and the conditions suggested by the manufacturer. The number of amplification cycles was adjusted to avoid reaching a plateau during PCR. For amplification of *POM34*, *POM152*, *SIC1*, and *ASH1* RNAs, we used 25 cycles, for amplification of *NDC1*, *MLP1*, and *MLP2* RNAs we used 24 cycles, whereas 22 cycles were used for amplification of *ADH1* RNA. The primers in Supplemental Table 3 were used for amplification.

RT-quantitative PCR

RNA was extracted from cells grown to logarithmic phase using Qiagen RNeasy kit following the manufacturer's protocol. A total of 4 μ g of RNA was reverse transcribed using 7 μ M oligo-dT and reverse transcriptase for 1 h at 37°C. The product was diluted 1:10 and used in the subsequent quantitative PCR reactions using *POM34* and *SEC61* primers in a Roche LightCycler using SYBR Green. Standard curves for each primer were generated using serial dilutions of yeast genomic DNA. Quantification of cDNA template concentrations was done using the standard curve for each primer.

Acknowledgments

We thank C. Mayer and I. Grummt for their help with quantitative PCR analyses. We thank K. Nasmyth, E. Hurt, and M. Winey for yeast strains; M. Ashe and M. Knop for plasmids and antibodies; P. Walter and K. Weis for plasmids; D. Kirkpatrick for the yeast genomic DNA library; and members of the Schiebel laboratory, I. Hagan, and G. Pereira for comments on the manuscript. We thank C. Schramm for the initial *mps2Δ* 2- μ m screen performed at the Beatson Institute, Glasgow. The GRK1188 and SFB638 supported this work.

References

- Adams IR, Kilmartin JV. 1999. Localization of core spindle pole body (SPB) components during SPB duplication in *Saccharomyces cerevisiae*. *J Cell Biol* **145**: 809–823.
- Alber F, Dokudovskaya S, Veenhoff LM, Zhang W, Kipper J, Devos D, Suprpto A, Karni-Schmidt O, Williams R, Chait BT, et al. 2007a. Determining the architectures of macromolecular assemblies. *Nature* **450**: 683–694.
- Alber F, Dokudovskaya S, Veenhoff LM, Zhang W, Kipper J, Devos D, Suprpto A, Karni-Schmidt O, Williams R, Chait BT, et al. 2007b. The molecular architecture of the nuclear pore complex. *Nature* **450**: 695–701.
- Araki Y, Lau CK, Maekawa H, Jaspersen SL, Giddings JTH, Schiebel E, Winey M. 2006. The *Saccharomyces cerevisiae* spindle pole body (SPB) component Nbp1p is required for SPB membrane insertion and interacts with the integral membrane proteins Ndc1p and Mps2p. *Mol Biol Cell* **17**: 1959–1970.
- Baum S, Bittins M, Frey S, Seedorf M. 2004. Asc1p, a WD40-domain containing adaptor protein, is required for the interaction of the RNA-binding protein Scp160p with polysomes. *Biochem J* **380**: 823–830.
- Bohl F, Kruse C, Frank A, Ferring D, Jansen RP. 2000. She2p, a novel RNA-binding protein tethers *ASH1* mRNA to the Myo4p myosin motor via She3p. *EMBO J* **19**: 5514–5524.
- Brenner C, Nakayama N, Goebel M, Tanaka K, Toh-e A, Matsumoto K. 1988. *CDC33* encodes mRNA cap-binding protein eIF-4E of *Saccharomyces cerevisiae*. *Mol Cell Biol* **8**: 3556–3559.
- Brodersen DE, Nissen P. 2005. The social life of ribosomal proteins. *FEBS J* **272**: 2098–2108.
- Chekulaeva M, Hentze MW, Ephrussi A. 2006. Bruno acts as a dual repressor of oskar translation, promoting mRNA oligomerization and formation of silencing particles. *Cell* **124**: 521–533.
- Chial HJ, Rout MP, Giddings TH, Winey M. 1998. *Saccharomyces cerevisiae* Ndc1p is a shared component of nuclear pore complexes and spindle pole bodies. *J Cell Biol* **143**: 1789–1800.
- Chial HJ, Stemm-Wolf AJ, McBratney S, Winey M. 2000. Yeast Eap1p, an eIF4E-associated protein, has a separate function involving genetic stability. *Curr Biol* **10**: 1519–1522.
- Cosentino GP, Schmelzle T, Haghighat A, Helliwell SB, Hall MN, Sonenberg N. 2000. Eap1p, a novel eukaryotic translation initiation factor 4E-associated protein in *Saccharomyces cerevisiae*. *Mol Cell Biol* **20**: 4604–4613.
- De Benedetti A, Rhoads RE. 1990. Overexpression of eukaryotic protein synthesis initiation factor 4E in HeLa cells results in aberrant growth and morphology. *Proc Natl Acad Sci* **87**: 8212–8216.
- de la Cruz Munoz-Centeno M, McBratney S, Monterrosa A, Byers B, Mann C, Winey M. 1999. *Saccharomyces cerevisiae* MPS2 encodes a membrane protein localized at the spindle pole body and the nuclear envelope. *Mol Biol Cell* **10**: 2393–2406.
- Deloche O, de la Cruz J, Kressler D, Doere M, Linder P. 2004. A membrane transport defect leads to a rapid attenuation of translation initiation in *Saccharomyces cerevisiae*. *Mol Cell* **13**: 357–366.
- Deshaies RJ, Sanders SL, Feldheim DA, Schekman R. 1991. Assembly of yeast Sec proteins involved in translocation into the endoplasmic reticulum into a membrane-bound multisubunit complex. *Nature* **349**: 806–808.
- Frey S, Pool M, Seedorf M. 2001. Scp160p, an RNA-binding, polysome-associated protein, localizes to the endoplasmic reticulum of *Saccharomyces cerevisiae* in a microtubule-dependent manner. *J Biol Chem* **276**: 15905–15912.
- Galy V, Gadal O, Fromont-Racine M, Romano A, Jacquier A, Nehrbass U. 2004. Nuclear retention of unspliced mRNAs in yeast is mediated by perinuclear Mlp1. *Cell* **116**: 63–73.
- Georgiev A, Sjostrom M, Wieslander A. 2007. Binding specificities of the GYF domains from two *Saccharomyces cerevisiae* paralogs. *Protein Eng Des Sel* **20**: 443–452.
- Gerbasi VR, Weaver CM, Hill S, Friedman DB, Link AJ. 2004. Yeast Asc1p and mammalian RACK1 are functionally orthologous core 40S ribosomal proteins that repress gene expression. *Mol Cell Biol* **24**: 8276–8287.
- Gingras AC, Raught B, Sonenberg N. 1999. eIF4 initiation factors: Effectors of mRNA recruitment to ribosomes and regulators of translation. *Annu Rev Biochem* **68**: 913–963.
- Guo M, Aston C, Burchett SA, Dyke C, Fields S, Rajarao SJ, Uetz P, Wang Y, Young K, Dohlman HG. 2003. The yeast G protein α subunit Gpa1 transmits a signal through an RNA binding effector protein Scp160. *Mol Cell* **12**: 517–524.
- Higashio H, Sato K, Nakano A. 2008. Smy2p participates in COPII vesicle formation through the interaction with Sec23p/Sec24p subcomplex. *Traffic* **9**: 79–93.
- Hogan DJ, Riordan DP, Gerber AP, Herschlag D, Brown PO. 2008. Diverse RNA-binding proteins interact with functionally related sets of RNAs, suggesting an extensive regulatory system. *PLoS Biol* **6**: 2297–2313.
- Ibrahim S, Holmes LE, Ashe MP. 2006. Regulation of translation initiation by the yeast eIF4E binding proteins is required for the pseudohyphal response. *Yeast* **23**: 1075–1088.
- Janke C, Magiera MM, Rathfelder N, Taxis C, Reber S, Maekawa H, Moreno-Borchart A, Doenges G, Schwob E, Schiebel E, et al. 2004. A versatile toolbox for PCR-based tagging of yeast genes: New fluorescent proteins, more markers and promoter substitution cassettes. *Yeast* **21**: 947–962.
- Jaspersen SL, Stearns T. 2008. Exploring the pole: An EMBO conference on centrosomes and spindle pole bodies. *Nat Cell Biol* **10**: 1375–1378.
- Jaspersen SL, Winey M. 2004. The budding yeast spindle pole body: Structure, duplication, and function. *Annu Rev Cell Dev Biol* **20**: 1–28.
- Kofler M, Motzny K, Freund C. 2005. GYF domain proteomics reveals interaction sites in known and novel target proteins. *Mol Cell Proteomics* **4**: 1797–1811.
- Lang BD, Li A, Black-Brewster HD, Fridovich-Keil JL. 2001. The brefeldin A resistance protein Bfr1p is a component of polyribosome-associated mRNP complexes in yeast. *Nucleic Acids Res* **29**: 2567–2574.
- Lau CK, Giddings THJ, Winey M. 2004. A novel allele of *Saccharomyces cerevisiae* *NDC1* reveals a potential role for the spindle pole body component Ndc1p in nuclear pore assembly. *Eukaryot Cell* **3**: 447–458.
- Lazaris-Karatzas A, Montine KS, Sonenberg N. 1990. Malignant transformation by a eukaryotic initiation factor subunit that binds to mRNA 5' cap. *Nature* **345**: 544–547.
- Li AM, Watson A, Fridovich-Keil JL. 2003. Scp160p associates with specific mRNAs in yeast. *Nucleic Acids Res* **31**: 1830–1837.
- Li AM, Vargas CA, Brykailo MA, Openo KK, Corbett AH, Fridovich-Keil JL. 2004. Both KH and non-KH domain

- sequences are required for polyribosome association of Scp160p in yeast. *Nucleic Acids Res* **32**: 4768–4775.
- Lillie SH, Brown SS. 1994. Immunofluorescence localization of the unconventional myosin, Myo2p, and the putative kinesin-related protein, Smy1p, to the same regions of polarized growth in *Saccharomyces cerevisiae*. *J Cell Biol* **125**: 825–842.
- Long RM, Gu W, Lorimer E, Singer RH, Chartrand P. 2000. She2p is a novel RNA-binding protein that recruits the Myo4p–She3p complex to ASH1 mRNA. *EMBO J* **19**: 6592–6601.
- Madrid AS, Mancuso J, Cande WZ, Weis K. 2006. The role of the integral membrane nucleoporins Ndc1p and Pom152p in nuclear pore complex assembly and function. *J Cell Biol* **173**: 361–371.
- Mansfeld J, Guttinger S, Hawryluk-Gara LA, Pante N, Mall M, Galy V, Haselmann U, Muhlhauser P, Wozniak RW, Mattaj JW, et al. 2006. The conserved transmembrane nucleoporin NDC1 is required for nuclear pore complex assembly in vertebrate cells. *Mol Cell* **22**: 93–103.
- Mascarenhas C, Edwards-Ingram LC, Zeef L, Shenton D, Ashe MP, Grant CM. 2008. Gcn4 is required for the response to peroxide stress in the yeast *Saccharomyces cerevisiae*. *Mol Biol Cell* **19**: 2995–3007.
- Matsuo R, Kubota H, Obata T, Kito K, Ota K, Kitazono T, Ibayashi S, Sasaki T, Iida M, Ito T. 2005. The yeast eIF4E-associated protein Eap1p attenuates GCN4 translation upon TOR-inactivation. *FEBS Lett* **579**: 2433–2438.
- Mendelsohn BA, Li AM, Vargas CA, Riehman K, Watson A, Fridovich-Keil JL. 2003. Genetic and biochemical interactions between SCP160 and EAP1 in yeast. *Nucleic Acids Res* **31**: 5838–5847.
- Miao M, Ryan KJ, Wente SR. 2006. The integral membrane protein Pom34p functionally links nucleoporin subcomplexes. *Genetics* **172**: 1441–1457.
- Minshall N, Reiter MH, Weil D, Standart N. 2007. CPEB interacts with an ovary-specific eIF4E and 4E-T in early *Xenopus* oocytes. *J Biol Chem* **282**: 37389–37401.
- Munchow S, Sauter C, Jansen RP. 1999. Association of the class V myosin Myo4p with a localised messenger RNA in budding yeast depends on She proteins. *J Cell Sci* **112**: 1511–1518.
- Nakamura A, Sato K, Hanyu-Nakamura K. 2004. *Drosophila* cup is an eIF4E binding protein that associates with Bruno and regulates oskar mRNA translation in oogenesis. *Dev Cell* **6**: 69–78.
- Niepel M, Strambio-de-Castillia C, Fasalo J, Chait BT, Rout MP. 2005. The nuclear pore complex-associated protein, Mlp2p, binds to the yeast spindle pole body and promotes its efficient assembly. *J Cell Biol* **170**: 225–235.
- Nilsson J, Sengupta J, Frank J, Nissen P. 2004. Regulation of eukaryotic translation by the RACK1 protein: A platform for signalling molecules on the ribosome. *EMBO Rep* **5**: 1137–1141.
- Onischenko E, Stanton LH, Madrid AS, Kieselbach T, Weis K. 2009. Role of the Ndc1 interaction network in yeast nuclear pore complex assembly and maintenance. *J Cell Biol* **185**: 475–491.
- Pereira G, Schiebel E. 2001. The role of the yeast spindle pole body and the mammalian centrosome in regulating late mitotic events. *Curr Opin Cell Biol* **13**: 762–769.
- Pereira G, Schiebel E. 2003. Separase regulates INCENP–Aurora B anaphase spindle function through Cdc14. *Science* **302**: 2120–2124.
- Rapoport TA. 1990. Protein transport across the ER membrane. *Trends Biochem Sci* **15**: 355–358.
- Richter JD, Sonenberg N. 2005. Regulation of cap-dependent translation by eIF4E inhibitory proteins. *Nature* **433**: 477–480.
- Schramm C, Elliott S, Shevchenko A, Shevchenko A, Schiebel E. 2000. The Bbp1p–Mps2p complex connects the SPB to the nuclear envelope and is essential for SPB duplication. *EMBO J* **19**: 421–433.
- Sidrauski C, Walter P. 1997. The transmembrane kinase Ire1p is a site-specific endonuclease that initiates mRNA splicing in the unfolded protein response. *Cell* **90**: 1031–1039.
- Sikorski RS, Hieter P. 1989. A system of shuttle vectors and yeast host strains designed for efficient manipulation of DNA in *Saccharomyces cerevisiae*. *Genetics* **122**: 19–27.
- Sonenberg N, Gingras AC. 1998. The mRNA 5' cap-binding protein eIF4E and control of cell growth. *Curr Opin Cell Biol* **10**: 268–275.
- Sonenberg N, Hinnebusch AG. 2009. Regulation of translation initiation in eukaryotes: Mechanisms and biological targets. *Cell* **136**: 731–745.
- Stavru F, Hulsman BB, Spang A, Hartmann E, Cordes VC, Gorlich D. 2006. NDC1: A crucial membrane-integral nucleoporin of metazoan nuclear pore complexes. *J Cell Biol* **173**: 509–519.
- Stebbins-Boaz B, Cao Q, de Moor CH, Mendez R, Richter JD. 1999. Maskin is a CPEB-associated factor that transiently interacts with eIF-4E. *Mol Cell* **4**: 1017–1027.
- Stegmeier F, Amon A. 2004. Closing mitosis: The function of the Cdc14 phosphatase and its regulation. *Annu Rev Genet* **38**: 203–232.
- Strambio-de-Castillia C, Blobel G, Rout MP. 1999. Proteins connecting the nuclear pore complex with the nuclear interior. *J Cell Biol* **144**: 839–855.
- Tcheperegine SE, Marelli M, Wozniak RW. 1999. Topology and functional domains of the yeast pore membrane protein Pom152p. *J Biol Chem* **274**: 5252–5258.
- Tzur YB, Wilson KL, Gruenbaum Y. 2006. SUN-domain proteins: 'Velcro' that links the nucleoskeleton to the cytoskeleton. *Nat Rev Mol Cell Biol* **7**: 782–788.
- Winey M, Goetsch L, Baum P, Byers B. 1991. MPS1 and MPS2: Novel yeast genes defining distinct steps of spindle pole body duplication. *J Cell Biol* **114**: 745–754.
- Winey M, Hoyt MA, Chan C, Goetsch L, Botstein D, Byers B. 1993. NDC1: A nuclear periphery component required for yeast spindle pole body duplication. *J Cell Biol* **122**: 743–751.
- Wozniak R, Blobel G, Rout MP. 1994. POM152 is an integral protein of the pore membrane domain of the yeast nuclear envelope. *J Cell Biol* **125**: 31–42.
- Xue Z, Shan X, Sinelnikov A, Melese T. 1996. Yeast mutants that produce a novel type of ascus containing asci instead of spores. *Genetics* **144**: 979–989.
- Zeller CE, Parnell SC, Dohlman HG. 2007. The RACK1 ortholog Asc1 functions as a G-protein β subunit coupled to glucose responsiveness in yeast. *J Biol Chem* **282**: 25168–25176.

The SESA network links duplication of the yeast centrosome with the protein translation machinery

Bengü Sezen, Matthias Seedorf and Elmar Schiebel

Genes Dev. 2009, **23**:

Access the most recent version at doi:[10.1101/gad.524209](https://doi.org/10.1101/gad.524209)

Supplemental Material

<http://genesdev.cshlp.org/content/suppl/2009/06/18/23.13.1559.DC1>

References

This article cites 68 articles, 32 of which can be accessed free at:
<http://genesdev.cshlp.org/content/23/13/1559.full.html#ref-list-1>

License

Email Alerting Service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).