REVIEW

MicroRNAs and gene regulatory networks: managing the impact of noise in biological systems

Héctor Herranz¹ and Stephen M. Cohen^{1,2,3}

¹Institute of Molecular and Cell Biology, Singapore 138673; ²Department of Biological Sciences, National University of Singapore, Singapore 117604

Biological systems are continuously challenged by an environment that is variable. Yet, a key feature of developmental and physiological processes is their remarkable stability. This review considers how microRNAs contribute to gene regulatory networks that confer robustness.

The molecular processes that drive cellular functions are inherently noisy. Intrinsic and extrinsic factors can affect levels of gene expression, protein stability, protein modification, enzymatic reactions rates, and signal transduction processes. The ways in which cells and organisms use noise and deal with its impact have become the focus of considerable interest (for review, see Raser and O'Shea 2005; Alon 2007; Barkai and Shilo 2007; Losick and Desplan 2008). This review considers some of the ways in which microRNAs (miRNAs) contribute to these processes. It focuses on the roles of miRNAs in small-scale gene regulatory networks with defined topology, with emphasis on the relationship between miRNAs and noise. The biological properties of miRNAs have been reviewed extensively in recent years, so we assume that the reader is familiar with what miRNAs are and how they work (for reviews, see Bushati and Cohen 2007; Eulalio et al. 2008; Stefani and Slack 2008; Bartel 2009).

Some biological processes involve stochastic decisions, which make use of noise. Examples include the switch between growth and competence phases in *Bacillus subtilis*, and the choice of Rhodopsin type during photoreceptor differentiation in the *Drosophila* eye (Losick and Desplan 2008). In other contexts, noisy gene expression has the potential to be harmful, as is illustrated in some of the examples that follow. Gene regulatory networks involving positive and negative feedback have the potential to control the effects of noise, by buffering its impact on gene expression (e.g., Lee et al. 2002; Milo et al. 2002). Computational analysis has suggested that miRNAs are overrepresented in gene regulatory networks, suggesting

that they confer useful regulatory possibilities (Tsang et al. 2007; Martinez et al. 2008).

Complex networks often include a small set of recurring circuits called network motifs (for review, see Alon 2007). These motifs are found more often than would be expected by chance in gene regulatory networks, from bacteria and yeast to plants and animals. Network motifs can be subdivided into two broad categories: feedback and feedforward. The utility of positive and negative feedback regulation is intuitively obvious (Fig. 1A). For example, a transcription factor can limit its own activity by inducing a repressor. Alternately, a positive feedback loop can produce a robust developmental switch by ensuring an "all-or-none" outcome. Feedforward motifs are less intuitive, and are based on regulators that act both directly and indirectly on their downstream targets (Fig. 2A). In the following sections, we consider the roles that miRNAs play in feedback and feedforward networks.

Feedback motifs as switches

Using noise

Evidence is beginning to emerge that miRNAs can be involved in the mechanisms by which noise is used to trigger stochastic developmental decisions. Sense organ specification in the *Drosophila* peripheral nervous system involves a key element of stochasticity. Sense organ primordia are initially defined as small groups of cells that express a set of "proneural" basic helix-loop-helix (bHLH) transcription factors of the Hairy-E(spl), or HES, family. Definition of the proneural cluster at this level is deterministic, based on programmed control of gene expression, but the choice of which cell will become the sense organ precursor (SOP) is stochastic. One cell is selected by chance, as a consequence of fluctuation in the level of the transcription factor Senseless (Nolo et al. 2000). Higher Senseless levels lead to an increase in proneural gene expression in the SOP, which feeds back to increase *Senseless* expression (Fig. 1B). As well, Senseless increases expression of the Notch ligand *Delta*, which in turn leads to increased Notch signaling activity in the adjacent cells (Fig. 1B). Notch activity represses

[Keywords: MicroRNAs; noise; robustness; regulatory networks]

³Corresponding author.

E-MAIL scohen@imcb.a-star.edu.sg; FAX 65-6872-7089.

Article is online at <http://www.genesdev.org/cgi/doi/10.1101/gad.1937010>.

Figure 1. miRNAs involved in feedback motifs. (A) Simple positive and negative feedback loops. (B) *miR-9a* in a positive feedback loop in SOP specification in *Drosophila*. The boxes represent cells within the proneural cluster. In the blue cell, the Senseless–proneural gene positive feedback circuit dominates and leads to SOP differentiation. Senseless activates Delta, increasing Notch activity in adjacent cells. Notch inhibits proneural genes, allowing *miR-9a* repression of Senseless to dominate in these cells. (C) *lsy-6* and *miR-273* in a negative feedback loop controlling left/right asymmetry in ASE neurons of *C. elegans*. The transcription factors encoded by *cog-1* and *die-1* are active in ASER and ASEL, respectively. *cog-1* activates *miR-273* miRNAs, which repress *die-1* in ASER. *die-1* activates *lsy-6*, which represses *cog-1* in ASEL. This initial asymmetry is stabilized by positive autoregulation of *cog-1* in ASER. (D) *miR-14* acts in a negative feedback loop modulating EcR activity. EcR positively autoregulates. *miR-14* limits EcR levels, which in turn negatively regulates *miR-14*. (E) *miR-124* in a negative feedback loop in the mouse cortex. The open boxes represent cells. Ephrin-B1 activity represses *miR-124* in neural progenitors. *miR-124* inhibits Ephrin-B1 and induces neuronal differentiation. (F) *miR-375* acts in a negative feedback regulating insulin secretion. High glucose levels inhibit *miR-375*. *miR-375* inhibits myotrophin and Pdkp-1, reducing insulin secretion. Circulating glucose promotes insulin secretion directly, and indirectly represses *miR-375*.

proneural gene expression in these cells, and reinforces the advantage of the *Senseless*- and *Delta*-expressing cell toward becoming the SOP (Jafar-Nejad et al. 2003). This system combines positive feedback with lateral inhibition to select a single cell and promote its differentiation into the SOP. However, more recent studies indicate this circuit is not enough to ensure a reliable outcome (Li et al. 2006; for review, see Cohen et al. 2006). Flies lacking *miR-9a* sporadically produce extra sense organs from individual proneural clusters. This indicates that more than one cell in the cluster can cross the threshold level of *Senseless* activity required to trigger the SOP switch. Feedback regulation is an essential element of the mechanisms that transform noise into switches (for review, see Losick and Desplan 2008), but positive feedback loops are inherently labile. Additional mechanisms are needed to ensure that random fluctuation in transcription factor levels does trigger positive feedback inappropriately. In this biological context, the presence of *miR-9a* sets a threshold that *Senseless* transcription must overcome in order to trigger the *Senseless*–proneural feedback loop (Fig. 1B).

An important feature of this circuit is that *miR-9a* expression is under proneural gene control. *miR-9a* levels are initially uniform in the proneural cluster, but as

Senseless activity increases proneural gene expression in the SOP, *miR-9a* levels are reduced. *miR-9a* levels remain high in the surrounding cells, where proneural gene activity is kept low through Notch activity. The opposing regulation of *Senseless* and *miR-9a* is an integral element of the switch. The outcome is a situation in which initially overlapping expression domains resolve into spatially reciprocal domains of the miRNA, and its target, Reciprocity, is a common feature of miRNA and target expression (Stark et al. 2005) and serves to reinforce the differences between the cell types. The transcriptional feedback system is triggered by random fluctuations in gene expression, and the miRNA helps to reinforce selection of only one peak of *Senseless* expression while filtering out others that might have been able to surpass the threshold in the absence of the miRNA. Thus, *miR-9a* contributes to the robustness of the noise-dependent switch.

lsy-6 and *miR-273* contribute to a bistable switch

The nematode *Caenorhabditis elegans* makes a pair of chemosensory neurons that are functionally asymmetric on the left and right sides of the animal. Work from the Hobert laboratory (for review see Hobert 2006) has shown

Figure 2. miRNAs involved in feedforward motifs. (A) Feedforward motifs where X regulates Z directly and indirectly through regulation of Y. Coherent motifs have the direct and indirect paths from X acting on the target Z in the same direction. Incoherent motifs have opposite signs for the two paths. (B) *miR-7* acts in two coherent feedforward motifs. Yan is a *miR-7* target. In one motif (gray), Yan represses *miR-7* directly and indirectly. In the indirect arm, Yan represses Phyllopod, alleviating repression of Ttk-69, which represses *miR-7*. In the second motif (red), Pnt-P1 directly activates *miR-7*, which represses Yan. Pnt-P1 represses *yan* directly. (C) *let-7* acts in a coherent feedforward motif (gray), modulating oncogenic transformation. NF- κ B induces IL6 expression directly. In the indirect arm, NF- κ B activity inhibits *let-7*, alleviating repression of IL6. *let-7* prevents noise in the activity of NF- κ B or IL6 from triggering the positive feedback loop (red). (D) *miR-7* acts in an incoherent feedforward motif. Atonal activates E(spl) directly. Atonal also represses E(spl) via *miR-7*. Atonal activity results in a pulse of E(spl) followed by a lower level of steady-state expression due to Atonal-induced *miR-7* activity. (E) *miR-20* and *miR-17-5p* act in an incoherent feedforward motif. c-Myc activates E2F directly. c-Myc also represses E2F via *miR-17-5p* and *miR-20*. c-Myc activity results in a pulse of E2F to promote cell cycle progression.

that miRNAs work together with transcription factors to produce a switch that allows these cells to adopt one of two alternative fates, called ASER/R (Fig. 1C). The homeobox-containing transcription factor encoded by *cog-1* is active in ASER, and the zinc finger transcription factor encoded by *die-1* is active in ASEL. In ASEL cells, *die-1* induces expression of the *lsy-6* miRNA, which has been shown to limit *cog-1* activity in vivo. Reciprocally, in ASER, *cog-1* induces expression of *miR-273* family miRNAs, which are thought to act by repressing *die-1*. The asymmetry between the two cells is initially stabilized by positive autoregulation of *cog-1* in ASER. The combination of miRNA-mediated repression and positive autoregulation produces a switch that drives the cell into one of two states. However, genetic evidence indicates that the two cells are initially identical. Both cells can adopt either ASER or ASEL identity if components of the feedback loops are missing. This initial state is labile, and is potentially subject to perturbation by noisy gene expression. A key issue that remains to be fully

understood is how an initial bias is introduced into the system to allow the switch to be thrown with a consistent left/right outcome.

miR-14 limits the impact of noise on a developmental switch

Drosophila miR-14 has been implicated in a different type of regulatory switch, which serves to buffer against noise (Varghese and Cohen 2007). The steroid hormone ecdysone and the ecdysone receptor (EcR) regulate major transformations in gene expression programs as the animal transitions from one developmental stage to the next (for review, see Thummel 2001). An important feature of these transitions is their all-or-none character. This depends on a positive feedback loop involving transcriptional autoregulation by EcR. A loop in which a transcription factor controls its own expression is inherently labile. Transcription often occurs in bursts (Raj and van Oudenaarden 2008), which are expected to lead to fluctuation in activity levels. In principle, a transcription burst for an autoregulating transcription factor such as EcR could trigger a self-amplifying response. Linking EcR autoregulation to a negative feedback loop involving *miR-14* provides a means to limit the potential impact of noise (Fig. 1D). *miR-14* acts directly to reduce EcR levels, and, reciprocally, EcR negatively regulates *miR-14* transcription. Mutual repression keeps the levels of *miR-14* and EcR in balance, until the input of a hormonal cue activates EcR. This will shut off production of new *miR-14*, but the preexisting miRNA takes time to decay. This delay permits the cell to discriminate between sustained input of hormone-induced EcR activation and transient fluctuation in EcR levels due to transcriptional bursts. The use of a miRNA to minimize the impact of noise permits the safe use of a positive autoregulatory loop in a switch capable of profoundly changing the gene expression program of the animal.

Repression of a repressor in a positive feedback loop: miR-9a

The regulatory network involving *miR-9a* can also be seen as an indirect feedback loop. *miR-9a* represses Senseless, which serves as an activator of the proneural genes. By repressing *miR-9a*, the proneural transcription factors remove repression of their activator, Senseless. Because the proneural genes promote *Senseless* expression, the net result is a positive autoregulatory loop (Fig. 1B). As in the case of *miR-14*, the preexisting expression of the miRNA limits the impact of noise until it rises above a threshold level. Thereafter, the cell commits to a program of differentiation. An independent loop involving Delta and Notch conveys this information to influence the behavior of the other cells in the group.

Negative feedback in neuronal differentiation: miR-124 and ephrin-B1

Mutual inhibition of *ephrin-B1* and *miR-124* in the developing mouse brain illustrates a negative feedback

interaction controlling neural differentiation (Arvanitis et al. 2010). Ephrins serve as both ligands and receptors that convey bidirectional signals reflecting cell–cell contact (for review, see Davy and Soriano 2005). *Ephrin-B1* and *miR-124* show reciprocal expression during cortex differentiation. *ephrin-B1* is expressed in neural progenitor layers, whereas *miR-124* expression is elevated in more differentiated cells. Ephrin-B1 signaling limits *miR-124* levels, and, reciprocally, *miR-124* expression can inhibit *ephrin-B1* expression. Thus, mutual inhibition between *miR-124* and *ephrin-B1* controls neuronal differentiation, where ephrin-B1 is required for maintaining the progenitor fate and *miR-124* promotes differentiation (Fig. 1E). Mutual repression may help to sharpen expression boundaries and confer robustness during differentiation.

Negative feedback in epithelial–mesenchymal transition (EMT): the miR-200 family and ZEB/Sip1

EMT is a fundamental cellular transformation that contributes to shaping tissues during development. EMT also plays a key role during metastasis. Members of the *miR-200* miRNA family are expressed in epithelial cells, where they limit expression of the E-box transcription factors ZEB1 and SIP1, thereby allowing expression of E-Cadherin (Gregory et al. 2008; Park et al. 2008). ZEB1 and SIP1 are induced by TGF- β to promote EMT through repression of E-Cadherin. ZEB1 and SIP1 also repress expression of the *miR-200* family of miRNAs (Bracken et al. 2008). The mutually repressive regulatory relationship is similar to that between *miR-124* and Ephrin-B1. Mutual inhibition helps to reinforce the switch between cell fates, and may help to ensure that low-level noisy expression of ZEB1 or SIP1 cannot accidentally trigger EMT prior to the onset of robust TGF- β -induced expression. During EMT, the cells do not remain together as part of a tissue, so the reciprocity may be thought of as being more temporal than spatial.

A miRNA-regulated feedback loop in glucose homeostasis

Energy homeostasis depends on insulin signaling. Insulin levels reflect the nutritional status of the animal to control circulating sugar levels and to regulate glycogen and fat metabolism. Insulin production and secretion as well as cellular responsiveness to insulin are subject to regulation by miRNAs. miRNAs have been shown to act in many tissues involved in glucose metabolism, including the pancreatic islet, liver, skeletal muscle, adipose tissue, and brain (for review, see Lynn 2009). In vivo, mice mutant for *miR-375* show reduced pancreatic β -cell mass and defects in energy homeostasis (Poy et al. 2009). In assays using insulin-producing cell lines, *miR-375* has been shown to negatively regulate glucose-stimulated insulin release (Poy et al. 2004). *miR-375* has been reported to act via its target myotrophin to control a late step during insulin exocytosis. Although the mechanism is not fully understood, it probably involves regulation of actin polymerization and vesicular fusion. miRNAs are also thought to regulate insulin levels by other means.

Pdk1, a 3'-phosphoinositide-dependent protein kinase, is a direct target of *miR-375*. Depletion of *miR-375* has been reported to increase Pdk-1 levels and to increase insulin expression, also in a cell-based assay system (El Ouaamari et al. 2008). Intriguingly, *miR-375* expression appears to be regulated by glucose levels in rat pancreatic islets. Thus, glucose sensing appears to act via repression of *miR-375* to promote insulin secretion, contributing to glucose homeostasis (Fig. 1F).

Managing noise: coherent feedforward motifs as buffers

Feedforward motifs involve at least three elements: an upstream regulator and two targets (illustrated in Fig. 2A; for review, see Alon 2007). The upstream element X acts on one target, Z, directly as well as indirectly, via the second target, Y. In cases in which both branches act in the same direction on Z, the motif is called “coherent.” If they act oppositely, the motif is called “incoherent.” Motifs of this sort can have many possible outcomes, depending on the nature of the relationships between the elements. For example, a coherent motif in which X and Y are both needed to activate Z introduces a delay in the induction of Z until Y can accumulate to the required level. A coherent motif in which either X or Y can activate Z allows for rapid onset of Z but delays turning Z off, to an extent that depends on the stability of Y. Incoherent motifs have different properties; for example, they can allow for rapid onset followed by repression to produce a pulse-like activation profile. Alon (2007) provides an in-depth review of the properties of these motifs. In the following sections, we focus on examples that involve miRNAs.

Coherent feedforward motifs in photoreceptor differentiation

miR-7 has been implicated in two coherent feedforward motifs involved in the specification of photoreceptor cells in the *Drosophila* eye (Fig. 2B; Li et al. 2009). Yan, a transcription factor essential for the development of photoreceptor cells, is a direct target of *miR-7*. In one circuit, Yan represses *miR-7* transcription directly as well as indirectly (Fig. 2B, shown in gray). The indirect arm in this case involves three elements: Yan represses Phyllopod, thereby alleviating repression of Ttk-69, which allows Ttk-69 to repress *miR-7* (the triple negative, repression of a repressor of a repressor results in repression). Repression by either branch is sufficient to limit *miR-7* expression. This makes the system less sensitive to transient fluctuation in the levels of either Yan or Ttk-69. Concurrently, *miR-7* is involved in a second coherent feedforward motif in the same cells. The transcription factor Pnt-P1 directly activates *miR-7* transcription, which in turn represses Yan. Pnt-P1 also acts directly to represses *yan* transcription. This circuit buffers Yan expression to variations in the levels of Pnt-P1 or *miR-7*. The Yan OFF state is buffered against transient fluctuation in Pnt-P1 levels, but can switch if activation is prolonged. Likewise the Yan ON state is stable. It is likely that use of two

interlinked motifs with elements in common provides considerable stability to photoreceptor specification. Indeed, Li et al. (2009) provide experimental evidence that the system is normally very stable, but that it can be perturbed by environmental fluctuation—temperature shifts in this instance—if *miR-7* function is compromised.

A coherent feedforward motif limits the impact of noise in an oncogenic switch

A recent study has implicated the human *let-7* miRNA in a molecular pathway underlying oncogenic transformation in response to an inflammatory signal (Iliopoulos et al. 2009). Loss of *let-7* has been implicated previously in cancer (e.g., Johnson et al. 2005). *let-7* activity normally serves to limit expression of the cytokine interleukin-6 (IL6), a key mediator of the inflammatory response. Activation of NF- κ B can block *let-7* family miRNA biogenesis through up-regulation of lin-28 (which prevents *let-7* processing) (Viswanathan et al. 2008). Sustained NF- κ B activation thereby leads to depletion of *let-7* and alleviates repression of IL6. This comprises the indirect arm of the feedforward motif (Fig. 2C, gray). NF- κ B also acts directly to induce IL6 expression. Activation of both arms of the feedforward motif is required to achieve the high levels of IL6 activity required for cellular transformation (Iliopoulos et al. 2009), which is aided by positive feedback regulation of IL6 on NF- κ B (Fig. 2C, red). In essence, the role of *let-7* is comparable with that of *miR-14* in the steroid hormone switch described above. By limiting levels of IL6, *let-7* prevents noise in the activity of NF- κ B or IL6 from triggering the positive feedback loop that leads to cellular transformation. Only sustained activation of NF- κ B can allow the decay of miRNA expression required for the IL6/NF- κ B feedback loop to become self-sustaining. Comparing *let-7* and *miR-14* illustrates how similar biological outputs can be obtained by regulatory networks of different topology.

Incoherent feedforward motifs as triggers

The *Drosophila* sensory nervous system is proving to be a rich source of feedforward regulatory motifs. Work from the Carthew laboratory (Li et al. 2009) has also implicated *miR-7* in an incoherent feedforward motif that contributes to sense organ specification. In this context, the transcription factor Atonal acts directly to activate expression of the proneural gene *E(spl)*, illustrated in Figure 2D. Atonal acts concurrently via *miR-7* to repress *E(spl)* genes. Activation of Atonal results in a pulse of *E(spl)* expression followed by a lower level of steady-state expression. The pulse of *E(spl)* can serve as a trigger for other regulatory events, including those involving *miR-9a* and *Senseless*.

A second example of an incoherent regulatory motif involves two miRNAs of the human *miR-17* cluster in regulating cell cycle progression (Fig. 2E; O'Donnell et al. 2005). c-Myc is a potent inducer of cell proliferation, and is sufficient to promote tumor formation (Pelengaris et al. 2002). c-Myc acts directly to induce expression of the

transcription factor *E2F*, which in turn regulates an array of cell cycle genes. c-Myc acts concurrently to induce expression of *miR-17-5p* and *miR-20*, which then repress *E2F*. This leads to a pulse of c-Myc-induced *E2F* expression to promote cell cycle progression. Incoherent feedforward motifs are well suited as triggers for oscillatory systems. A potentially important implication of the use of such motifs is that the output can be shorter-lived than the input.

Conclusions

The studies discussed here represent only a small fraction of what has been learned in recent years about the roles of miRNAs in vivo. They represent the cases in which it has been possible to place the miRNA into a gene regulatory circuit, and from this it has been possible to understand how the miRNA confers a specific mode of biological regulation. These circuits can have quite different roles, from using noise to throw a developmental switch to buffering the consequences of noise in order to confer robustness to environmental perturbation.

Acknowledgments

We thank Ruifen Weng, Xin Hong, Jan-Michael Kugler, and Jishy Varghese for discussions and comments on the manuscript.

References

- Alon U. 2007. Network motifs: Theory and experimental approaches. *Nat Rev Genet* **8**: 450–461.
- Arvanitis DN, Jungas T, Behar A, Davy A. 2010. Ephrin-B1 reverse signaling controls a post-transcriptional feedback mechanism via miR-124. *Mol Cell Biol* **30**: 2508–2517.
- Barkai N, Shilo BZ. 2007. Variability and robustness in biochemical systems. *Mol Cell* **28**: 755–760.
- Bartel DP. 2009. MicroRNAs: Target recognition and regulatory functions. *Cell* **136**: 215–233.
- Bracken CP, Gregory PA, Kolesnikoff N, Bert AG, Wang J, Shannon MF, Goodall GJ. 2008. A double-negative feedback loop between ZEB1-SIP1 and the microRNA-200 family regulates epithelial-mesenchymal transition. *Cancer Res* **68**: 7846–7854.
- Bushati N, Cohen SM. 2007. MicroRNA functions. *Annu Rev Cell Dev Biol* **23**: 175–205.
- Cohen SM, Brennecke J, Stark A. 2006. Denoising feedback loops by thresholding—a new role for microRNAs. *Genes Dev* **20**: 2769–2772.
- Davy A, Soriano P. 2005. Ephrin signaling in vivo: Look both ways. *Dev Dyn* **232**: 1–10.
- El Ouamari A, Baroukh N, Martens GA, Lebrun P, Pipeleers D, van Obberghen E. 2008. miR-375 targets 3'-phosphoinositide-dependent protein kinase-1 and regulates glucose-induced biological responses in pancreatic β -cells. *Diabetes* **57**: 2708–2717.
- Eulalio A, Huntzinger E, Izaurralde E. 2008. Getting to the root of miRNA-mediated gene silencing. *Cell* **132**: 9–14.
- Gregory PA, Bert AG, Paterson EL, Barry SC, Tsykin A, Farshid G, Vadas MA, Khew-Goodall Y, Goodall GJ. 2008. The miR-200 family and miR-205 regulate epithelial to mesenchymal transition by targeting ZEB1 and SIP1. *Nat Cell Biol* **10**: 593–601.
- Hobert O. 2006. Architecture of a microRNA-controlled gene regulatory network that diversifies neuronal cell fates. *Cold Spring Harb Symp Quant Biol* **71**: 181–188.

- Iliopoulos D, Hirsch HA, Struhl K. 2009. An epigenetic switch involving NF- κ B, Lin28, Let-7 microRNA, and IL6 links inflammation to cell transformation. *Cell* **139**: 693–706.
- Jafar-Nejad H, Acar M, Nolo R, Lacin H, Pan H, Parkhurst SM, Bellen HJ. 2003. Senseless acts as a binary switch during sensory organ precursor selection. *Genes Dev* **17**: 2966–2978.
- Johnson SM, Grosshans H, Shingara J, Byrom M, Jarvis R, Cheng A, Labourier E, Reinert KL, Brown D, Slack FJ. 2005. RAS is regulated by the let-7 microRNA family. *Cell* **120**: 635–647.
- Lee TI, Rinaldi NJ, Robert F, Odom DT, Bar-Joseph Z, Gerber GK, Hannett NM, Harbison CT, Thompson CM, Simon I, et al. 2002. Transcriptional regulatory networks in *Saccharomyces cerevisiae*. *Science* **298**: 799–804.
- Li Y, Wang F, Lee JA, Gao FB. 2006. MicroRNA-9a ensures the precise specification of sensory organ precursors in *Drosophila*. *Genes Dev* **20**: 2793–2805.
- Li X, Cassidy JJ, Reinke CA, Fischboeck S, Carthew RW. 2009. A microRNA imparts robustness against environmental fluctuation during development. *Cell* **137**: 273–282.
- Losick R, Desplan C. 2008. Stochasticity and cell fate. *Science* **320**: 65–68.
- Lynn FC. 2009. Meta-regulation: MicroRNA regulation of glucose and lipid metabolism. *Trends Endocrinol Metab* **20**: 452–459.
- Martinez NJ, Ow MC, Barrasa MI, Hammell M, Sequerra R, Doucette-Stamm L, Roth FP, Ambros VR, Walhout AJ. 2008. A *C. elegans* genome-scale microRNA network contains composite feedback motifs with high flux capacity. *Genes Dev* **22**: 2535–2549.
- Milo R, Shen-Orr S, Itzkovitz S, Kashtan N, Chklovskii D, Alon U. 2002. Network motifs: Simple building blocks of complex networks. *Science* **298**: 824–827.
- Nolo R, Abbott LA, Bellen HJ. 2000. Senseless, a Zn finger transcription factor, is necessary and sufficient for sensory organ development in *Drosophila*. *Cell* **102**: 349–362.
- O'Donnell KA, Wentzel EA, Zeller KI, Dang CV, Mendell JT. 2005. c-Myc-regulated microRNAs modulate E2F1 expression. *Nature* **435**: 839–843.
- Park SM, Gaur AB, Lengyel E, Peter ME. 2008. The miR-200 family determines the epithelial phenotype of cancer cells by targeting the E-cadherin repressors ZEB1 and ZEB2. *Genes Dev* **22**: 894–907.
- Pelengaris S, Khan M, Evan GI. 2002. Suppression of Myc-induced apoptosis in β cells exposes multiple oncogenic properties of Myc and triggers carcinogenic progression. *Cell* **109**: 321–334.
- Poy MN, Eliasson L, Krutzfeldt J, Kuwajima S, Ma X, Macdonald PE, Pfeffer S, Tuschl T, Rajewsky N, Rorsman P, et al. 2004. A pancreatic islet-specific microRNA regulates insulin secretion. *Nature* **432**: 226–230.
- Poy MN, Hausser J, Trajkovski M, Braun M, Collins S, Rorsman P, Zavolan M, Stoffel M. 2009. miR-375 maintains normal pancreatic α - and β -cell mass. *Proc Natl Acad Sci* **106**: 5813–5818.
- Raj A, van Oudenaarden A. 2008. Nature, nurture, or chance: Stochastic gene expression and its consequences. *Cell* **135**: 216–226.
- Raser JM, O'Shea EK. 2005. Noise in gene expression: Origins, consequences, and control. *Science* **309**: 2010–2013.
- Stark A, Brennecke J, Bushati N, Russell RB, Cohen SM. 2005. Animal microRNAs confer robustness to gene expression and have a significant impact on 3'UTR evolution. *Cell* **123**: 1133–1146.
- Stefani G, Slack FJ. 2008. Small non-coding RNAs in animal development. *Nat Rev Mol Cell Biol* **9**: 219–230.
- Thummel CS. 2001. Molecular mechanisms of developmental timing in *C. elegans* and *Drosophila*. *Dev Cell* **1**: 453–465.
- Tsang J, Zhu J, van Oudenaarden A. 2007. MicroRNA-mediated feedback and feedforward loops are recurrent network motifs in mammals. *Mol Cell* **26**: 753–767.
- Varghese J, Cohen SM. 2007. MicroRNA miR-14 acts to modulate a positive autoregulatory loop controlling steroid hormone signaling in *Drosophila*. *Genes Dev* **21**: 2277–2282.
- Viswanathan SR, Daley GQ, Gregory RI. 2008. Selective blockade of microRNA processing by Lin28. *Science* **320**: 97–100.

MicroRNAs and gene regulatory networks: managing the impact of noise in biological systems

Héctor Herranz and Stephen M. Cohen

Genes Dev. 2010, **24**:

Access the most recent version at doi:[10.1101/gad.1937010](https://doi.org/10.1101/gad.1937010)

References

This article cites 36 articles, 16 of which can be accessed free at:
<http://genesdev.cshlp.org/content/24/13/1339.full.html#ref-list-1>

License

Email Alerting Service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).