Growth factor stimulation induces a distinct ER α cistrome underlying breast cancer endocrine resistance

Mathieu Lupien,^{1,2,3,8} Clifford A. Meyer,^{4,8} Shannon T. Bailey,^{1,2} Jérôme Eeckhoutte,⁵ Jennifer Cook,^{1,2} Thomas Westerling,^{1,2} Xiaoyang Zhang,³ Jason S. Carroll,⁶ Daniel R. Rhodes,⁷ X. Shirley Liu,^{2,10} and Myles Brown^{1,2,9}

¹Division of Molecular and Cellular Oncology, Department of Medical Oncology, Dana-Farber Cancer Institute, Boston, Massachusetts 02115, USA; ²Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts 02115, USA; ³Department of Genetics, Norris Cotton Cancer Center, Dartmouth Medical School, Lebanon, New Hampshire 03756, USA; ⁴Department of Biostatistics and Computational Biology, Dana-Farber Cancer Institute and Harvard School of Public Health, Boston, Massachusetts 02115, USA; ⁵UMR CNRS 6026, Université de Rennes 1, 35042 Rennes, France; ⁶Cancer Research UK, Cambridge Research Institute, Cambridge CB2 0RE, United Kingdom; ⁷Compendia Bioscience, Inc., Ann Arbor, Michigan 48104, USA

Estrogen receptor α (ER α) expression in breast cancer is predictive of response to endocrine therapy; however, resistance is common in ER α -positive tumors that overexpress the growth factor receptor ERBB2. Even in the absence of estrogen, ER α can be activated by growth factors, including the epidermal growth factor (EGF). EGF induces a transcriptional program distinct from estrogen; however, the mechanism of the stimulus-specific response is unknown. Here we show that the EGF-induced ER α genomic targets, its cistromes, are distinct from those induced by estrogen in a process dependent on the transcription factor AP-1. The EGF-induced ER α cistrome specifically regulates genes found overexpressed in ERBB2-positive human breast cancers. This provides a potential molecular explanation for the endocrine therapy resistance seen in ER α -positive breast cancers that overexpress ERBB2. These results suggest a central role for ER α in hormone-refractory breast tumors dependent on growth factor pathway activation and favors the development of therapeutic strategies completely antagonizing ER α , as opposed to blocking its estrogen responsiveness alone.

[**Keywords:** ERBB2; breast cancer; cistrome; estrogen receptor; growth factors; transcription]

Supplemental material is available at <http://www.genesdev.org>.

Received September 10, 2009; revised version accepted August 16, 2010.

More than two-thirds of human breast cancers overexpress the estrogen receptor α (ER α), where it is both a target of endocrine therapy and a predictor of response (Sorlie et al. 2001, 2003). Upon activation by estrogen, ER α is recruited to thousands of sites across the genome of human breast cancer cells, defining its cistrome (Carroll et al. 2005, 2006; Lin et al. 2007; Hua et al. 2008; Hurtado et al. 2008; Liu et al. 2008; Lupien et al. 2008; Fullwood et al. 2009). This process is highly organized through specific epigenetic events that restrict the recruitment of the receptor to a subset of its potential binding sites (Lupien et al. 2008). Accordingly, this ER α cistrome guides the response to estrogen in breast cancer cells by favoring the implementation of an ER α -positive breast tumor-specific transcrip-

tional program (Carroll et al. 2006). The importance of the cistrome in defining a specific transcriptional program is further supported by work in osteosarcoma cells. Indeed, estrogen stimulation in these cells results in a distinct expression profile (Monroe et al. 2003; Krum et al. 2008) that is directly related to a unique ER α cistrome (Krum et al. 2008). Such lineage-specific recruitment patterns were also reported recently for other factors. Indeed, cell type-specific transcriptional programs associated with FoxA1 and p300 are linked to their lineage-specific cistromes (Lupien et al. 2008; Visel et al. 2009). Thus, the contribution of a given transcription factor to the execution of a specific transcriptional program is highly dependent on its cell type-specific cistrome.

In addition to cell type-specific transcription programs, ER α can also respond to a variety of stimuli in a given cell type. In breast cancer cells, ER α can be stimulated even in the absence of estrogen by a variety of growth factors, including epidermal growth factor (EGF) (Kato et al. 1995; Bunone et al. 1996; Joel et al. 1998; Smith 1998; Kato 2001;

⁸These authors contributed equally to this work.

Corresponding authors.

⁹E-MAIL myles_brown@dfci.harvard.edu; FAX (617) 632-5417.

¹⁰E-MAIL xslu@jimmy.harvard.edu; FAX (617) 632-2444.

Article is online at <http://www.genesdev.org/cgi/doi/10.1101/gad.1944810>.

Kurokawa and Arteaga 2003). Indeed, growth factor-stimulated breast cancer proliferation as well as normal uterine growth is dependent on ER α (Ignar-Trowbridge et al. 1992; Kato et al. 1995; Lee and Yee 1995; Curtis et al. 1996; Lupu et al. 1996; Knowlden et al. 2003; Schiff et al. 2005). However, the transcriptional response induced by growth factor pathway stimulation in breast cancer cells differs from that of estrogen (Cunliffe et al. 2003; Dudek and Picard 2008). Furthermore, ER α -positive breast cancers that overexpress the EGF receptor-2 (ERBB2/HER2) are resistant to endocrine therapies that disrupt the estrogen-dependent ER α program. In the present study, we addressed the role of growth factor-specific ER α -dependent transcriptional responses in breast cancer cells.

Results

Growth factor-induced unique ER α -dependent transcriptional program

Given that growth factor pathway stimulation activates a number of transcription factors (Moasser 2007), we first assessed the contribution of ER α in EGF-mediated breast cancer cell proliferation. ER α depletion using the full antagonist fulvestrant (100 nM) or siRNA against ER α (Lupien et al. 2007, 2008) abrogated the EGF-mediated proliferation of MCF7 breast cancer cells (Fig. 1A; Supplemental Fig. S1A,B). Partial depletion using a lower dose of fulvestrant had reduced effects (Fig. 1A). ER α depletion also completely abrogated cellular proliferation triggered by activation of ERBB2 by heregulin in the ER α -

positive BT474 breast cancer cell line (Supplemental Fig. S1C). Hence, ER α contributes significantly to growth factor pathway-mediated proliferation of breast cancer cells, including those overexpressing ERBB2.

While EGF induced proliferation of MCF7 breast cancer cells to the same extent as estrogen, this involved a distinct transcriptional program (Fig. 1A,B; Cunliffe et al. 2003; Dudek and Picard 2008). ER α depletion using fulvestrant in EGF-treated cells revealed that >39% and 24% of EGF up-regulated and down-regulated genes in MCF7 cells, respectively, were fully or partially dependent on ER α (Fig. 1B,C). Similar results were obtained when silencing ER α expression using siRNA (Supplemental Fig. S1D,E). Noteworthy, by defining the EGF up-regulated transcriptional program in MCF7 cells, we found, using OncoPrint Concepts Map analysis (Rhodes et al. 2007), that this program was correlated with the most highly expressed genes in ERBB2-positive breast tumors (odds ratio ≥ 2 , $P \leq 1e-2$) (Fig. 1D). Conversely the EGF down-regulated genes in MCF7 cells correlated with the most repressed genes from ERBB2-positive breast tumors (Supplemental Fig. S2). Importantly, these associations were also observed for the ER α -dependent EGF up-regulated or down-regulated transcriptional program (Fig. 1D; Supplemental Fig. S2). In

Figure 1. ER α is required for growth factor-mediated breast cancer cell proliferation. (A) Proliferation of MCF7 breast cancer cells pretreated or not with the full anti-estrogen fulvestrant (Ful) was measured following EGF stimulation. (*) $P \leq 0.05$; (**) $P \leq 0.01$; (***) $P \leq 0.001$. (B) Comparison of estrogen-up (E2), EGF-up, and ER α -dependent EGF up-regulated as well as down-regulated transcriptional programs in MCF7 breast cancer cells. (C) mRNA levels derived from RT-qPCR of the ER α -dependent EGF target genes *LIF* and *ACP6* are presented under EGF stimulation in MCF7 cells pretreated or not with Ful (100 nM). *TFF1* is used as a negative control. (D) OncoPrint Concepts Map analysis (Compendia Biosciences, Inc.; <https://www.oncoPrint.com>) was used to compare the EGF-induced gene signatures from primary breast tumors. This revealed significant correlations between EGF-up as well as ER α -dependent EGF-up gene signatures from MCF7 cells with gene signatures from poor-outcome (metastasis, recurrence, death, and high grade) as well as ERBB2-positive breast tumors ($P \leq 1e-2$, odds ratio [O.R.] ≥ 2). No significant correlations were revealed between EGF-up or ER α -dependent EGF-up gene signatures and expression signatures from ER α -positive primary breast tumors. Each green circle in the *left* figure corresponds to the gene signature from primary breast tumors established in an independent study. The red circle corresponds to the EGF-up gene signature in MCF7 breast cancer cells. The purple circle corresponds to the ER α -dependent EGF-up gene signature from MCF7 breast cancer cells. Genes significantly correlated with each other are linked to each other by a straight line. The *right* figure presents an example of how significant correlation between EGF-up and ER α -dependent EGF-up gene signatures with the ERBB2-positive breast cancer gene signature were established in one published study (Richardson et al. 2006). Specifically, the expression profile in primary breast tumors from different patients (vertical axis) is presented for the genes found in the EGF-up and ER α -dependent EGF-up gene signature from MCF7 breast cancer cells (horizontal axis).

addition, both the EGF-up and ER α -dependent EGF-up transcriptional programs were associated with poor-outcome expression signatures such as relapse, death, metastasis, and high tumor grade (Fig. 1D). Overall, these new results reveal the capacity of ER α to elicit stimuli-specific transcriptional programs in breast cancer cells. Furthermore, the association of the ER α -dependent EGF-up gene expression signature with ERBB2-positive and poor-outcome breast tumors suggests a role for ER α in these tumors and with endocrine therapy-resistant breast cancers dependent on growth factor pathway activation.

Stimulus-specific ER α cistrome leads to unique transcriptional program

To address whether differential ER α recruitment to the genome mediates the stimulus-specific transcriptional responses under growth factor stimulation, we compared EGF and estrogen-induced ER α cistromes from MCF7 breast cancer cells (Fig. 2A; Supplemental Figs. S3–S5; Carroll et al. 2005, 2006; Lupien et al. 2008). Although both estrogen and EGF induced ER α recruitment to shared sites (Fig. 2A), growth factor pathway activation induced ER α recruitment to a significant number of unique sites (Fig. 2A; Supplemental Fig. S5A). In keeping with the central role of FoxA1 in ER α -positive breast cancer, the Forkhead (FKH) motif was highly enriched in both the estrogen and EGF-induced ER α cistromes (Fig. 2B). In fact, >45% of the EGF-unique ER α cistromes overlapped with the previously characterized FoxA1 cistromes in MCF7 cells (Supplemental Fig. S6; Lupien et al. 2008). However, the EGF-unique sites were more highly enriched for the AP-1 as opposed to estrogen-responsive element (ERE) motif (Fig. 2B). This suggests that recruitment of ER α following growth factor stimulation is occurring preferentially through an indirect tethering mechanism involving AP-1 family members. Chromatin immunoprecipitation (ChIP)–reChIP assays directed against ER α and AP-1 demonstrate that these two transcription factors are corecruited upon EGF stimulation to ER α -binding sites (Fig. 2C). Proliferation assays in the presence of a dominant-negative AP-1 mutant (TAM67) (Dhar et al. 2004) reveal the central role of AP-1 in EGF-mediated MCF7 breast cancer cell proliferation and in ER α -dependent EGF target gene regulation (Fig. 2D,E). Taken together, these results suggest that AP-1 is a critical partner in ER α signaling favorable to the growth of breast cancer cells under growth factor stimulation.

Of interest, EGF up-regulated genes were significantly associated with EGF-unique but not estrogen-unique ER α -binding sites (Fig. 3A). In fact, the estrogen-unique ER α -binding sites did not associate with the estrogen-responsive genes (Fig. 3A). This is in agreement with their poor overlap with FoxA1-binding regions (Supplemental Fig. S6), which we previously showed is typical of sites not driving an estrogen response in breast cancer cells (Lupien et al. 2008). The role of EGF-unique ER α -binding sites on the EGF-induced transcriptional response is exemplified by analyzing ER α recruitment in MCF7 breast cancer cells near the EGF-responsive *TNFRSF21* and *LIF* genes (Fig.

Figure 2. AP-1 is central to the growth factor-induced ER α cistromes. (A) Genome-wide ChIP-on-chip analysis following EGF stimulation in MCF7 breast cancer cells reveals 12,550 ER α -binding sites (false discovery rate [FDR] 1%), 31% overlapping with the estrogen (E2)-induced ER α cistrome. (B) Sequence analysis of EGF-unique (orange), shared (purple), or E2-unique (green) ER α -binding sites reveals the preferential enrichment of EREs in the center of the E2-unique and shared binding sites, while the FKH and AP-1 motifs are preferentially enriched in the center of the shared and EGF-unique ER α cistromes. (C) ChIP–reChIP assays directed against ER α and AP-1 were performed to reveal the corecruitment of these factors following EGF stimulation on ER α -binding sites. (*) $P \leq 0.05$; (**) $P \leq 0.01$; (***) $P \leq 0.001$. (D) Proliferation of MCF7 breast cancer cells transfected with the mock (pcDNA3.1) or AP-1 dominant-negative (TAM67) vectors was measured following EGF stimulation (*) $P \leq 0.05$; (**) $P \leq 0.01$; (***) $P \leq 0.001$. (E) mRNA levels of EGF target genes were measured following EGF stimulation in MCF7 breast cancer cells transfected with the mock (pcDNA3.1) or AP-1 dominant-negative (TAM67) vectors. (*) $P \leq 0.05$; (**) $P \leq 0.01$; (***) $P \leq 0.001$.

3B,C; Supplemental Fig. S7A). Indeed, ER α was recruited to all three regulatory elements associated with *TNFRSF21* and *LIF* following EGF stimulation, while estrogen treatment induced only ER α recruitment to *TNFRSF21* enh1 and *LIF* enh2 (Fig. 3B; Supplemental Fig. S7A). These results suggest that the stimuli-specific transcriptional response in a given cell is in part dependent on a unique ER α cistrome. Furthermore, the increased number of ER α -binding sites following EGF stimulation near EGF target genes (Supplemental Fig. S7A) is in agreement with the notion that genes are more likely regulated when ER α -binding sites cluster near them (Krum et al. 2008).

Activation of ER α by growth factor pathway stimulation has been shown to be dependent on its phosphorylation at specific N-terminal residues, including, predominantly,

Figure 3. Growth factor-induced transcriptional response relates to stimuli-specific ER α cistromes. (A) Correlation between E2, EGF-up, or ER α -dependent EGF-up target genes with EGF-unique, shared, or E2-unique ER α -binding sites from MCF7 breast cancer cells. The occurrence of ER α -binding sites within 20 kb of the TSS of regulated genes was compared with that on nonregulated genes. (*) $P \leq 0.05$; (**) $P \leq 0.01$; (***) $P \leq 0.001$. (B) ChIP-qPCR results against ER α performed under EGF or E2 stimulation in MCF7 cells on the regulatory element associated with the *TNFRSF21* and *LIF* genes. (C) RT-qPCR results measuring expression of the EGF-specific responsive genes *TNFRSF21* and *LIF* following 3 h of E2 or EGF stimulation.

Ser 118 (S118phos) (Kato et al. 1995; Bunone et al. 1996; Joel et al. 1998; Smith 1998; Kato 2001; Kurokawa and Arteaga 2003). In contrast, estrogen-mediated activation of ER α is dependent on an activation domain overlapping its ligand-binding domain (LBD) located in its C-terminal region that also leads to S118 phosphorylation (Chen et al. 2002). We found that recruitment of S118-phosphorylated ER α to the genome was detected following EGF stimulation at EGF-unique as well as EGF-estrogen shared ER α -binding sites (Supplemental Fig. S7B,C). This suggests that ER α phosphorylation alone is not sufficient to explain its stimulus-specific cistromes.

Stimulus-specific ER α coactivation

We demonstrated previously that FoxA1-binding sites shared across cells of different lineages can exhibit cell type-specific activities (Eeckhoutte et al. 2009). Here, although a significant proportion of the EGF-induced ER α cistrome is unique, close to 4000 sites are shared with the estrogen-induced ER α cistrome (Fig. 2A). Furthermore, EGF as well as estrogen-regulated genes are enriched around these shared binding sites (Fig. 3A). Hence, this suggests that different subsets of the shared ER α -binding sites are active under EGF or estrogen stimulation. We demonstrated recently that only a fraction of the ER α cistrome undergoes coactivation and associates with regulated genes following estrogen stimulation in breast cancer cells (Lupien et al. 2009). Hence, a number of estrogen-induced ER α -binding events appear to be futile for the establishment of the estrogen-induced transcriptional response. Furthermore, coactivators were reported previously to undergo specific regulation following growth factor stimulation (Font De Mora and Brown 2000; Lopez et al. 2001). We therefore addressed the contribution of stimuli-specific coactivation of the shared ER α -binding sites in the distinct transcriptional response generated in MCF7 breast cancer cells following EGF or estrogen treatment. In order to address

coactivation instead of coactivator recruitment, we decided to measure a post-translational modification induced by CBP/p300, a well-established ER α coactivator (Hanstein et al. 1996). This was accomplished by measuring the level of histone H3 acetylation on Lys 18 (H3K18ac) following either EGF or estrogen treatment in MCF7 breast cancer cells. Comparison of H3K18ac levels with those in control untreated cells revealed the stimuli-specific coactivation of shared ER α -binding sites (Fig. 4A). Indeed, a fraction of shared ER α -binding sites associated with strong induction of H3K18ac following EGF treatment, while a distinct fraction associated with the estrogen-induced H3K18ac (Fig. 4A). Importantly, EGF-regulated genes were specifically enriched near shared ER α -binding sites preferentially coactivated under that same treatment (Fig. 4B). Conversely, estrogen-regulated genes were significantly enriched near shared ER α -binding sites specifically coactivated following estrogen treatment (Fig. 4B). This was exemplified by the *ACP6* and *LIF* EGF-regulated genes and the *TFF1* and *XBPI* estrogen-regulated genes (Fig. 4C). Both EGF and estrogen could induce ER α recruitment to the series of regulatory elements near these genes in MCF7 breast cancer cells (Fig. 4D; Supplemental Fig. S8); however, their coactivation was stimuli-specific. Indeed, histone acetylation (both H3K18ac and H4K12ac) could be induced only on the ER α -binding sites near *ACP6* and *LIF* following EGF treatment (Fig. 4D; Supplemental Fig. S8). Similarly, estrogen could induce histone acetylation only on *TFF1*- and *XBPI*-associated ER α regulatory elements (Fig. 4D; Supplemental Fig. S8). Hence, this suggests that growth factor pathway activation leads to the coactivation of a different set of ER α -binding sites shared with other stimuli.

ER α -dependent growth factor response relates to poor-outcome breast cancers

To address the relevance of growth factor-dependent ER α cistrome in breast tumors, we established the correlation

Figure 4. Selective coactivation of ER α -binding sites under growth factor stimulation. (A) Coactivation as measured by H3K18ac levels established following either EGF or estrogen (E2) stimulation by ChIP-on-chip of chromosomes 8, 11, and 12 in MCF7 breast cancer cells. The MAT score for H3K18ac (EGF/control and E2/control) was established on all ER α -binding sites shared under E2 and EGF stimulation found on chromosomes 8, 11, and 12. K-means cluster analysis was used to identify ER α -binding sites preferentially associated with induced H3K18ac under E2 or EGF treatment. (B) Correlation between gene expression and the three subsets of shared ER α -binding sites based on H3K18ac levels was performed as described for Figure 2D. (C) Specific examples validating the concept of stimulus-specific coactivation at ER α -binding sites common to estrogen and EGF stimulation. RT-qPCR studies validate the stimuli-specific expression of *ACP6*, *LIF*, *TFF1*, and *XBP1* in MCF7 breast cancer cells following E2 or EGF treatment in MCF7 cells. (D) ChIP-qPCR analysis demonstrates E2 and EGF equivalent induction of ER α recruitment to regulatory regions associated with *ACP6*, *LIF*, *TFF1*, and *XBP1*. Coactivation measured through acetylation of H4K12ac or H3K18ac (Supplemental Material) is specific to ER α -binding sites near genes induced under the same treatment in MCF7 cells.

between the genes with growth factor pathway-specific ER α -binding sites within 20 kb of their transcription start sites (TSSs) and breast tumor expression signatures using OncoPrint Concepts Map analysis. Significant correlations were revealed between genes associated with EGF-ER α -specific sites and gene expression signatures found in poor-outcome (relapse, death, metastasis, and high tumor grade) as well as ERBB2-positive breast tumors (Fig. 5A). Genes with estrogen-associated ER α -binding sites within 20 kb of their TSSs were not associated with these gene expression signatures, but were associated with ER α -positive overexpressed gene signatures (Supplemental Fig. S9). Therefore, the growth factor pathway-specific ER α cistrome supports its role in the transcriptional response associated with breast tumors overexpressing ERBB2 and with poor outcomes.

Discussion

Taken together, our results demonstrate that differential recruitment and coactivation of ER α is a fundamental mechanism allowing for stimulus-specific transcriptional programs (Fig. 5B). Furthermore, as growth factor pathway activation is commonly associated with the development of hormone-refractory breast tumors (Dowsett

2001), our results suggest that ER α can play a fundamental role in their proliferation and involves the transcription factor AP-1. Indeed, hormone-refractory tumors are typically dependent on the overexpression of the EGFR or ERBB2 (Benz et al. 1992; Pietras et al. 1995; Kurokawa et al. 2000; Nicholson et al. 2001; Shou et al. 2004). In addition, a subset of hormone-refractory breast tumors are responsive to the full ER α antagonist fulvestrant (Howell et al. 1995; Martin et al. 2005). Furthermore, fulvestrant results in clinical benefits in ERBB2-overexpressing advanced breast cancers (Robertson et al. 2010). Additionally, in model systems, treatment of breast cancer cells with the combination of fulvestrant and growth factor pathway inhibitors more significantly represses growth than either treatment alone, and prevents the development of endocrine resistance (Kunisue et al. 2000; Gee et al. 2003; Pietras et al. 2003; Macedo et al. 2008). Considering that ER α - and ERBB2-positive breast cancers are resistant to endocrine therapies targeting estrogen stimulation of ER α , such as aromatase inhibitors or selective ER modulators, our results provide a mechanistic understanding for this clinical observation. Our results suggest that complete ER α antagonists that would block both its estrogen- and growth factor-stimulated activities would overcome this problem. Hence, the EGF-induced ER α cistrome reveals

Figure 5. Growth factor ERα cistrome relates to poor-outcome expression signatures in breast tumors. (A) OncoPrint Concepts Map analysis reveals significant association between genes specifically associated with an EGF-specific ERα-binding site within 20 kb of their TSS and gene expression signatures from ERBB2-positive, poor-outcome (metastasis, recurrence, death, and high grade), or ERα-positive breast tumors characterized in eight independent studies (each represented by individual circles). Only significantly associated gene lists are linked by a straight line ($P \leq 1e-4$, O.R. ≥ 1.35). The gene list associated with estrogen (E2)-induced ERα-binding sites within 20 kb of their TSS does not significantly associate with any of the gene expression signatures linked to EGF-ERα-associated genes. The right panel presents an example of the expression profile for EGF-ERα-associated genes differentially expressed in ERBB2-positive breast tumors according to one independent study. Only genes significantly differentially expressed in ERBB2-positive versus -negative breast tumors are presented ($P \leq 5e-2$). (B) Schematic representation of mechanisms involved in stimuli-specific transcriptional response acting through a shared transcription factor. E2 and the growth factor pathway (GF) induce a unique transcriptional response dependent on the ERα. This involves stimuli-specific ERα cistromes and coactivation. Current therapies for ERα-positive primary breast tumors, including aromatase inhibitors (AI) or selective ERα modulators such as tamoxifen (T), block only E2-mediated activation of the receptor. Full antiestrogens (FA) such as fulvestrant lead to ERα degradation and can therefore block E2- as well as GF-mediated activation of ERα. Growth factor inhibitors (GFI) should also block ERα activation following growth factor pathway activation.

key features to be considered in the development of therapeutic strategies for hormone-refractory ERα-positive breast tumors.

Materials and methods

ChIP-on-chip and ChIP-qPCR and ChIP-reChIP-qPCR

Prior to stimulation, MCF7 cells were maintained for 3 d in phenol red-free medium (Invitrogen) supplemented with 10% charcoal dextran-treated fetal bovine serum (CDT-FBS) as described previously (Lupien et al. 2008). Cells were stimulated with the EGF (100 ng/mL) for 90 min and crosslinked using 1% formaldehyde (Kato et al. 1995; Cunliffe et al. 2003). Samples were sonicated (Fisher Sonic Dismembrator, model 500) and immunoprecipitated as described previously (Carroll et al. 2005, 2006) using antibodies against ERα (Santa Cruz Biotechnology, Inc., HC-20; Neomarkers, Ab-10), and H3K18ac (Millipore, 07-354). Three independent assays were performed. Purified samples were labeled and hybridized to microarrays (Affymetrix GeneChip Human Tiling 2.0R array sets). Genome-wide ChIP-on-Chip analysis was conducted using the model-based analysis of tiling arrays program (MAT) based on the latest human genomic sequence (Hg18) (Johnson et al. 2006). All ChIP-on-chip data used in this study can be accessed at <http://research.dfci.harvard.edu/brownlab/datasets>. ChIP-qPCR experiments were

performed as in Carroll et al. (2005). Antibodies against ERα (Santa Cruz Biotechnology, Inc., HC-20; Neomarkers, Ab-10); ERα S118P (Millipore, 07-487), H3K18ac (Upstate Biotechnologies, Inc., 07-354), and H4K12ac (Upstate Biotechnologies, Inc., 07-595) were used for this assay. ChIP-reChIP was performed as described previously (Ross-Innes et al. 2010). AP-1 was re-ChIPed using a mix of anti-AP-1 antibodies (Santa Cruz Biotechnology, Inc., SC-44 and SC-253). Statistically significant differences were established using a Student's *t*-test comparison for unpaired data. Primer sequences used in this assay are found in Supplemental Table 1.

Gene expression profiling

Prior to stimulation, MCF7 cells were maintained for 3 d in phenol red-free medium (Invitrogen) supplemented with 10% CDT-FBS as described previously (Lupien et al. 2008). Cells were pretreated with fulvestrant (100 nM; ICI182,780) or control vehicle for 3 h and then stimulated with the EGF (100 μg/mL) for 3 h before RNA extraction using Qiagen RNeasy kit (Qiagen). Triplicate experiments were performed using Affymetrix U133Plus2.0 expression microarrays. The Robust Multichip Average (RMA) algorithm was used to analyze the data as described previously (Carroll et al. 2006), and level of differential expression for each time point relative to 0 h was established as in Lupien et al. (2008). Gene lists are found in Supplemental Table 2A–E. Statistically differentially expressed genes (*t*-test, $P \geq 1e-3$) were

defined as EGF-up genes. ER α -dependent EGF-up genes correspond to the EGF-up genes that were no longer significantly expressed when cells were treated with fulvestrant prior to EGF stimulation (EGF-responsive genes, t -test $P \geq 1e-3$, minus fulvestrant + EGF-responsive genes, t -test, $P \geq 1e-2$). Estrogen-responsive genes were presented previously (Carroll et al. 2006).

Transfection of MCF7 cells

MCF7 cells were maintained in phenol red-free medium (Invitrogen) supplemented with 10% CDT-FBS as described previously (Lupien et al. 2008) prior to transfection. MCF7 cells were transfected with the mock (pcDNA3.1) or AP-1 dominant-negative (TAM67) vectors (1 μ g per well) using Lipofectamine 2000 DNA transfection reagent according to the manufacturer's instructions (SigmaGen). Forty-eight hours after transfection, cells were stimulated with control (ddH₂O) or the EGF (100 ng/mL). For cell proliferation assays, cell number was determined every 24 h after EGF addition. For expression assays, RNA was extracted 3 h following EGF stimulation.

RT-qPCR

Collected RNA was processed for RT-qPCR as described previously (Krum et al. 2008). Primer sequences used for in RT-qPCR are listed in Supplemental Table 1.

Sequence analysis and cluster analysis

Genome-wide distribution as well as sequence conservation analysis of the different clusters derived from the ER α estrogen versus EGF cistromes was determined using the Cis-Elements Annotation Systems (CEAS) (Ji et al. 2006). Enriched motifs within clusters as well as the association of trends in gene expression with cluster binding sites were identified as described in Lupien et al. (2008).

Oncomine Concepts Map

We compared our various gene lists (list of genes in each gene list can be found in Supplemental Table 3A,B) with expression profiles from breast tumors compiled on Oncomine (Compendia Bioscience; <http://www.oncomine.org>). Using the Oncomine Concepts Map tools, we established significant association between our gene lists and Oncomine overexpressed or under-expressed Gene Expression Signature derived from independent breast cancer studies. Node connection figures can be generated with Cytoscape (<http://www.cytoscape.org>). Gene Expression Signatures used in Figures 1D and 4A and Supplemental Figure S2 are derived from van de Vijver et al. (2002), Zhao et al. (2004), Miller et al. (2005), Minn et al. (2005), Wang et al. (2005), Chin et al. (2006), Ginestier et al. (2006), Hess et al. (2006), Ivshina et al. (2006), Richardson et al. (2006), Sotiriou et al. (2006), Yu et al. (2006), Desmedt et al. (2007), Saal et al. (2007), Boersma et al. (2008), and Finak et al. (2008).

Acknowledgments

We thank Dr. Nancy Colburn (NCI-Frederick) for providing the dominant-negative AP-1 construct (TAM67). This work was supported by grants from the NIDDK (R01DK074967 to M.B.), the NCI (P01 CA8011105, and the DF/HCC Breast Cancer SPORE Grant P50C89393 to M.B.), the DFCI Women's Cancers Program, and the US Department of Defense Breast Cancer Research Program Awards (W81XWH-08-1-0214 to M.L.).

References

- Benz CC, Scott GK, Sarup JC, Johnson RM, Tripathy D, Coronado E, Shepard HM, Osborne CK. 1992. Estrogen-dependent, tamoxifen-resistant tumorigenic growth of MCF-7 cells transfected with HER2/neu. *Breast Cancer Res Treat* **24**: 85–95.
- Boersma BJ, Reimers M, Yi M, Ludwig JA, Luke BT, Stephens RM, Yfantis HG, Lee DH, Weinstein JN, Ambs S. 2008. A stromal gene signature associated with inflammatory breast cancer. *Int J Cancer* **122**: 1324–1332.
- Bunone G, Briand PA, Miksicek RJ, Picard D. 1996. Activation of the unliganded estrogen receptor by EGF involves the MAP kinase pathway and direct phosphorylation. *EMBO J* **15**: 2174–2183.
- Carroll JS, Liu XS, Brodsky AS, Li W, Meyer CA, Szary AJ, Eeckhoutte J, Shao W, Hestermann EV, Geistlinger TR, et al. 2005. Chromosome-wide mapping of estrogen receptor binding reveals long-range regulation requiring the forkhead protein FoxA1. *Cell* **122**: 33–43.
- Carroll JS, Meyer CA, Song J, Li W, Geistlinger TR, Eeckhoutte J, Brodsky AS, Keeton EK, Fertuck KC, Hall GF, et al. 2006. Genome-wide analysis of estrogen receptor binding sites. *Nat Genet* **38**: 1289–1297.
- Chen D, Washbrook E, Sarwar N, Bates GJ, Pace PE, Thirunuvakkarasu V, Taylor J, Epstein RJ, Fuller-Pace FV, Egly J-M, et al. 2002. Phosphorylation of human estrogen receptor α at serine 118 by two distinct signal transduction pathways revealed by phosphorylation-specific antisera. *Oncogene* **21**: 4921–4931.
- Chin K, DeVries S, Fridlyand J, Spellman PT, Roydasgupta R, Kuo WL, Lapuk A, Neve RM, Qian Z, Ryder T, et al. 2006. Genomic and transcriptional aberrations linked to breast cancer pathophysiology. *Cancer Cell* **10**: 529–541.
- Cunliffe HE, Ringner M, Bilke S, Walker RL, Cheung JM, Chen Y, Meltzer PS. 2003. The gene expression response of breast cancer to growth regulators: Patterns and correlation with tumor expression profiles. *Cancer Res* **63**: 7158–7166.
- Curtis SW, Washburn T, Sewall C, DiAugustine R, Lindzey J, Couse JF, Korach KS. 1996. Physiological coupling of growth factor and steroid receptor signaling pathways: Estrogen receptor knockout mice lack estrogen-like response to epidermal growth factor. *Proc Natl Acad Sci* **93**: 12626–12630.
- Desmedt C, Piette F, Loi S, Wang Y, Lallemand F, Haibe-Kains B, Viale G, Delorenzi M, Zhang Y, d'Assignies MS, et al. 2007. Strong time dependence of the 76-gene prognostic signature for node-negative breast cancer patients in the TRANSBIG multicenter independent validation series. *Clin Cancer Res* **13**: 3207–3214.
- Dhar A, Hu J, Reeves R, Resar LM, Colburn NH. 2004. Dominant-negative c-Jun (TAM67) target genes: HMGA1 is required for tumor promoter-induced transformation. *Oncogene* **27**: 4466–4476.
- Dowsett M. 2001. Overexpression of HER-2 as a resistance mechanism to hormonal therapy for breast cancer. *Endocr Relat Cancer* **8**: 191–195.
- Dudek P, Picard D. 2008. Genomics of signaling crosstalk of estrogen receptor α in breast cancer cells. *PLoS ONE* **3**: e1859. doi: 10.1371/journal.pone.0001859.
- Eeckhoutte J, Lupien M, Meyer CA, Verzi MP, Shivdasani RA, Liu XS, Brown M. 2009. Cell-type selective chromatin remodeling defines the active subset of FOXA1-bound enhancers. *Genome Res* **19**: 372–380.
- Finak G, Bertos N, Pepin F, Sadekova S, Souleimanova M, Zhao H, Chen H, Omeroglu G, Meterissian S, Omeroglu A, et al.

2008. Stromal gene expression predicts clinical outcome in breast cancer. *Nat Med* **14**: 518–527.
- Font De Mora J, Brown M. 2000. AIB1 is a conduit for kinase-mediated growth factor signaling to the estrogen receptor. *Mol Cell Biol* **20**: 5041–5047.
- Fullwood MJ, Liu MH, Pan YF, Liu J, Xu H, Mohamed YB, Orlov YL, Velkov S, Ho A, Mei PH, et al. 2009. An oestrogen-receptor- α -bound human chromatin interactome. *Nature* **462**: 58–64.
- Gee JM, Harper ME, Hutcheson IR, Madden TA, Barrow D, Knowlden JM, McClelland RA, Jordan N, Wakeling AE, Nicholson RI. 2003. The antiepidermal growth factor receptor agent gefitinib (ZD1839/Iressa) improves antihormone response and prevents development of resistance in breast cancer in vitro. *Endocrinology* **144**: 5105–5117.
- Ginestier C, Cervera N, Finetti P, Esteyries S, Esterni B, Adelaide J, Xerri L, Viens P, Jacquemier J, Charafe-Jauffret E, et al. 2006. Prognosis and gene expression profiling of 20q13-amplified breast cancers. *Clin Cancer Res* **12**: 4533–4544.
- Hanstein B, Eckner R, DiRenzo J, Halachmi S, Liu H, Searcy B, Kurokawa R, Brown M. 1996. p300 is a component of an estrogen receptor coactivator complex. *Proc Natl Acad Sci* **93**: 11540–11545.
- Hess KR, Anderson K, Symmans WF, Valero V, Ibrahim N, Mejia JA, Booser D, Theriault RL, Buzdar AU, Dempsey PJ, et al. 2006. Pharmacogenomic predictor of sensitivity to preoperative chemotherapy with paclitaxel and fluorouracil, doxorubicin, and cyclophosphamide in breast cancer. *J Clin Oncol* **24**: 4236–4244.
- Howell A, DeFriend D, Robertson J, Blamey R, Walton P. 1995. Response to a specific antioestrogen (ICI 182780) in tamoxifen-resistant breast cancer. *Lancet* **345**: 29–30.
- Hua S, Kallen CB, Dhar R, Baquero MT, Mason CE, Russell BA, Shah PK, Liu J, Khramtsov A, Tretiakova MS, et al. 2008. Genomic analysis of estrogen cascade reveals histone variant H2A.Z associated with breast cancer progression. *Mol Syst Biol* **4**: 188. doi: 10.1038/msb.2008.25.
- Hurtado A, Holmes KA, Geistlinger TR, Hutcheson IR, Nicholson RI, Brown M, Jiang J, Howat WJ, Ali S, Carroll JS. 2008. Regulation of ERBB2 by oestrogen receptor-PAX2 determines response to tamoxifen. *Nature* **456**: 663–666.
- Ignar-Trowbridge DM, Nelson KG, Bidwell MC, Curtis SW, Washburn TE, McLachlan JA, Korach KS. 1992. Coupling of dual signaling pathways: Epidermal growth factor action involves the estrogen receptor. *Proc Natl Acad Sci* **89**: 4658–4662.
- Ivshina AV, George J, Senko O, Mow B, Putti TC, Smeds J, Lindahl T, Pawitan Y, Hall P, Nordgren H, et al. 2006. Genetic reclassification of histologic grade delineates new clinical subtypes of breast cancer. *Cancer Res* **66**: 10292–10301.
- Ji X, Li W, Song J, Wei L, Liu XS. 2006. CEAS: Cis-regulatory element annotation system. *Nucleic Acids Res* **34**: W551–W554. doi: 10.1093/nar/gkl322.
- Joel PB, Smith J, Sturgill TW, Fisher TL, Blenis J, Lannigan DA. 1998. pp90rsk1 regulates estrogen receptor-mediated transcription through phosphorylation of Ser-167. *Mol Cell Biol* **18**: 1978–1984.
- Johnson WE, Li W, Meyer CA, Gottardo R, Carroll JS, Brown M, Liu XS. 2006. Model-based analysis of tiling-arrays for ChIP-chip. *Proc Natl Acad Sci* **103**: 12457–12462.
- Kato S. 2001. Estrogen receptor-mediated cross-talk with growth factor signaling pathways. *Breast Cancer* **8**: 3–9.
- Kato S, Endoh H, Masuhiro Y, Kitamoto T, Uchiyama S, Sasaki H, Masushige S, Gotoh Y, Nishida E, Kawashima H, et al. 1995. Activation of the estrogen receptor through phosphorylation by mitogen-activated protein kinase. *Science* **270**: 1491–1494.
- Knowlden JM, Hutcheson IR, Jones HE, Madden T, Gee JM, Harper ME, Barrow D, Wakeling AE, Nicholson RI. 2003. Elevated levels of epidermal growth factor receptor/c-erbB2 heterodimers mediate an autocrine growth regulatory pathway in tamoxifen-resistant MCF-7 cells. *Endocrinology* **144**: 1032–1044.
- Krum SA, Miranda-Carboni GA, Lupien M, Eeckhoute J, Carroll JS, Brown M. 2008. Unique ER α cistromes control cell type-specific gene regulation. *Mol Endocrinol* **22**: 2393–2406.
- Kunisue H, Kurebayashi J, Otsuki T, Tang CK, Kurosumi M, Yamamoto S, Tanaka K, Doihara H, Shimizu N, Sonoo H. 2000. Anti-HER2 antibody enhances the growth inhibitory effect of anti-oestrogen on breast cancer cells expressing both estrogen receptors and HER2. *Br J Cancer* **82**: 46–51.
- Kurokawa H, Arteaga CL. 2003. ErbB (HER) receptors can abrogate antiestrogen action in human breast cancer by multiple signaling mechanisms. *Clin Cancer Res* **9**: 511S–515S.
- Kurokawa H, Lenferink AE, Simpson JE, Pisacane PI, Sliwkowski MX, Forbes JT, Arteaga CL. 2000. Inhibition of HER2/neu (erbB-2) and mitogen-activated protein kinases enhances tamoxifen action against HER2-overexpressing, tamoxifen-resistant breast cancer cells. *Cancer Res* **60**: 5887–5894.
- Lee AV, Yee D. 1995. Insulin-like growth factors and breast cancer. *Biomed Pharmacother* **49**: 415–421.
- Lin CY, Vega VB, Thomsen JS, Zhang T, Kong SL, Xie M, Chiu KP, Lipovich L, Barnett DH, Stossi F, et al. 2007. Whole-genome cartography of estrogen receptor α binding sites. *PLoS Genet* **3**: e87. doi: 10.1371/journal.pgen.0030087.
- Liu Y, Gao H, Marstrand TT, Strom A, Valen E, Sandelin A, Gustafsson JA, Dahlman-Wright K. 2008. The genome landscape of ER α - and ER β -binding DNA regions. *Proc Natl Acad Sci* **105**: 2604–2609.
- Lopez GN, Turck CW, Schaufele F, Stallcup MR, Kushner PJ. 2001. Growth factors signal to steroid receptors through mitogen-activated protein kinase regulation of p160 coactivator activity. *J Biol Chem* **276**: 22177–22182.
- Lupien M, Jeyakumar M, Hebert E, Hilmi K, Cotnoir-White D, Loch C, Auger A, Dayan G, Pinard GA, Wurtz JM, et al. 2007. Raloxifene and ICI182,780 increase estrogen receptor- α association with a nuclear compartment via overlapping sets of hydrophobic amino acids in activation function 2 helix 12. *Mol Endocrinol* **21**: 797–816.
- Lupien M, Eeckhoute J, Meyer CA, Wang Q, Zhang Y, Li W, Carroll JS, Liu XS, Brown M. 2008. FoxA1 translates epigenetic signatures into enhancer-driven lineage-specific transcription. *Cell* **132**: 958–970.
- Lupien M, Eeckhoute J, Meyer CA, Krum SA, Rhodes DR, Liu XS, Brown M. 2009. Coactivator function defines the active estrogen receptor- α cistrome. *Mol Cell Biol* **29**: 3413–3423.
- Lupu R, Cardillo M, Cho C, Harris L, Hijazi M, Perez C, Rosenberg K, Yang D, Tang C. 1996. The significance of heregulin in breast cancer tumor progression and drug resistance. *Breast Cancer Res Treat* **38**: 57–66.
- Macedo LF, Sabnis GJ, Golubeva OG, Brodie A. 2008. Combination of anastrozole with fulvestrant in the intratumoral aromatase xenograft model. *Cancer Res* **68**: 3516–3522.
- Martin LA, Pancholi S, Chan CM, Farmer I, Kimberley C, Dowsett M, Johnston SR. 2005. The anti-oestrogen ICI 182,780, but not tamoxifen, inhibits the growth of MCF-7 breast cancer cells refractory to long-term oestrogen deprivation through down-regulation of oestrogen receptor and IGF signalling. *Endocr Relat Cancer* **12**: 1017–1036.

- Miller LD, Smeds J, George J, Vega VB, Vergara L, Ploner A, Pawitan Y, Hall P, Klaar S, Liu ET, et al. 2005. An expression signature for p53 status in human breast cancer predicts mutation status, transcriptional effects, and patient survival. *Proc Natl Acad Sci* **102**: 13550–13555.
- Minn AJ, Gupta GP, Siegel PM, Bos PD, Shu W, Giri DD, Viale A, Olshen AB, Gerald WL, Massague J. 2005. Genes that mediate breast cancer metastasis to lung. *Nature* **436**: 518–524.
- Moasser MM. 2007. The oncogene HER2: Its signaling and transforming functions and its role in human cancer pathogenesis. *Oncogene* **26**: 6469–6487.
- Monroe DG, Getz BJ, Johnsen SA, Riggs BL, Khosla S, Spelsberg TC. 2003. Estrogen receptor isoform-specific regulation of endogenous gene expression in human osteoblastic cell lines expressing either ER α or ER β . *J Cell Biochem* **90**: 315–326.
- Nicholson RI, Hutcheson IR, Harper ME, Knowlden JM, Barrow D, McClelland RA, Jones HE, Wakeling AE, Gee JM. 2001. Modulation of epidermal growth factor receptor in endocrine-resistant, oestrogen receptor-positive breast cancer. *Endocr Relat Cancer* **8**: 175–182.
- Pietras RJ, Arboleda J, Reese DM, Wongvipat N, Pegram MD, Ramos L, Gorman CM, Parker MG, Sliwkowski MX, Slamon DJ. 1995. HER-2 tyrosine kinase pathway targets estrogen receptor and promotes hormone-independent growth in human breast cancer cells. *Oncogene* **10**: 2435–2446.
- Pietras RJ, Marquez DC, Chen HW, Ayala R, Ramos LB, Slamon DJ. 2003. Improved antitumor therapy with Herceptin and Faslodex for dual targeting of HER-2 and estrogen receptor signalling pathways in human breast cancers with overexpression of HER-2/neu gene. *Breast Cancer Res Treat* **82**: S12, Abstract 22.
- Rhodes DR, Kalyana-Sundaram S, Tomlins SA, Mahavisno V, Kasper N, Varambally R, Barrette TR, Ghosh D, Varambally S, Chinnaiyan AM. 2007. Molecular concepts analysis links tumors, pathways, mechanisms, and drugs. *Neoplasia* **9**: 443–454.
- Richardson AL, Wang ZC, De Nicolo A, Lu X, Brown M, Miron A, Liao X, Iglehart JD, Livingston DM, Ganesan S. 2006. X chromosomal abnormalities in basal-like human breast cancer. *Cancer Cell* **9**: 121–132.
- Robertson JFR, Steger GG, Neven P, Barni S, Gieseking F, Nole F, Pritchard KI, O'Malley FP, Simon SD, Kaufman B, et al. 2010. Activity of fulvestrant in HER2-overexpressing advanced breast cancer. *Ann Oncol* **21**: 1246–1253.
- Ross-Innes CS, Stark R, Holmes KA, Schmidt D, Spyrou C, Russel R, Massie CE, Vowler SL, Eldridge M, Carroll JS. 2010. Cooperative interaction between retinoic acid receptor- α and estrogen receptor in breast cancer. *Genes Dev* **24**: 171–182.
- Saal LH, Johansson P, Holm K, Gruvberger-Saal SK, She QB, Maurer M, Koujak S, Ferrando AA, Malmstrom P, Memeo L, et al. 2007. Poor prognosis in carcinoma is associated with a gene expression signature of aberrant PTEN tumor suppressor pathway activity. *Proc Natl Acad Sci* **104**: 7564–7569.
- Schiff R, Massarweh SA, Shou J, Bharwani L, Arpino G, Rimawi M, Osborne CK. 2005. Advanced concepts in estrogen receptor biology and breast cancer endocrine resistance: Implicated role of growth factor signaling and estrogen receptor coregulators. *Cancer Chemother Pharmacol* **56**: 10–20.
- Shou J, Massarweh S, Osborne CK, Wakeling AE, Ali S, Weiss H, Schiff R. 2004. Mechanisms of tamoxifen resistance: Increased estrogen receptor-HER2/neu cross-talk in ER/HER2-positive breast cancer. *J Natl Cancer Inst* **96**: 926–935.
- Smith CL. 1998. Cross-talk between peptide growth factor and estrogen receptor signaling pathways. *Biol Reprod* **58**: 627–632.
- Sorlie T, Perou CM, Tibshirani R, Aas T, Geisler S, Johnsen H, Hastie T, Eisen MB, van de Rijn M, Jeffrey SS, et al. 2001. Gene expression patterns of breast carcinomas distinguish tumor subclasses with clinical implications. *Proc Natl Acad Sci* **98**: 10869–10874.
- Sorlie T, Tibshirani R, Parker J, Hastie T, Marron JS, Nobel A, Deng S, Johnsen H, Pesich R, Geisler S, et al. 2003. Repeated observation of breast tumor subtypes in independent gene expression data sets. *Proc Natl Acad Sci* **100**: 8418–8423.
- Sotiriou C, Wirapati P, Loi S, Harris A, Fox S, Smeds J, Nordgren H, Farmer P, Praz V, Haibe-Kains B, et al. 2006. Gene expression profiling in breast cancer: Understanding the molecular basis of histologic grade to improve prognosis. *J Natl Cancer Inst* **98**: 262–272.
- van de Vijver MJ, He YD, van't Veer LJ, Dai H, Hart AA, Voskuil DW, Schreiber GJ, Peterse JL, Roberts C, Marton MJ, et al. 2002. A gene-expression signature as a predictor of survival in breast cancer. *N Engl J Med* **347**: 1999–2009.
- Visel A, Blow MJ, Li Z, Zhang T, Akiyama JA, Holt A, Plajzer-Frick I, Shoukry M, Wright C, Chen F, et al. 2009. ChIP-seq accurately predicts tissue-specific activity of enhancers. *Nature* **457**: 854–858.
- Wang Y, Klijn JG, Zhang Y, Sieuwerts AM, Look MP, Yang F, Talantov D, Timmermans M, Meijer-van Gelder ME, Yu J, et al. 2005. Gene-expression profiles to predict distant metastasis of lymph-node-negative primary breast cancer. *Lancet* **365**: 671–679.
- Yu K, Ganesan K, Miller LD, Tan P. 2006. A modular analysis of breast cancer reveals a novel low-grade molecular signature in estrogen receptor-positive tumors. *Clin Cancer Res* **12**: 3288–3296.
- Zhao H, Langerod A, Ji Y, Nowels KW, Nesland JM, Tibshirani R, Bukholm IK, Karesen R, Botstein D, Borresen-Dale AL, et al. 2004. Different gene expression patterns in invasive lobular and ductal carcinomas of the breast. *Mol Biol Cell* **15**: 2523–2536.

Growth factor stimulation induces a distinct ER α cistrome underlying breast cancer endocrine resistance

Mathieu Lupien, Clifford A. Meyer, Shannon T. Bailey, et al.

Genes Dev. 2010, **24**:

Access the most recent version at doi:[10.1101/gad.1944810](https://doi.org/10.1101/gad.1944810)

Supplemental Material

<http://genesdev.cshlp.org/content/suppl/2010/09/27/24.19.2219.DC1>

References

This article cites 69 articles, 29 of which can be accessed free at:
<http://genesdev.cshlp.org/content/24/19/2219.full.html#ref-list-1>

License

Email Alerting Service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).