Tissue-specific splicing of a ubiquitously expressed transcription factor is essential for muscle differentiation

Soji Sebastian,¹ Hervé Faralli,^{1,4} Zizhen Yao,^{2,4} Patricia Rakopoulos,^{1,3} Carmen Palii,¹ Yi Cao,² Kulwant Singh,¹ Qi-Cai Liu,¹ Alphonse Chu,¹ Arif Aziz,¹ Marjorie Brand,^{1,3} Stephen J. Tapscott,² and F. Jeffrey Dilworth^{1,3,5}

¹Sprott Center for Stem Cell Research, Ottawa Hospital Research Institute, Ottawa, Ontario K1H 8L6, Canada; ²Human Biology Division, Fred Hutchinson Cancer Research Center, Seattle, Washington 98109, USA; ³Department of Cellular and Molecular Medicine, University of Ottawa, Ontario K1H 8L6, Canada

Alternate splicing contributes extensively to cellular complexity by generating protein isoforms with divergent functions. However, the role of alternate isoforms in development remains poorly understood. Mef2 transcription factors are essential transducers of cell signaling that modulate differentiation of many cell types. Among *Mef2* family members, *Mef2D* is unique, as it undergoes tissue-specific splicing to generate a muscle-specific isoform. Since the ubiquitously expressed (*Mef2D* α 1) and muscle-specific (*Mef2D* α 2) isoforms of *Mef2D* are both expressed in muscle, we examined the relative contribution of each *Mef2D* isoform to differentiation. Using both *in vitro* and *in vivo* models, we demonstrate that *Mef2D* isoforms act antagonistically to modulate differentiation. While chromatin immunoprecipitation (ChIP) sequencing analysis shows that the *Mef2D* isoforms bind an overlapping set of genes, only *Mef2D* α 2 activates late muscle transcription. Mechanistically, the differential ability of *Mef2D* isoforms to activate transcription depends on their susceptibility to phosphorylation by protein kinase A (PKA). Phosphorylation of *Mef2D* α 1 by PKA provokes its association with corepressors. Conversely, exon switching allows *Mef2D* α 2 to escape this inhibitory phosphorylation, permitting recruitment of Ash2L for transactivation of muscle genes. Thus, our results reveal a novel mechanism in which a tissue-specific alternate splicing event has evolved that permits a ubiquitously expressed transcription factor to escape inhibitory signaling for temporal regulation of gene expression.

[**Keywords:** alternative splicing; Mef2; PKA signaling; gene expression; HDAC; Ash2L]

Supplemental material is available for this article.

Received February 4, 2013; revised version accepted April 26, 2013.

Genome-wide studies have highlighted the capacity of cells to exploit the 20,000 genes of the human genome to produce a far more complex proteome through the generation of alternate transcripts (The ENCODE Project Consortium et al. 2012). A major contributor to this cellular complexity is alternative splicing of exons, which has been proposed to occur universally in multiexon genes (Wang et al. 2008). The generation of these alternatively spliced transcripts can occur through several mechanisms, including (1) the inclusion/exclusion of an exon, (2) alternate 3' or 5' splice donor/acceptor sites, or (3) the alternate use of mutually exclusive exons (Kalsotra and Cooper 2011). In the case of mutually exclusive exons, the con-

trolled use of alternate exons allows for the expression of functionally distinct protein isoforms that have the potential to perform specific tasks in a temporally or spatially restricted manner. Furthermore, ontology analysis suggests that transcripts incorporating mutually exclusive exons are likely to play a role in tissue-specific functions such as differentiation (Wang et al. 2008). However, our knowledge concerning the role of specific alternative splicing events in regulating differentiation remains limited.

During myogenesis, myoblasts undergo a temporally ordered series of changes in gene expression, giving rise to multinucleated myotubes (Bergstrom et al. 2002). These alterations are extensive, as RNA sequencing (RNA-seq) analysis has shown that 7770 genes change their expression during myotube formation (Trapnell et al. 2010). In addition to these changes in transcript abundance, alternate mRNA processing leads to a switch in the major isoform of 7% of alternative spliced genes (Trapnell et al.

⁴These authors contributed equally to this work.

⁵Corresponding author

E-mail jdilworth@ohri.ca

Article published online ahead of print. Article and publication date are online at <http://www.genesdev.org/cgi/doi/10.1101/gad.215400.113>. Freely available online through the *Genes & Development* Open Access option.

2010). This drastic change in the myogenic transcriptome suggests an important role for alternative splicing in establishing the myotube fate. Among the genes that undergo a major isoform switch during myogenesis is the transcription factor *Mef2D* (Trapnell et al. 2010). *Mef2D* is a member of the *Mef2* family of transcription factors, which act as key mediators of signal-dependent transcription in many different cell types to control developmental processes such as differentiation (Potthoff and Olson 2007). Interestingly, the switch in *Mef2D* isoforms observed during muscle differentiation is a result of an alternate use of mutually exclusive exons (see Fig. 1) that generates an isoform that is expressed exclusively in skeletal and cardiac muscle (Martin et al. 1994). However, the functional relevance of this muscle-specific isoform of *Mef2D* to muscle development remains unknown. Here we sought to elucidate the differential roles of the ubiquitously expressed (*Mef2D* α 1) and muscle-specific (*Mef2D* α 2) isoforms of *Mef2D* in skeletal myogenesis.

Results

The muscle-specific isoform of Mef2D drives the late stages of myogenesis

Advances in our understanding of the relative roles of *Mef2D* isoforms in myogenesis have been hampered by the lack of reagents that distinguish between the endogenous proteins. Consequently, we generated antibodies that discriminate between the muscle-specific *Mef2D* α 2 isoform and its ubiquitously expressed counterpart, *Mef2D* α 1 (Fig. 1A). Western blot of recombinant *Mef2D* proteins demonstrate the specificity of our antibodies (Supplemental Fig. 1). Furthermore, these antibodies recognize the endogenous *Mef2D* isoforms in differentiating myoblasts using Western blot, immunofluorescence, and immunoprecipitation (Supplemental Fig. 1; data not shown). Importantly, a similar efficiency of immunoprecipitation is observed with both isoform-specific antibodies.

Initially, we characterized the expression of *Mef2D* isoforms during murine myogenesis. Temporal analysis demonstrates a clearly distinct pattern of expression for the two *Mef2D* isoforms (Fig. 1B,C). In the case of the ubiquitously expressed *Mef2D* α 1, the protein is present in proliferating myoblasts and continues to accumulate throughout differentiation. In contrast, the muscle-specific *Mef2D* α 2 is absent from proliferating myoblasts and is strongly induced between 48 and 72 h of differentiation (Fig. 1B,C). This pattern of expression suggests a role for *Mef2D* α 2 in late myogenesis.

To further explore the role of *Mef2D* isoforms in myogenesis, we created isogenic C2 myoblast lines where the exogenous expression of either *Mef2D* α 1 or *Mef2D* α 2 can be induced upon doxycycline (Dox) treatment (Supplemental Fig. 2A). Interestingly, myoblasts induced to express *Mef2D* α 2 formed multinucleated, myosin heavy chain (MHC)-positive cells more quickly than the control cells (Fig. 2A; Supplemental Fig. 2B), suggesting an accelerated differentiation. In contrast, expression of *Mef2D* α 1 appeared to block myogenesis, as fewer MHC⁺ cells were

Figure 1. *Mef2D* undergoes alternative splicing during muscle differentiation to up-regulate the *Mef2D* α 2 isoform. (A) Schematic representation of the intron–exon structure of the *Mef2D* gene. Alternative incorporation of the mutually exclusive third (α 1) or fourth (α 2) exon results in the generation of an mRNA encoding *Mef2D* α 1 or *Mef2D* α 2, respectively. The DNA-binding domain of *Mef2D* proteins is shown in brown (MADS). (B,C) The *Mef2D* α 2 isoform is up-regulated during muscle differentiation. (B) Whole-cell extracts were prepared from C2 cells that were differentiated for various lengths of time (0–96 h). Western blot was then performed using antibodies specifically recognizing *Mef2D* α 1, *Mef2D* α 2, *Mef2D* (pan), and Ash2L as indicated. (C) Immunofluorescence analysis was performed on C2 cells that were differentiated for various lengths of time (0–96 h). Slides were stained using antibodies recognizing *Mef2D* α 1, *Mef2D* α 2, or *Mef2D* (pan) as indicated.

observed in this population at late stages (120 h) of differentiation. Having observed opposing effects of *Mef2D* α 1 and *Mef2D* α 2 on differentiation in cultured myoblasts, we next examined whether these disparate roles of *Mef2D* isoforms are also observed in vivo using a model of mouse muscle regeneration. Indeed, injection of

Figure 2. Mef2D α 1 and Mef2D α 2 have opposing effects on muscle differentiation and regeneration. (A) Stable isogenic C2iFRT myoblast lines were induced to express Mef2D α 1, Mef2D α 2, or control (parental cells). After 24 h of Dox treatment, cells were incubated in low-serum medium supplemented with Dox to permit differentiation in the presence of exogenous Mef2D isoforms (or control). Immunofluorescence was performed using antibodies recognizing MHC or DAPI as indicated. (B) Cardiotoxin-injured mouse TA muscles were infected with retrovirus expressing Mef2D α 1, Mef2D α 2, or vector control. After 15 d of regeneration, cross-sections were stained using hematoxylin and eosin. The cross-sectional area of individual fibers was calculated and plotted. $n = 3$.

retrovirus expressing Mef2D α 1 into cardiotoxin-injured tibialis anterior (TA) muscle resulted in severely impaired regeneration where myofiber repair was delayed (Fig. 2B; Supplemental Fig. 3A), resulting in an accumulation of undifferentiated MyoD $^{+}$ muscle cells in the interstitial space (Supplemental Fig. 4). In contrast, injection of retrovirus expressing Mef2D α 2 resulted in improved muscle regeneration with an increased muscle fiber caliber compared with the contralateral control TA muscle (Fig. 2B; Supplemental Fig. 3B). Furthermore, only the injection of Mef2D α 2 could mediate the previously described Mef2D-dependent fiber type switching in the TA muscle (Potthoff et al. 2007) from a predominantly fast type MHC-IIb myofiber toward a slower muscle isotype that contained a mixture of MHC-IIb and MHC-IIa myofibers (Supplemental Fig. 5). Thus, Mef2D α 1 and Mef2D α 2 have antagonistic effects on myogenesis and muscle regeneration.

Examination of Mef2D α 2-expressing myoblasts showed that MHC proteins were expressed at the earliest stages of differentiation (0 h) (Fig. 2A). This temporal shift in MHC expression led us to examine whether Mef2D α 2 expression would allow for the expression of muscle genes in proliferating myoblasts. To our surprise, RT-qPCR analysis demonstrates that expression of Mef2D α 2, but

not Mef2D α 1, can activate *Myh3* along with other late myogenic genes (*Myh3* and *Tnnt1*) in proliferating myoblasts (Supplemental Fig. 6). To determine whether the expression of these late markers of myogenesis was due to a precocious differentiation, we examined the ability of Mef2D isoforms to activate expression of *Myog* and *Ckm*—two muscle-specific genes normally up-regulated at intermediate stages of myogenesis. In contrast to *Myh3*, which is strongly up-regulated by ectopic expression of Mef2D α 2 in proliferating myoblasts, we observed no change in transcription of *Myog* or *Ckm* upon expression of the muscle-specific isoform (Supplemental Fig. 6). Taken together, our findings suggest that alternate splicing generates the muscle-specific Mef2D α 2 isoform to temporally regulate activation of late muscle genes.

We next sought to understand the mechanism underlying the differential ability of Mef2D isoforms to activate late muscle gene expression. Two potential models were envisaged: (1) Mef2D α 1 and Mef2D α 2 recruit similar coregulatory proteins to an alternate subset of muscle genes or (2) Mef2D α 1 and Mef2D α 2 bind to a similar set of genomic loci, leading to the recruitment of alternate coregulatory proteins. We therefore explored the contribution of these two models to the functional differences observed between Mef2D isoforms.

Genome-wide binding shows that Mef2D isoforms bind to a largely overlapping set of genes

We first explored the contribution of altered genomic targeting to the differential ability of Mef2D isoforms to activate transcription. For this, we performed chromatin immunoprecipitation (ChIP) sequencing (ChIP-seq) to examine the genomic distribution of Mef2D α 1 and Mef2D α 2 in myotubes. Fragmented chromatin was immunoprecipitated using our isoform-specific antibodies to isolate DNA associated with the endogenous Mef2D α 1 and Mef2D α 2 proteins (Supplemental Figs. 7, 8). Bioinformatic analysis of ChIP-seq data identified 7910 binding sites for Mef2D α 1, while the muscle-specific isoform Mef2D α 2 bound to 9103 sites that are not present in the control (Fig. 3A; Supplemental Table 1). Comparison of the Mef2D-binding sites using a highly stringent cutoff of $P < 10^{-6}$ revealed that 66% of Mef2D α 1-bound peaks are also bound by Mef2D α 2 (Fig. 3A). However, when the stringency is modestly reduced to $P < 10^{-4}$, the overlap in binding of Mef2D α 1 at Mef2D α 2-bound peaks increases to 85% (data not shown). Consistent with the fact that Mef2D α 1 and Mef2D α 2 bind to a largely overlapping set of genomic loci, position weight matrices for Mef2D-binding elements observed under these peaks show that both isoforms of the protein recognize a nearly identical consensus sequence of CTAAAAATAG (Fig. 3B). A de novo search for DNA sequences enriched under peaks identified Mef2 and E-box sequences (i.e., MyoD/Myog/Myf5/MRF4-binding sites) as the two most highly represented DNA elements at both Mef2D α 1 and Mef2D α 2 genomic targets (data not shown). Interestingly, examination of genomic loci bound preferentially by either Mef2D α 1 or Mef2D α 2 did not identify Mef2-binding sites as the most highly represented DNA sequences found under these peaks (Supplemental Fig. 9). This finding suggests that the differential binding observed between Mef2D α 1 and Mef2D α 2 at a small subset of genes is likely due to altered targeting of the transcription factor to DNA through an indirect tethering mechanism.

To assess the importance of overlapping Mef2D α 1 and Mef2D α 2 genomic binding to myogenesis, we performed gene ontology (GO) analysis of peaks bound by both isoforms. Genes targeted by both Mef2D α 1 and Mef2D α 2 are strongly associated with GO annotations defining a role in muscle development (Fig. 3C; Supplemental Table 1). Among these genes, we examined Mef2D binding at the *HDAC9* gene, which is known to participate in a negative feedback loop with Mef2 proteins (Haberland et al. 2007). At this locus, we observed peaks within the *HDAC9* promoter for both Mef2D α 1 and Mef2D α 2 that overlap with MyoD binding (Fig. 3D). Similarly, we observed that the promoters of *Myh3*, *Titin*, *Myog*, *Tnni1*, *Ckm*, and *Mylpf* are bound equally well by both Mef2D α 1 and Mef2D α 2 in differentiating myotubes (Supplemental Fig. 8). RNA-seq analysis of transcripts that are up-regulated at 24 h of differentiation when expression of Mef2D isoforms is induced in C2 cells identified 227 genes activated by Mef2D α 2 (Supplemental Table 1). Combining the RNA-seq data set with Mef2D peaks within 50 kb of the

transcription start site identified 107 genes whose expression may be directly modulated by Mef2D α 2 (Supplemental Table 1; Supplemental Fig. 10). Consistent with Mef2D α 2 modulating the late muscle gene expression program, GO analysis of putative Mef2D α 2 direct target genes showed a highly significant enrichment for genes involved in muscle development and muscle contraction (Supplemental Fig. 10A). Importantly, although Mef2D α 1 bound to most of the direct transcriptional targets of Mef2D α 2, exogenous expression of the ubiquitous Mef2D isoform did not lead to a significant increase in expression at any of these genes (Supplemental Table 1; Supplemental Fig. 10B). Similarly, ChIP-qPCR analysis of Mef2D isoform binding in proliferating myoblasts demonstrates that both Mef2D α 1 and Mef2D α 2 bind to the *Myh3* and *Mylpf* promoters (Supplemental Fig. 6B), although only Mef2D α 2 binding gives rise to transcriptional activation. Taken together, these results demonstrate that altered DNA binding is not the major determinant of the differential activation of transcription by Mef2D isoforms.

The muscle-specific Mef2D α 2 isoform preferentially interacts with coactivator complexes

To assess the possibility that Mef2D α 1 and Mef2D α 2 associate with alternate coregulatory proteins to regulate muscle gene expression, we performed a proteomic analysis of proteins that interact with each Mef2D isoform. Initially, we examined proteins that interact with recombinant Mef2D α 1 and Mef2D α 2. In a pull-down assay, Flag-tagged Mef2D α 1 or Mef2D α 2 proteins were used as bait to isolate interacting proteins from nuclear extracts prepared from myotubes (Supplemental Fig. 11). Mass spectrometry analysis of Mef2D α 1- or Mef2D α 2-interacting proteins (Table 1) identified components of the coactivator Ash2L methyltransferase complexes, corepressor HDAC deacetylase complexes, and the SWI/SNF ATP-dependent chromatin remodeling complexes that have previously been reported to associate with Mef2D (Ohkawa et al. 2006, 2007; Rampalli et al. 2007; Du et al. 2008). Qualitative analysis of the identified peptides showed that recombinant Mef2D α 1 and Mef2D α 2 interact with the same sets of proteins (Table 1). This finding suggests that the exchange of mutually exclusive exons in Mef2D does not generate proteins with distinct protein-protein interaction domains. However, the experiment was performed with recombinant Mef2D isoforms. To determine whether Mef2D α 1 and Mef2D α 2 isoforms also interact with the same sets of proteins in their native cellular context, we immunoprecipitated endogenous Mef2D α 1 and Mef2D α 2 proteins from myotube extracts and tested the interaction with the previously identified cofactors by Western blot (Fig. 4A). Strikingly, in these conditions, we observed differential interactions for Mef2D α 1 versus Mef2D α 2. Precisely, the muscle-specific Mef2D α 2 interacts with the Ash2L coactivator complex but not the repressive HDAC4 and HDAC9 proteins. Conversely, the ubiquitously expressed Mef2D α 1 does not associate with

Figure 3. Mef2D isoforms bind a largely overlapping set of genes involved in muscle development. ChIP-seq analysis was performed on C2 myotubes that had been differentiated for 120 h. Chromatin was immunoprecipitated with antibodies recognizing either Mef2D α 1 or Mef2D α 2. To control for nonspecific binding to the antibodies, control ChIP-seq experiments were performed with antibodies that were preincubated with peptides corresponding to the epitope and labeled as Mef2D α 1 (blocked) or Mef2D α 2 (blocked). (A) Venn diagram showing the overlap between the 9103 Mef2D α 2 peaks and the 7910 Mef2D α 1 peaks identified by ChIP-seq. (B) Logo diagrams of the position weight matrix of the sequence generated based on the enrichment of Mef2-binding sites observed in peaks bound by either Mef2D α 1 or Mef2D α 2. (C) The top GO terms for biological processes (BP) are shown for the group of 5204 genes that are bound equally well by both Mef2D α 1 and Mef2D α 2. (D) The sequencing reads are mapped across the *HDAC9* locus for ChIP experiments performed using Mef2D α 1, Mef2D α 2, Mef2D α 1 (blocked), Mef2D α 2 (blocked), and MyoD. Peaks for both Mef2D α 1 and Mef2D α 2 that overlap with MyoD binding are observed in the proximal promoter region (boxed in red).

Ash2L but interacts strongly with HDAC4 and HDAC9. Therefore, in contrast to their recombinant counterparts, endogenous Mef2D isoforms interact with distinct

protein complexes in differentiating myotubes—Mef2D α 1 interacts with corepressors, while Mef2D α 2 interacts with coactivators.

Table 1. Proteomics analysis of Mef2D α 1- and Mef2D α 2-interacting proteins

Identified proteins	Mef2D α 1		Mef2D α 2	
	Peptides	Mascot score	Peptides	Mascot score
Transcription factors				
Mef2D	35	2848	24	1535
Myogenin	4	119	3	74
Jun	4	217	5	208
Jun-B	4	197	4	275
Maf-G	4	78	2	49
Pbx3	2	39	—	—
Ash2L complex				
Ash2L	6	89	10	145
WDR5	4	185	8	180
DPY-30	2	85	—	—
HCFC1	4	88	2	54
HDAC complexes				
HDAC2	—	—	3	23
SAP18	1	25	1	21
SAP30	1	26	1	31
RbBP4	9	261	11	341
RbBP7	7	198	8	249
SWI/SNF complex				
BRG1	—	—	2	38
BAF250a/Arid1a	15	236	9	50
BAF180	1	44	—	—
BAF170	18	455	15	345
BAF155	18	400	12	202
BAF60c	10	359	14	409
BAF57	7	296	5	223
Brd7	2	53	2	37

Recombinant Flag-tagged Mef2D α 1 and Mef2D α 2 proteins were used as bait to pull down interacting proteins from nuclear extracts prepared from C2 myotubes. Proteins interacting with the Mef2D isoforms were separated by SDS-PAGE and analyzed by mass spectrometry as described in the Supplemental Material. (Peptides) The number of unique peptides identified for each protein; (mascot score) the confidence in the proper identification of the protein; (—) no peptides were identified for that particular protein.

Mef2D α 2 protein evades inhibitory protein kinase A (PKA) signaling

Based on the difference in protein-binding specificity observed between recombinant and endogenous proteins, we reasoned that the altered interactions could be modulated by post-translational modifications (PTMs). Indeed, several PTMs have been shown to regulate Mef2D function in vivo (Potthoff and Olson 2007). Important to transcriptional activation, phosphorylation of Mef2D by p38 α MAPK (p38 α) at positions Thr308 (T308) and T315 (see Fig. 4B) is required for recruitment of the Ash2L coactivator complexes to muscle genes (Rampalli et al. 2007). We therefore examined the propensity of Mef2D α 1 and Mef2D α 2 to be phosphorylated by p38 α . Using recombinant proteins, in vitro kinase assays demonstrate that both Mef2D α 1 and Mef2D α 2 are efficiently phosphorylated by p38 α (Fig. 4C). Having observed that the activating phosphorylation of Mef2D by p38 α occurs

in both Mef2D α 1 and Mef2D α 2, we next explored the possibility that the isoforms may be differentially affected by inhibitory PTMs. Transactivation by Mef2D is known to be inhibited through phosphorylation of Ser119 (S119) and S190 by PKA, which enables an association with class II HDACs (Du et al. 2008). Importantly, alternative exon usage in Mef2D α 2 removes the S119 residue targeted by PKA in the Mef2D α 1 isoform (see Fig. 4B). Thus, we examined whether the Mef2D α 1 and Mef2D α 2 isoforms are differentially phosphorylated by PKA. Consistent with previous observations (Du et al. 2008), Mef2D α 1 is strongly phosphorylated by PKA (Fig. 4C). In contrast, Mef2D α 2 is resistant to phosphorylation by PKA even though the phosphorylation target site S189(190) remains intact (Fig. 4C). These findings confirm the hypothesis that Mef2D isoforms are differentially modified by PTMs. Thus, overall, our results show that alternative exon usage regulates the ability of Mef2D protein to be phosphorylated by PKA.

Mutation of PKA phosphorylation sites allows Mef2D α 1 to activate myogenic transcription

To better understand the importance of the differential phosphorylation of Mef2D isoforms by PKA, we examined how PKA activity changes during differentiation. Previous studies have shown that PKA signaling is required at various stages of myogenesis to mediate CREB-dependent gene expression (Berdeaux and Stewart 2012). Using antibodies that recognize CREB phosphorylated at S133 to measure PKA activity, we observed that the PKA phosphorylates CREB in proliferating myoblasts, and the kinase remains active as myoblasts transition to myotubes (Fig. 4D). Thus, consistent with the requirement for activated CREB in myogenesis (Berdeaux and Stewart 2012), PKA activity is maintained throughout myotube formation.

The fact that PKA activity is maintained throughout differentiation suggests that the differential propensity to be phosphorylated by PKA could be responsible for the differential interaction of Mef2D isoforms with co-repressor versus coactivators. To test this possibility, we generated the mutant Mef2D α 1(S119A/S190A), which is resistant to phosphorylation by PKA (Du et al. 2008). Similar to results observed with the endogenous protein, overexpressed Mef2D α 1 did not interact with Ash2L but showed a strong interaction with HDAC9 (Fig. 5A). However, inhibition of phosphorylation by PKA using Mef2D α 1(S119/S190A) led to a marked reduction in the interaction between the mutant protein and the repressive HDAC9 protein. Surprisingly, the loss in interaction with HDAC9 is concurrent with the appearance of an association with the coactivator Ash2L that normally interacts with Mef2D α 2. Thus, phosphorylation of Mef2D proteins by PKA is a primary determinant of the interaction with transcriptional coactivators or corepressors.

Finally, we examined whether the switch in Mef2D isoforms could represent a requirement for the transcription factor to evade inhibitory PKA signaling to mediate

Coomassie blue. (D) PKA activity in C2 cells was determined at different stages of differentiation by examining the amount of phosphorylated CREB present in whole-cell extracts. Western blots were performed on whole-cell extracts using the antibodies indicated.

activation of late gene transcription. Using RT-qPCR, we observed that, unlike wild-type Mef2Dα1, the PKA phosphorylation mutant Mef2Dα1(S119/S190A) gives rise to a strong activation of both *Myh3* and *Mylpf* expression (Fig. 5B). Consistent with the differential ability to interact with coregulatory proteins, ChIP analysis showed that expression of Mef2Dα2 or Mef2Dα1(S119/S190A) leads to an enrichment of Ash2L at both the *Mylpf* and *Myh3* gene promoters, while expression of Mef2Dα1 results in an increased enrichment of HDAC9 at these same loci (Fig. 5C). These findings demonstrate that Mef2Dα1(S119/S190A) and Mef2Dα2 both activate expression of late muscle genes through the recruitment of coactivator proteins led us to examine whether the inability of the ubiquitously expressed Mef2Dα1 protein to stimulate myotube formation may be due to PKA phosphorylation. As observed above, immunofluorescence shows that myoblasts expressing Mef2Dα1 are resistant to the formation of MHC⁺ myotubes at 48 h of differentiation (Fig. 5D). In contrast, expression of the PKA mutant Mef2Dα1(S119/S190) gave rise to MHC⁺ myotubes with a level of differentiation similar to that observed with Mef2Dα2 (Fig. 5D; Supplemental Fig. 12). This demonstrates that the differential ability of Mef2Dα1 and Mef2Dα2 to drive myogenesis depends on their propensity for PKA-mediated phosphorylation. Thus, our results reveal a new role for alternative use of mutually exclusive exons in generating distinct transcription factors that have antagonizing roles in regulating differentiation based on their susceptibility to cell signaling-induced PTMs.

Discussion

Alternative splicing of transcription factors has been shown to direct tissue-specific gene expression by modifying the specificity of DNA-binding element recognition of the protein, resulting in the activation of an alternate set of genes (Reed et al. 2010; Gabut et al. 2011). Performing the first comparative genome-wide analysis of alternate transcription factor isoforms, we determined that the generation of a muscle-specific Mef2Dα2 isoform does not alter the DNA-binding element recognition of the transcription factor compared with the more ubiquitous Mef2Dα1 isoform. Instead, we found that alternate usage of mutually exclusive exons plays a crucial role in controlling the transcriptional activity of Mef2D during differentiation by altering its susceptibility to phosphorylation by PKA. Specifically, the Mef2Dα1 isoform that is phosphorylated by PKA acts as a transcriptional repressor, while the PKA-resistant Mef2Dα2 isoform activates late muscle gene expression. Thus, we identified a previously unappreciated mechanism by which a ubiquitously expressed transcription factor undergoes tissue-specific alternative exon usage to permit temporal activation of gene expression during differentiation.

A role for the Mef2 family of proteins in modulating gene expression in response to cell signaling pathways is well established (Potthoff and Olson 2007). Among the signaling pathways known to regulate Mef2 activity in development, PKA and p38 MAPK signaling are known to converge on Mef2D function during myogenesis. Studies using reagents that did not distinguish between

Figure 5. Mutation of PKA phosphorylation sites in Mef2Dα1 facilitates its ability to activate MHC expression. (A) Ash2L complex interacts with forms of Mef2D that are not phosphorylated by PKA. Nuclear extracts were prepared from C2iFRT lines induced to express a Flag-tagged version of either Mef2Dα2(WT), Mef2Dα2(S189A), Mef2Dα1(WT), Mef2Dα1(S190A), or Mef2Dα1(S119/190A) that had been incubated in differentiation medium for 48 h. Western blots were performed using the antibodies indicated. (B,C) C2iFRT cells induced to express either Mef2Dα2(WT), Mef2Dα1(WT), Mef2Dα1(S190A), or Mef2Dα1(S119/190A) were incubated in differentiation medium for 24 h. (B) Total RNA was converted to cDNA and analyzed using qPCR with primers specific for Myh3, Mylpf, p21^{cip}, or the ubiquitously expressed DDX5 mRNA. Values are expressed as the mean $\pm$ SEM relative to the internal control DDX5 where the expression in the control was normalized to 1. $n = 3$. (C) Cross-linked chromatin from C2iFRT cells expressing exogenous Flag-tagged Mef2D proteins were subjected to ChIP analysis using antibodies directed against the Flag epitope. Immunopurified DNA was quantitated by qPCR using probes that recognize the promoter of the Myh3 or Mylpf genes or the upstream enhancer of IgH. (D) C2iFRT cells induced to express Mef2Dα2(WT), Mef2Dα1(WT), Mef2Dα1(S190A), or Mef2Dα1(S119/190A) were incubated in differentiation medium for 48 h. Immunofluorescence was performed using antibodies recognizing MHC (green) or DAPI (blue) as indicated.

Mef2Dα1 and Mef2Dα2 have shown that PKA represses Mef2D function (Du et al. 2008), while p38α stimulated its transactivation (Penn et al. 2004; Rampalli et al. 2007). In the case of p38α, phosphorylation of Mef2D allows the recruitment of Ash2L/MLL2-containing methyltransferase complexes to establish trimethylation of histone H3 (H3K4me3) within promoters of MyoD target genes to facilitate transcription (Rampalli et al. 2007; Aziz et al. 2010). In contrast, PKA-mediated phosphorylation of Mef2D engages an interaction between the transcription factor and class II HDACs to mediate transcriptional repression (Du et al. 2008). The use of antibodies that allow us to distinguish between Mef2Dα1 and Mef2Dα2 confirm these original findings, as Mef2D proteins can interact with both corepressors and coactivators during myogenesis. However, our analysis of specific isoforms demonstrates that it is not the same Mef2D molecules that interact with these antagonizing coregulatory molecules. Instead, we found that the ubiquitous Mef2Dα1 interacts with the corepressors HDAC4 and HDAC9 through an interaction enabled by PKA-mediated phosphorylation. Splicing of the Mef2D transcripts during myogenesis then generates a muscle-specific protein that is

resistant to PKA phosphorylation and associates with the Ash2L coactivator complex. This represents a significant advance in our understanding of myogenesis, as previous studies showed that PKA phosphorylates Mef2D to block its ability to drive myogenesis (Du et al. 2008). This inhibitory effect of PKA on Mef2D function was perplexing due to the fact that PKA signaling is required in myogenesis to facilitate CREB-dependent transactivation and myoblast fusion (Chen et al. 2005; Mukai and Hashimoto 2008; Stewart et al. 2011; Berdeaux and Stewart 2012). Our finding that Mef2D undergoes a splicing event during differentiation that renders the transcription factor resistant to PKA inhibition resolves this paradox and highlights the need to study isoform-specific functions of proteins during development.

Resolving the paradox of continued PKA signaling during myogenesis is an important finding that identifies an alternate mechanism that evolved for cells to overcome inhibitory signaling pathways during differentiation. Indeed, common models for development describe the down-regulation of signaling pathways that are inhibitory to differentiation concomitant with the up-regulation of prodifferentiation signals to permit progression toward a

new cell fate (Iglesias-Bartolome and Gutkind 2011). However, PKA signaling is both promyogenic through phosphorylation of CREB (Chen et al. 2005; Mukai and Hashimoto 2008; Stewart et al. 2011; Berdeaux and Stewart 2012) and inhibitory to myogenesis through phosphorylation of Mef2D (Du et al. 2008). Thus, the muscle-specific switch in exon usage for Mef2D α 2 evolved to evade the inhibitory effects of PKA signaling, allowing transcriptional activation of Mef2D target genes in the presence of promyogenic PKA signaling. While we showed that the exon switching is used by Mef2D to evade phosphorylation by PKA, it remains to be determined how extensively splicing is used to evade signaling pathways in development. However, we propose that isoform switching may be a widely used mechanism to modify susceptibility of a protein to cell signaling, as recent studies suggest that tissue-specific exons often encode disordered peptides that are enriched for sites of phosphorylation (Buljan et al. 2012).

Profiling of Mef2D α 1 and Mef2D α 2 protein levels during myogenesis demonstrated a temporal control of isoform abundance where a clear switch in exon usage allows accumulation of the muscle-specific Mef2D α 2 starting at an intermediate stage of differentiation. The timing of this splicing event suggests a major role for Mef2D α 2 in

ensuring the proper temporal activation of late gene expression during myogenesis. Previous studies of myogenesis have identified a role for Mef2D in establishing the proper temporal activation of a subset of muscle genes (Penn et al. 2004). Furthermore, studies in zebrafish have demonstrated that Mef2 proteins are important to the activation of the late stages of the myogenic program (Hinitz and Hughes 2007). Based on these observations, we propose a model for the temporal regulation of late muscle gene expression through Mef2D isoform switching (Fig. 6). In this model, Mef2D α 1 is the predominant isoform of the transcription factor expressed in the early myogenesis. Active PKA signaling in the cell catalyzes phosphorylation of Mef2D α 1 to establish an HDAC-dependent transcriptional repression at late muscle genes. As differentiation progresses, Mef2D transcripts undergo alternative splicing, leading to the establishment of two distinct isoforms: the ubiquitous Mef2D α 1 isoform that continues to act as a transcriptional repressor and a muscle-specific Mef2D α 2 isoform that is resistant to PKA signaling and permits the transcriptional activation of the late muscle gene expression program. As our ChIP-seq studies have shown that Mef2D α 1 and Mef2D α 2 bind to overlapping subsets of target genes in the late stages of myogenesis, we propose that an ongoing competition between the two isoforms

Figure 6. Model for the temporal regulation of late muscle gene expression through the use of alternate exons. During the early stages of differentiation, the ubiquitous isoform of Mef2D (Mef2D α 1) is expressed almost exclusively. The majority of the Mef2D α 1 are phosphorylated by active PKA signaling, leading to a recruitment of class II HDAC4/9 onto muscle promoters to maintain a transcriptionally repressed state. At later stages of differentiation, alternative splicing of Mef2D transcripts gives rise to a second population of Mef2D proteins (Mef2D α 2). The Mef2D α 2 isoform, which cannot be phosphorylated by PKA, competes with Mef2D α 1 for binding to target genes. While binding of PKA-phosphorylated Mef2D α 1 leads to recruitment of class II HDACs to repress the promoter, Mef2D α 2 (and unphosphorylated Mef2D α 1) leads to recruitment of the Ash2L complexes to muscle promoters to activate gene expression. This competition would result in an ongoing dynamic exchange of repressive and activating Mef2D transcription factors that is proposed to act as a rheostat to ensure an optimal level of transcription to mediate formation of function myotubes.

In conclusion, we showed that the muscle-specific isoform of Mef2D plays a key role in establishing the late muscle gene expression program. This role is enabled through a temporally regulated switch in exon usage that allows Mef2D to evade inhibitory phosphorylation by PKA. Thus, we identified a novel role for the alternate use of mutually exclusive exons to modulate the transition in function of a transcription factor from repressor to activator through altered susceptibility to an inhibitory PTM.

Antibodies

Cell culture

Generation of isogenic C2 lines expressing Mef2D isoforms

ChIPs

High-throughput DNA sequencing analysis

Sequences were extracted using GAPIipeline. Reads were aligned by Burrows-Wheeler alignment (BWA) to the mouse genome (mm9). Duplicate sequences were discarded to minimize the effects of PCR amplification. Each read was extended in the sequencing orientation to a total of 200 bases to infer the coverage at

each genomic position. Peak calling was performed by an in-house-developed R package as previously described (Palii et al. 2011; Fong et al. 2012). Motif analyses were performed using in-house-developed Bioconductor package motifRG (Palii et al. 2011; Fong et al. 2012). To detect differential peaks between two samples, we first took the square root of the peak heights in both samples and fit a local regression curve on the difference of the two versus the mean. The residual is defined as the difference subtracted by the fitted value, which adjusts for mean-dependent bias. We fit another local regression model on squared residual versus mean to estimate mean-dependent variance. The Z-value is defined by the residual divided by standard deviation (square root of the estimated variance). This approach draws some similarity to the mean-dependent dispersion estimate by DESeq (Anders and Huber 2010), except that we used a nonparametric approach instead of modeling by a negative binomial distribution. ChIP-seq data were deposited in Gene Expression Omnibus (GEO; accession no. GSE43223).

RNA-seq

C2iFRT-Mef2D α 1, C2iFRT-Mef2D α 2, or control (C2iFRT) cells were grown in proliferative conditions for 24 h in the presence of Dox. Near confluence, cells were then differentiated for a further 24 h in the presence of Dox. Total RNA was isolated from cells using the RNeasy minikit (Qiagen) using standard conditions. Library preparation and 100-bp paired-end RNA-seq were performed using standard Illumina procedures for the HiSeq 2000 platform.

The RNA-seq data were analyzed using the Bowtie, TopHat, CuffLinks, and CummeRbund software suite (Trapnell et al. 2012). Briefly, paired-end 100-bp reads were mapped to the mm9 genome (University of California at Santa Cruz) and known mm9 transcripts (RefSeq mm9 build37.2) using Bowtie version 2.0.2 (Langmead and Salzberg 2012) with default parameters. Splice junctions were determined from the mapped reads using TopHat version 2.0.6 (Trapnell et al. 2009) with default values with the exception of $-r = 0$. Transcript assembly and quantification was performed with CuffLinks version 2.0.2 (Trapnell et al. 2010) using default values. Identification of differential gene expression was performed with the CummeRbund bioconductor package (<http://compbio.mit.edu/cummeRbund>). Genes with a Q-value ≤ 0.05 were considered to be significant. RNA-seq data were deposited in GEO (accession no. GSE46363).

Regeneration in mice

Mouse studies were performed in accordance with the University of Ottawa regulations for animal care and handling. The left and right TA muscles of wild-type CD-1 mice (obtained from Charles River) were injured by injection of cardiotoxin as previously described (Liu et al. 2012). After 48 h of recovery, retrovirus expressing either Mef2D α 1 or Mef2D α 2 (multiplicity of infection [MOI] 1×10^6) was injected into one of the regenerating TA muscles, while the contralateral regenerating muscle was injected with a control retrovirus. After 3, 7, or 15 d of recovery, TA muscles were isolated and sectioned (12 μ m thickness). For TA samples taken at 15 d of regeneration, the muscles were prepared for paraffin inclusion as described in Liu et al. (2012). For TA samples prepared at 3 and 7 d of regeneration, unfixed muscles were embedded in OCT compound, frozen in isopentane cooled with liquid nitrogen, and then sectioned (12 μ m thickness). To assess regeneration, cross-sections were prepared for histological examination on a ScanScope CS microscope (Aperio) after staining with hematoxylin and eosin. Changes in fiber size distribution between muscles injected with Mef2D-expressing virus and the

control were determined using a nonparametric (Wilcoxon) test. Fiber type switching was assessed by immunofluorescence using antibodies recognizing different isoforms of MHC. After immunostaining, coverslips were mounted in mounting medium with DAPI (Vector Laboratories, Inc., H-1500) and imaged using a Zeiss LSM 510 Meta confocal microscope.

Protein–protein interaction analysis

Nuclear extracts were prepared from differentiated C2 myotubes (5 d) as previously described (Weil et al. 1979) with the following modifications. Nuclear pellets were resuspended in one pellet volume of modified buffer C (20 mM Hepes at pH 7.6, 1.5 mM MgCl₂, 650 mM KCl, benzonase [2.5 U per 10^7 cells], 0.2 mM PMSF, 0.5 mM DTT, 5 mM β -glycerolphosphate, 1 mM sodium orthovanadate) and rotated for 30 min at 4°C. Nuclear homogenates were then diluted with one pellet volume of buffer E (20 mM Hepes at pH 7.6, 1.5 mM MgCl₂, 0.2 mM EDTA, 1 mM PMSF, 0.5 mM DTT, 5 mM β -glycerolphosphate, 1 mM sodium orthovanadate). Extracted proteins were recovered from the nuclei by centrifugation at 15,000g for 30 min at 4°C. Nuclear extracts were then quantitated for protein content prior to interaction studies.

For immunoprecipitation studies of endogenous proteins, 10 μ g of specific antibody or rabbit IgG were prebound to 20 μ L of protein A Dynabeads. Flag immunoprecipitations were performed using M2-agarose (Sigma). Mef2D-associated proteins were immunoprecipitated as previously described (Chaturvedi et al. 2012) and eluted by incubating with 5% acetic acid for 5 min at 22°C. The acetic acid was removed from the recovered proteins by evaporation using a SpeedVac (Eppendorf) before proceeding with Western blot analysis.

In the pull-down studies, baculovirus-expressed Flag-Mef2D α 1 or Flag-Mef2D α 2 proteins were purified from Sf9 insect cells using M2-agarose (Sigma) as previously described (Rampalli et al. 2007). Briefly, M2-agarose-bound proteins were washed in stringent conditions (500 mM KCl) and then equilibrated in immunoprecipitation buffer (20 mM Hepes at pH 7.6, 150 mM KCl, 1.5 mM MgCl₂, 0.2 mM EDTA, 0.2 mM PMSF, 0.5 mM DTT). Flag-Mef2D-bound M2 agarose beads were then incubated with nuclear extracts for 4 h at 4°C. After washing, proteins were eluted with 3X-Flag peptide (Sigma) and separated by SDS-PAGE. After visualization with Coomassie blue, protein bands were excised for mass spectrometry analysis.

Mass spectrometry

Protein identification was performed at the Ottawa Hospital Research Institute Proteomics Core Facility (Ottawa, Canada). Gel bands were in-gel-digested according to the method of Shevchenko et al. (2006). Peptide extracts were concentrated by SpeedVac (Eppendorf) and resuspended in 0.1% trifluoroacetic acid. Peptides were loaded onto a peptide trap (Zorbax SB300-C18, Agilent) for 5 min at 15 μ L/min using a Dionex Ultimate 3000 RSLC nano high-pressure liquid chromatography (HPLC). Peptides were eluted over a 60-min gradient of 3%–45% acetonitrile with 0.1% formic acid at 0.3 μ L/min onto a 10-cm analytical column (New Objective Picofrit self-packed with Zorbax C18) and sprayed directly into a LTQ Orbitrap XL hybrid mass spectrometer using a nanospray source (Thermo Scientific). Mass spectra were acquired in a data-dependent fashion, with MS scans acquired in the Fourier Transform cell, while MS² scans were acquired in the ion trap module.

Tandem mass spectrometry (MS/MS) spectra were matched against a custom database comprised of mouse sequences in SwissProt (2011_07 version of uniprot_sprot.fasta.gz from <ftp://>

ftp.uniprot.org) concatenated with a database of common contaminants or against the full SwissProt database using MASCOT 2.3.01 software (Matrix Science) with mass spectrometry tolerance of ± 5 parts per million (ppm) and MS/MS tolerance of 0.6 Da. For standard Mascot search, oxidation of methionine, carbamidomethylation of cysteine, deamidation, protein N-terminal acetylation, and conversion of peptide N-terminal Glu or Gln to Pyro-Glu were allowed as potential protein modifications.

Immunofluorescence

Cells were fixed using 4% paraformaldehyde, permeabilized with 0.5% Triton X-100/PBS, and then blocked with 5% BSA in PBS. Primary antibodies were diluted with PBS/5% BSA solution and incubated overnight at 4°C. After washing, cells were incubated with secondary antibodies conjugated with Alexa Fluor 488 (Invitrogen, A11001) and/or Alexa Fluor 546 (Invitrogen, A11010). Prior to visualization, coverslips were mounted in mounting medium with DAPI (Vector Laboratories, Inc., H-1500). Images were taken using an Olympus FluoView FV1000 confocal microscope.

Kinase assay

Baculovirus-expressed Flag-Mef2D α 1 or Flag-Mef2D α 2 proteins were purified from Sf9 insect cells using M2-agarose (Sigma). His-p38 α was purified from bacteria in its active (phosphorylated) form as previously described (Khokhlatchev et al. 1997). Active PKA was purchased from New England Biolabs. Recombinant Mef2D proteins (100 ng) were incubated with either His-p38 α (80, 250, or 750 fmol) or PKA (50, 100, or 200 U) in the presence of 1 pmol (4 μ Ci) of [32 P] γ -ATP in a buffer containing 10 mM Hepes (pH 7.4), 50 mM KCl, 1.5 mM MgCl₂, 10% glycerol, 0.5 mM EGTA, and 10 mM β -glycerophosphate for 30 min at 37°C. Reactions were then separated on 10% SDS-PAGE gels, dried, and exposed to autoradiographic film. Parallel reactions lacking [32 P] γ -ATP were run on 10% SDS-PAGE gels and stained for protein content with Coomassie blue.

Acknowledgments

We thank Lynn Megeney and Alexandre Blais for critically reading the manuscript, Alexandre Montpetit for ChIP-seq and RNA-seq library sequencing, and Lawrence Puente for mass spectrometry analysis. This work was funded by the Canadian Institute of Health Research (MOP-77778 to F.J.D., and MOP-89834 to M.B.) and National Institutes of Health (NIH National Institute of Arthritis and Musculoskeletal and Skin Diseases R01AR045113 to S.J.T.). S.S. and C.P. were funded by the Ontario Research Fund supporting the International Regulome Consortium. F.J.D. holds a Canada Research Chair in the Epigenetic Regulation of Transcription.

References

Anders S, Huber W. 2010. Differential expression analysis for sequence count data. *Genome Biol* **11**: R106.
 Aziz A, Liu QC, Dilworth FJ. 2010. Regulating a master regulator: Establishing tissue-specific gene expression in skeletal muscle. *Epigenetics* **5**: 692–696.
 Berdeaux R, Stewart R. 2012. cAMP signaling in skeletal muscle adaptation: Hypertrophy, metabolism, and regeneration. *Am J Physiol Endocrinol Metab* **303**: E1–E17.
 Bergstrom DA, Penn BH, Strand A, Perry RL, Rudnicki MA, Tapscott SJ. 2002. Promoter-specific regulation of MyoD

binding and signal transduction cooperate to pattern gene expression. *Mol Cell* **9**: 587–600.
 Brand M, Rampalli S, Chaturvedi CP, Dilworth FJ. 2008. Analysis of epigenetic modifications of chromatin at specific gene loci by native chromatin immunoprecipitation (N-ChIP) of nucleosomes isolated using hydroxyapatite chromatography. *Nat Protoc* **3**: 398–409.
 Buljan M, Chalancon G, Eustermann S, Wagner GP, Fuxreiter M, Bateman A, Babu MM. 2012. Tissue-specific splicing of disordered segments that embed binding motifs rewires protein interaction networks. *Mol Cell* **46**: 871–883.
 Chaturvedi CP, Somasundaram B, Singh K, Carpenedo RL, Stanford WL, Dilworth FJ, Brand M. 2012. Maintenance of gene silencing by the coordinate action of the H3K9 methyltransferase G9a/KMT1C and the H3K4 demethylase Jarid1a/KDM5A. *Proc Natl Acad Sci* **109**: 18845–18850.
 Chen AE, Ginty DD, Fan CM. 2005. Protein kinase A signalling via CREB controls myogenesis induced by Wnt proteins. *Nature* **433**: 317–322.
 Demers C, Chaturvedi CP, Ranish JA, Juban G, Lai P, Morle F, Aebersold R, Dilworth FJ, Groudine M, Brand M. 2007. Activator-mediated recruitment of the MLL2 methyltransferase complex to the β -globin locus. *Mol Cell* **27**: 573–584.
 Du M, Perry RL, Nowacki NB, Gordon JW, Salma J, Zhao J, Aziz A, Chan J, Siu KW, McDermott JC. 2008. Protein kinase A represses skeletal myogenesis by targeting myocyte enhancer factor 2D. *Mol Cell Biol* **28**: 2952–2970.
 The ENCODE Project Consortium, Dunham I, Kundaje A, Aldred SF, Collins PJ, Davis CA, Doyle F, Epstein CB, Frietze S, Harrow J, et al. 2012. An integrated encyclopedia of DNA elements in the human genome. *Nature* **489**: 57–74.
 Fong AP, Yao Z, Zhong JW, Cao Y, Ruzzo WL, Gentleman RC, Tapscott SJ. 2012. Genetic and epigenetic determinants of neurogenesis and myogenesis. *Dev Cell* **22**: 721–735.
 Gabut M, Samavarchi-Tehrani P, Wang X, Slobodeniuc V, O'Hanlon D, Sung HK, Alvarez M, Talukder S, Pan Q, Mazzoni EO, et al. 2011. An alternative splicing switch regulates embryonic stem cell pluripotency and reprogramming. *Cell* **147**: 132–146.
 Haberland M, Arnold MA, McAnally J, Phan D, Kim Y, Olson EN. 2007. Regulation of HDAC9 gene expression by MEF2 establishes a negative-feedback loop in the transcriptional circuitry of muscle differentiation. *Mol Cell Biol* **27**: 518–525.
 Hinitz Y, Hughes SM. 2007. Mef2s are required for thick filament formation in nascent muscle fibres. *Development* **134**: 2511–2519.
 Iglesias-Bartolome R, Gutkind JS. 2011. Signaling circuitries controlling stem cell fate: To be or not to be. *Curr Opin Cell Biol* **23**: 716–723.
 Kalsotra A, Cooper TA. 2011. Functional consequences of developmentally regulated alternative splicing. *Nat Rev Genet* **12**: 715–729.
 Khokhlatchev A, Xu S, English J, Wu P, Schaefer E, Cobb MH. 1997. Reconstitution of mitogen-activated protein kinase phosphorylation cascades in bacteria. Efficient synthesis of active protein kinases. *J Biol Chem* **272**: 11057–11062.
 Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**: 357–359.
 Liu QC, Zha XH, Faralli H, Yin H, Louis-Jeune C, Perdiguero E, Prancaviciene E, Munoz-Canoves P, Rudnicki MA, Brand M et al. 2012. Comparative expression profiling identifies differential roles for myogenin and p38 α MAPK signaling in myogenesis. *J Mol Cell Biol* **4**: 386–397.
 Martin JF, Miano JM, Hustad CM, Copeland NG, Jenkins NA, Olson EN. 1994. A Mef2 gene that generates a muscle-specific

- isoform via alternative mRNA splicing. *Mol Cell Biol* **14**: 1647–1656.
- McNally JG, Muller WG, Walker D, Wolford R, Hager GL. 2000. The glucocorticoid receptor: Rapid exchange with regulatory sites in living cells. *Science* **287**: 1262–1265.
- Mukai A, Hashimoto N. 2008. Localized cyclic AMP-dependent protein kinase activity is required for myogenic cell fusion. *Exp Cell Res* **314**: 387–397.
- Ohkawa Y, Marfella CG, Imbalzano AN. 2006. Skeletal muscle specification by myogenin and Mef2D via the SWI/SNF ATPase Brg1. *EMBO J* **25**: 490–501.
- Ohkawa Y, Yoshimura S, Higashi C, Marfella CG, Dacwag CS, Tachibana T, Imbalzano AN. 2007. Myogenin and the SWI/SNF ATPase Brg1 maintain myogenic gene expression at different stages of skeletal myogenesis. *J Biol Chem* **282**: 6564–6570.
- Palii CG, Perez-Iratxeta C, Yao Z, Cao Y, Dai F, Davison J, Atkins H, Allan D, Dilworth FJ, Gentleman R, et al. 2011. Differential genomic targeting of the transcription factor TAL1 in alternate haematopoietic lineages. *EMBO J* **30**: 494–509.
- Penn BH, Bergstrom DA, Dilworth FJ, Bengal E, Tapscott SJ. 2004. A MyoD-generated feed forward circuit temporally patterns gene expression during skeletal muscle differentiation. *Genes Dev* **18**: 2348–2353.
- Potthoff MJ, Olson EN. 2007. MEF2: A central regulator of diverse developmental programs. *Development* **134**: 4131–4140.
- Potthoff MJ, Wu H, Arnold MA, Shelton JM, Backs J, McAnally J, Richardson JA, Bassel-Duby R, Olson EN. 2007. Histone deacetylase degradation and MEF2 activation promote the formation of slow-twitch myofibers. *J Clin Invest* **117**: 2459–2467.
- Rampalli S, Li L, Mak E, Ge K, Brand M, Tapscott SJ, Dilworth FJ. 2007. p38 MAPK signaling regulates recruitment of Ash2L-containing methyltransferase complexes to specific genes during differentiation. *Nat Struct Mol Biol* **14**: 1150–1156.
- Reed HC, Hoare T, Thomsen S, Weaver TA, White RA, Akam M, Alonso CR. 2010. Alternative splicing modulates Ubx protein function in *Drosophila melanogaster*. *Genetics* **184**: 745–758.
- Seenundun S, Rampalli S, Liu QC, Aziz A, Palii C, Hong S, Blais A, Brand M, Ge K, Dilworth FJ. 2010. UTX mediates demethylation of H3K27me3 at muscle-specific genes during myogenesis. *EMBO J* **29**: 1401–1411.
- Shevchenko A, Tomas H, Havlis J, Olsen JV, Mann M. 2006. In-gel digestion for mass spectrometric characterization of proteins and proteomes. *Nat Protoc* **1**: 2856–2860.
- Stewart R, Flechner L, Montminy M, Berdeaux R. 2011. CREB is activated by muscle injury and promotes muscle regeneration. *PLoS ONE* **6**: e24714.
- Trapnell C, Pachter L, Salzberg SL. 2009. TopHat: Discovering splice junctions with RNA-Seq. *Bioinformatics* **25**: 1105–1111.
- Trapnell C, Williams BA, Pertea G, Mortazavi A, Kwan G, van Baren MJ, Salzberg SL, Wold BJ, Pachter L. 2010. Transcript assembly and quantification by RNA-Seq reveals unannotated transcripts and isoform switching during cell differentiation. *Nat Biotechnol* **28**: 511–515.
- Trapnell C, Roberts A, Goff L, Pertea G, Kim D, Kelley DR, Pimentel H, Salzberg SL, Rinn JL, Pachter L. 2012. Differential gene and transcript expression analysis of RNA-seq experiments with TopHat and Cufflinks. *Nat Protoc* **7**: 562–578.
- Voss TC, Schiltz RL, Sung MH, Yen PM, Stamatoyannopoulos JA, Biddie SC, Johnson TA, Miranda TB, John S, Hager GL. 2011. Dynamic exchange at regulatory elements during chromatin remodeling underlies assisted loading mechanism. *Cell* **146**: 544–554.
- Wang ET, Sandberg R, Luo S, Khrebukova I, Zhang L, Mayr C, Kingsmore SF, Schroth GP, Burge CB. 2008. Alternative isoform regulation in human tissue transcriptomes. *Nature* **456**: 470–476.
- Wang Z, Zang C, Cui K, Schones DE, Barski A, Peng W, Zhao K. 2009. Genome-wide mapping of HATs and HDACs reveals distinct functions in active and inactive genes. *Cell* **138**: 1019–1031.
- Weil PA, Luse DS, Segall J, Roeder RG. 1979. Selective and accurate initiation of transcription at the Ad2 major late promoter in a soluble system dependent on purified RNA polymerase II and DNA. *Cell* **18**: 469–484.
- Yaffe D, Saxel O. 1977. Serial passaging and differentiation of myogenic cells isolated from dystrophic mouse muscle. *Nature* **270**: 725–727.

Tissue-specific splicing of a ubiquitously expressed transcription factor is essential for muscle differentiation

Soji Sebastian, Hervé Faralli, Zizhen Yao, et al.

Genes Dev. 2013, **27**: originally published online May 30, 2013
Access the most recent version at doi:[10.1101/gad.215400.113](https://doi.org/10.1101/gad.215400.113)

Supplemental Material

<http://genesdev.cshlp.org/content/suppl/2013/05/29/gad.215400.113.DC1>

References

This article cites 42 articles, 12 of which can be accessed free at:
<http://genesdev.cshlp.org/content/27/11/1247.full.html#ref-list-1>

License

Freely available online through the Genes & Development Open Access option.

Email Alerting Service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).