REVIEW

Pioneer transcription factors in cell reprogramming

Makiko Iwafuchi-Doi and Kenneth S. Zaret

Institute for Regenerative Medicine, Department of Cell and Developmental Biology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA

A subset of eukaryotic transcription factors possesses the remarkable ability to reprogram one type of cell into another. The transcription factors that reprogram cell fate are invariably those that are crucial for the initial cell programming in embryonic development. To elicit cell programming or reprogramming, transcription factors must be able to engage genes that are developmentally silenced and inappropriate for expression in the original cell. Developmentally silenced genes are typically embedded in “closed” chromatin that is covered by nucleosomes and not hypersensitive to nuclease probes such as DNase I. Biochemical and genomic studies have shown that transcription factors with the highest reprogramming activity often have the special ability to engage their target sites on nucleosomal DNA, thus behaving as “pioneer factors” to initiate events in closed chromatin. Other reprogramming factors appear dependent on pioneer factors for engaging nucleosomes and closed chromatin. However, certain genomic domains in which nucleosomes are occluded by higher-order chromatin structures, such as in heterochromatin, are resistant to pioneer factor binding. Understanding the means by which pioneer factors can engage closed chromatin and how heterochromatin can prevent such binding promises to advance our ability to reprogram cell fates at will and is the topic of this review.

Cell fate control represents the most extreme form of gene regulation. Genes specific to the function of a particular type of cell may be antagonistic to the function of another type of cell, so nature has evolved diverse regulatory mechanisms to ensure stable patterns of gene activation and repression. In prokaryotes, self-sustaining regulatory networks of transcription factors bound to DNA can be sufficient to regulate gene activation and repression (Ptashne 2011). In eukaryotes, ~200-base-pair (bp) segments of the genome are wound nearly twice around an octamer of the four core histones to form nucleosomes, providing steric constraints on how transcription factors

can bind DNA (Kornberg and Lorch 1999). As described below, pioneer transcription factors have the special property of being able to overcome such constraints, enabling the factors to engage closed chromatin that is not accessible by other types of transcription factors (Fig. 1). However, further higher-order packaging of nucleosomes into heterochromatin can occlude access to DNA altogether, presenting an additional barrier to cell fate conversion (Fig. 1). This review focuses on the most recent studies of pioneer factors in cell programming and reprogramming, how pioneer factors have special chromatin-binding properties, and facilitators and impediments to chromatin binding. We project that such knowledge will greatly aid future efforts to change the fates of cells at will for research, diagnostic, and therapeutic purposes.

Who's on first?

Which transcription factors are the first to access a developmentally silent gene and initiate its expression to promote a cell fate change? Genetic studies alone cannot answer this, as a transcription factor may appear necessary or sufficient to elicit a regulatory change but dependent on the prior binding of other factors in the cell. Why does knowing “who's on first” matter? Because the hierarchical mechanisms by which transcription factors engage target sites in chromatin provide insights into ways to modulate the process and hence cell fate control.

In diverse contexts where groups of transcription factors have been tested for their ability to convert cell fate, a subset of factors consistently has the greatest effect in cell conversion. Biochemical and genomic studies indicate that such factors can be considered “pioneers” by virtue of their ability to engage target DNA sites in closed chromatin prior to the apparent engagement, opening, or modification of the site by other factors (Box 1). Nucleosome binding by pioneer factors typically enables the coordinate or subsequent binding of other transcription factors, cofactors, and chromatin-modifying and remodeling enzymes, culminating in the activation of genes of a

[**Keywords:** pioneer transcription factor; reprogramming; nucleosome; chromatin; transdifferentiation; development]

Corresponding author: zaret@upenn.edu

Article is online at <http://www.genesdev.org/cgi/doi/10.1101/gad.253443.114>. Freely available online through the *Genes & Development* Open Access option.

© 2014 Iwafuchi-Doi and Zaret This article, published in *Genes & Development*, is available under a Creative Commons License (Attribution 4.0 International), as described at <http://creativecommons.org/licenses/by/4.0>.

Figure 1. Initial targeting of closed chromatin by pioneer factors. The DNA-binding domain of pioneer factors allows the protein to recognize its target site on nucleosomal DNA. The initial targeting of nucleosomal DNA by pioneer factors occurs in closed, silent chromatin that lacks nuclease hypersensitivity and consistent histone modifications or other prior marks. This allows the pioneer factor to initiate reprogramming of silent genes that may be inappropriate to express in a given cell, enabling cell type conversion. Many transcription factors (nonpioneers) cannot initially target such genes but can do so coordinately with, or after, pioneer factors bind. However, certain heterochromatic regions of the genome, such as where H3K9me2 or H3K9me3 marks are deposited, are refractory to pioneer factor binding. Continuing research on how pioneer factors can target nucleosomal sites and how heterochromatic impediments can be broken down will inform ways to enhance our ability to control cell fates for biomedical purposes.

new cell fate (Fig. 2). The main point is that the nucleosome-binding activity of pioneer factors allows them to initiate regulatory events at particular sites in chromatin that have not been programmed for expression, as typically seen in developmentally silent genes.

We start by reviewing the latest genome-mapping studies of chromatin states before and after transcription factor engagement in the contexts of development and cell reprogramming, which provide evidence that pioneer factors have special chromatin-binding properties suitable for their pioneering function. We then review the molecular mechanisms that underlie chromatin binding, the limits of such binding, and the importance of overcoming chromatin impediments in order to control cell fate.

Pioneer factors in zygotic genome activation

Maternal factors in the oocyte trigger zygotic genome activation, which is perhaps the most dramatic reprogramming event in embryogenesis (Tadros and Lipshitz 2009). The maternal transcription factor Zelda (Zld, zinc finger early *Drosophila* activator) plays a primary role for the onset of zygotic genome activation in *Drosophila* embryos (Liang et al. 2008; Nien et al. 2011). Zld protein is present in nuclei considerably earlier than other key maternal transcription factors such as Bicoid (Bcd) and Dorsal (Dl) proteins and is bound to gene regulatory regions prior to zygotic genome activation (Harrison et al. 2011; Nien et al.

2011). Zld binding increases DNA accessibility and facilitates the binding of other transcription factors, including Bcd and Dl, to target enhancers (Foo et al. 2014; Xu et al. 2014). Furthermore, differential DNA accessibility established by different levels of Zld binding sets the threshold for responding to the Dl gradient: More open enhancers are activated even where Dl concentration is low, but fewer open enhancers are activated only where Dl concentration is high (Foo et al. 2014). While in vitro nucleosome-binding studies have yet to be reported for Zld, by all in vivo criteria, Zld appears to function as a pioneer factor for zygotic genome activation.

Although homologs of Zld have not been reported outside the insect clade, in zebrafish, Nanog, Pou5f3 (originally named Pou2 and Pou5f1; a member of the class V POU family, as is mammalian Oct3/4), and the functionally redundant SoxB1 group of transcription factors (Sox2, Sox3, Sox19a, and Sox19b) are highly enriched and bound to their target sites prior to zygotic genome activation. They also play primary roles in the onset of zygotic genome activation (Lee et al. 2013; Leichsenring et al. 2013). In mice, maternal Oct3/4 and Sox2 are also primary regulators of zygotic genome activation (Foygel et al. 2008; Pan and Schultz 2011). Altogether, factors that activate the zygotic genome can engage their target sites in chromatin that is not preprogrammed, can elicit local chromatin changes, and can enable subsequent gene expression, thus having the hallmarks of pioneer transcription factors (Table 1).

Box 1. Features of pioneer transcription factors

Engage their target sites in closed (nuclease-resistant), silent chromatin prior to gene activity.

Increase accessibility of a target site that makes other proteins (e.g., transcription factors, chromatin remodelers, chromatin modifiers, histone variants, and repressors) accessible to the site.

Play a primary role in cell programming and reprogramming and establish the competence for cell fate changes.

How to predict/validate pioneer factors

Observe the chromatin state of target sites for a transcription factor before and after the factor is expressed in a cell. Pioneer factors can target sites that are closed (nuclease-resistant; e.g., with ATAC-seq and DNase-seq) or shown to be nucleosomal (MNase-Seq) prior to binding and often result in chromatin opening upon binding. Merely correlating open chromatin at sites where a transcription factor is already bound does not predict pioneer factors.

Analyze direct binding between a transcription factor and a reconstituted mononucleosome or a nucleosomal array in vitro (e.g., with electrophoretic mobility shift assays, DNase I footprinting, and sequential transcription factor and core histone ChIP).

Analyze the effect of a transcription factor binding on DNA accessibility at the target sites in vitro (reconstituted nucleosomal array with a transcription factor) or in vivo (ectopic expression or deletion of a transcription factor).

Figure 2. Initial targeting of pioneer factor and subsequent events. (A) Pioneer transcription factors can target sites with high intrinsic nucleosome occupancy. (B) Pioneer factors initially engage nucleosomal target sites, which enable other factors (transcription factors, chromatin modifiers, and remodelers) to access the target sites. (C) Other transcription factor (TF)-binding and chromatin modifications could stabilize pioneer factor binding to the target sites.

Pioneer factors in cell reprogramming

Reprogramming of terminally differentiated cells was first shown by somatic cell nuclear transfer into enucleated oocytes (Gurdon 1962), indicating that factors in the oocyte cytoplasm can reprogram somatic nuclei to a pluripotent state. By screening diverse factors that are normally expressed in pluripotent stem cells (PSCs), but not fibroblasts, for their ability to convert fibroblasts to pluripotency, the transcription factors Oct3/4, Sox2, Klf4, and c-Myc (O, S, K, and M) were found to trigger endogenous expression of pluripotent factors and be sufficient to reprogram fibroblasts into induced PSCs (iPSCs) (Takahashi and Yamanaka 2006). Although alternative sets of transcription factors for iPSC reprogramming have been reported (e.g., Buganim et al. 2014), most studies include Oct3/4 and/or Sox2 (Yu et al. 2007; Feng et al. 2009; Han et al. 2010; Gao et al. 2013). How does this set of factors first interact with their target sites to initiate reprogramming? A snapshot of the initial binding events of OSKM in human fibroblasts indicates their preferential occupancy of promoter-distal (i.e., enhancer) target sites (Soufi et al. 2012). Many initial binding events occur at genes that elicit reprogramming to pluripotency as well as at genes that promote apoptosis during the early stages of iPSC reprogramming. Many more initial binding events are distinct from the definitive binding pattern in embryonic stem (ES) cells or iPSCs that maintains pluripotency (Soufi et al. 2012). Thus, the initial binding or scanning

of the genome is quite promiscuous during reprogramming of somatic cells to pluripotency, and subsequent reorganization of the factors in the genome must occur to establish the final pluripotent state (Hochedlinger and Plath 2009).

Of these factors, Oct3/4, Sox2, and Klf4 act as pioneer transcription factors in that they can access closed chromatin whether they bind together or alone (Fig. 1; Table 1; Soufi et al. 2012). “Closed chromatin” in this context means to lack DNase hypersensitivity and a lack of a consistent histone modification pattern. In contrast, c-Myc alone prefers to bind to “open chromatin” sites that are DNase-hypersensitive and contain activating histone modifications (Soufi et al. 2012). However, c-Myc can bind closed chromatin sites in conjunction with the other factors (Soufi et al. 2012). These findings illustrate how a cohort of factors, all of which are necessary in genetic tests of efficient conversion to pluripotency, can differ markedly with regard to whether they are pioneer factors. In addition, the pioneer factors can directly enable non-pioneers to engage closed chromatin sites. The exact mechanisms by which such facilitation occurs remain to be determined but could include interactions between the pioneer factors and the nucleosomes that enable other factors to bind, direct interactions between the factors themselves, and/or recruitment of cofactors or nucleosome remodelers that in turn alter the local chromatin landscape to facilitate secondary factor binding (Fig. 2).

Single-molecule tracking studies in live cells suggested that during iPSC reprogramming, Sox2 is bound to the target DNA first, followed by binding of Oct3/4 to stabilize the ternary complex (Chen et al. 2014). These live-cell results do not match the ChIP-seq (chromatin immunoprecipitation [ChIP] combined with deep sequencing) data, where the majority of the initial Oct3/4- and Sox2-binding events were found to be independent of one another (Soufi et al. 2012). The ChIP-seq studies with native OSKM proteins were performed under conditions where continued culture of the OSKM-induced cells led to iPSC conversion. In contrast, the single-molecule studies required apparently higher ectopic expression of HALO-tagged factors, which covalently bind a fluorescent ligand in short growth period experiments. It could be that the single-molecule studies were revealing chromatin state-dependent binding for the factors to engage DNase-sensitive, apparently free DNA sites in the nucleus, which represented a minority of events in the ChIP-seq studies. We need direct nucleosome- and chromatin-binding studies to assess exactly how the OSKM factors mechanistically and sequentially engage chromatin to initiate the dramatic reprogramming to pluripotency. That the Pou5f3 and Sox2 factors can serve a pioneering function during zygotic genome activation (see above) underscores the special abilities of these factors.

Pioneer factors in direct cell conversion (transdifferentiation)

Direct cell conversion (transdifferentiation) by individual transcription factors was shown for the first time when MyoD was found to induce muscle-specific properties

Table 1. Predicted/validated pioneer transcription factors

Pioneer factors	Cellular/biochemical context (model organisms)	Predicted/validated pioneer activity	References
FoxA	Transdifferentiation from fibroblast to iHep (mouse)	Inducing transdifferentiation	Huang et al. 2011; Sekiya and Suzuki 2011
	Liver/foregut development (mouse/ <i>Caenorhabditis elegans</i>)	Establishment of the competence for liver development	Lee et al. 2005
	Mitotic bookmarking (human)	Occupying silent liver enhancer in endoderm progenitors	Gualdi et al. 1996
	Hormone-dependent cancer progression (human)	Recruitment of histone variant H2A.Z Chromatin decompaction	Updike and Mango 2006 Fakhouri et al. 2010
	In vitro reconstituted nucleosome or nucleosome array	Binding to mitotic chromatin Recruitment of hormone receptor	Caravaca et al. 2013 Zaret and Carroll 2011; Jozwik and Carroll 2012
Zelda		Binding to mononucleosomes, dinucleosomes, and nucleosome array independent of other factors in vitro	Cirillo and Zaret 1999; Cirillo et al. 2002
	Zygotic genome activation (<i>Drosophila</i>)	Increasing DNA accessibility in compacted nucleosome array independent of other factors in vitro Inducing zygotic genome activation	Liang et al. 2008 Harrison et al. 2011; Foo et al. 2014; Xu et al. 2014
	Zygotic genome activation (zebrafish/mouse)	Increasing DNA accessibility in native chromatin Inducing zygotic genome activation	Foygel et al. 2008; Lee et al. 2013
Class V Pou (e.g., Oct3/4, Pou5f3)		Binding to target sites before zygotic genome activation Inducing reprogramming	Leichsenring et al. 2013 Takahashi and Yamanaka 2006
	Reprogramming from fibroblast to iPSCs (mouse/human)	Binding to chromatin that is not DNase-sensitive and not premarked by common histone modifications in vivo	Soufi et al. 2012
	Zygotic genome activation (zebrafish/mouse)	Inducing zygotic genome activation	Pan and Schultz 2011; Lee et al. 2013
Group B1 Sox (e.g., Sox2)	Reprogramming from fibroblast to iPSCs (mouse/human)	Inducing reprogramming	Takahashi and Yamanaka 2006
		Binding to chromatin that is nonhypersensitive and not premarked by common histone modifications in vivo	Soufi et al. 2012
	Cancer progression (mouse/human)	Inducing tumor formation	Bass et al. 2009; Boumahdi et al. 2014
Klf4	Reprogramming from fibroblast to iPSCs (mouse/human)	Inducing reprogramming	Takahashi and Yamanaka 2006
		Binding to chromatin that is nonhypersensitive and not premarked by common histone modifications in vivo	Soufi et al. 2012

(continued)

Table 1. Continued

Pioneer factors	Cellular/biochemical context (model organisms)	Predicted/validated pioneer activity	References
Ascl1	Transdifferentiation from fibroblast/hepatocyte to iN cells (mouse/human)	Inducing transdifferentiation	Vierbuchen et al. 2010
Pax7	Pituitary melanotrope development	Binding to nonhypersensitive chromatin in vivo	Caiazzo et al. 2011; Son et al. 2011; Wapinski et al. 2013
PU.1	Pituitary adenomas—Cushing's disease Transdifferentiation from fibroblast to macrophages (mouse) Myeloid and lymphoid development (mouse/human)	Establishment of the competence for melanotrope development Increasing DNA accessibility in native chromatin Inducing transdifferentiation	Budry et al. 2012; Drouin 2014
GATA4	Transdifferentiation from fibroblast to iHep (mouse) Liver development (mouse) Hormone-dependent cancer progression (human)	Increasing DNA accessibility in native chromatin Inducing transdifferentiation	Feng et al. 2008 Ghisletti et al. 2010; Heinz et al. 2010; Barozzi et al. 2014 Huang et al. 2011
GATA1 CLOCK:BMAL1 p53	In vitro reconstituted nucleosome array Mitotic bookmarking (mouse) Circadian clock (mouse) Tumor suppression by regulating cell cycle progression and apoptosis (human)	Occupying silent liver enhancer in endoderm progenitors Recruitment of hormone receptor Binding to nucleosome array independent of other factors in vitro (not as effective as FoxA) Increasing DNA accessibility in compacted nucleosome array independent of other factors in vitro (not as effective as FoxA) Binding to mitotic chromatin Increasing DNA accessibility in native chromatin Increasing DNA accessibility in native chromatin	Bossard and Zaret 1998 Zaret and Carroll 2011; Jozwik and Carroll 2012 Cirillo et al. 2002 Kadauke et al. 2012 Menet et al. 2014 Laptenko et al. 2011
	In vitro reconstituted mononucleosome or nucleosome array	Binding to chromatin that encodes high intrinsic nucleosome occupancy Binding to nucleosome independent of other factors in vitro Recruitment of histone acetyltransferase p300	Lidor Nili et al. 2010 Espinosa and Emerson 2001

when ectopically expressed in fibroblasts (Davis et al. 1987). However, exogenous MyoD in endodermal and ectodermal cells failed to induce a full phenotypic switch (Weintraub et al. 1989; Schäfer et al. 1990). Later studies revealed that MyoD induces *myogenin* through cooperation with the Pbx factor, which is constitutively bound to MyoD target sites in fibroblasts prior to MyoD expression (Berkas et al. 2004). There are many other examples of direct cell conversion within the same germ layer: the combination of PU.1 and C/EBP α into macrophage-like cells from fibroblasts (Feng et al. 2008) and GATA4, MEF2C, TBX5, Hand2, and NKX2.5 for cardiomyocyte-like cells from fibroblasts (Ieda et al. 2010; Addis et al. 2013).

The first evidence of efficient transdifferentiation across different germ layers was the generation of functional glutaminergic neurons (inducible neurons [iNs]) from fibroblasts by the three transcription factors Ascl1, Brn2, and Myt1l (Vierbuchen et al. 2010). Of these factors, Ascl1 plays a central role to initiate transdifferentiation because Ascl1 alone is sufficient to induce immature iN cells, but Brn2 and Myt1l are not. Soon after, dopaminergic and motor neurons were generated from fibroblasts using different sets of transcription factors, but both sets included Ascl1 (Caiazzo et al. 2011; Son et al. 2011). Recently, Wernig and colleagues (Wapinski et al. 2013) revealed a hierarchical mechanism governing the early stage of transdifferentiation to iNs: Ascl1 acts as a pioneer transcription factor to bind closed chromatin and recruit Brn2 to Ascl1 target sites (Table 1), and Brn2 is primarily required for the later stage of transdifferentiation by contributing to iN maturation. These experiments show that even when the factors are simultaneously overexpressed, they function in a hierarchical manner, as described above for Oct3/4, Sox2, Klf4, and c-Myc. Ascl1 acts first to establish competence for the neuronal lineage, followed by other transcription factors to specify neuronal subtypes. Like the nonpioneer c-Myc, Ascl1 is a basic helix-loop-helix (bHLH) family transcription factor whose core structure is thought to bind to both sides of the DNA helix (Bertrand et al. 2002). However, unlike c-Myc, Ascl1 appears to have properties of a pioneer factor. It will be important to determine how different members of the bHLH class, which exhibit differences in how they contact DNA, can or cannot function as pioneer factors.

Furthermore, Ascl1 was characterized as an “on-target” pioneer factor in that most of its initially bound sites, when ectopically expressed in fibroblasts, correspond to sites that remain bound when the cells finally differentiate into neurons (Wapinski et al. 2013). This in contrast to Oct3/4, Sox2, and Klf4 initially binding to the genome in fibroblasts, where, as described above, many of the initial binding events do not correspond to those after full conversion to iPSCs (Soufi et al. 2012). This could reflect inherent differences in how Ascl1 versus Oct3/4, Sox2, and Klf4 initially recognize their target sites in chromatin.

Alternatively, there could be an inherent difference in the overall chromatin structure of pluripotent cells that takes multiple phases to recapitulate by reprogramming

to iPSCs, whereas direct conversion among differentiated cells may not involve as dramatic an overall chromatin change. For example, fluorescence recovery after photobleaching experiments have suggested that pluripotent cell chromatin is more “hyperdynamic” than somatic cell chromatin, allowing for a more rapid exchange of chromatin components than in differentiated cells (Meshorer et al. 2006). Thus, global changes in chromatin structure may be necessary for reprogramming to iPSCs, and hence cells may have to go through various stages with different binding patterns at each stage.

Another example of transdifferentiation across germ layers is when hepatocyte-like cells are induced from fibroblasts (iHep) by exogenous expression of either forkhead box protein A1 (FoxA1), FoxA2, or FoxA3 with HNF4a (Sekiya and Suzuki 2011) or exogenous expression of FoxA3, GATA4, and HNF1a in combination with the inactivation of the tumor suppressor gene p19^{Arf} (Huang et al. 2011). These factors were discovered from pools of many transcription factors involved in liver development; the common component was FoxA. However, compared with primary hepatocytes, iHeps show significant differences in gene expression and are unable to fully rescue liver function in a transplantation model (Huang et al. 2011; Sekiya and Suzuki 2011). CellNet analysis, which is a bioinformatics tool to assess the fidelity of cell conversion, surprisingly revealed that iHeps induced by FoxA and HNF4a represent gut endoderm progenitors more than mature hepatocytes (Morris et al. 2014). They also show that iHeps are capable of engrafting both the mouse liver and colon, which are alternative fates from gut endoderm (Bort et al. 2006). It appears that the pioneer transcription factor FoxA and cofactors establish the competence to differentiate gut endodermal lineage from fibroblasts (Table 1), and additional lineage-determining factors are required for making mature hepatocytes. This model fits well with the original description of FoxA as endowing competence for the endoderm in embryos (Gualdi et al. 1996).

CellNet analyses further revealed that most transdifferentiation events fail to extinguish the expression programs of the starting cell type and thus impair the fidelity of cell type conversion (Cahan et al. 2014). Therefore, it is important to know both the hierarchical regulatory network required to activate a desired cell type and the means to extinguish the original gene regulatory program. This latter area appears to be a major gap in the field of cell type conversion. Along these lines, it would be important to know how to reset tissue-specific heterochromatic blocks that can be resistant to pioneer factor binding, as discussed below, and serve as an impediment to cell reprogramming (Soufi et al. 2012).

Pioneer factors in cell programming during development

Not surprisingly, the transcription factors that have the strongest effect on cell reprogramming and transdifferentiation have been shown to play central roles in embryonic development. FoxA1 and FoxA2 are expressed in foregut endoderm and are required for the establish-

ment of competence for liver development (Gualdi et al. 1996; Lee et al. 2005). GATA4 and GATA6 are expressed in foregut endoderm and are required for early liver development (Holtzinger and Evans 2005; Zhao et al. 2005; Watt et al. 2007). The liver-specific enhancer of the *albumin* (*Alb1*) gene contains six transcription factor-binding sites (McPherson et al. 1993). Of these, only binding sites for FoxA and GATA are occupied in the beginning of gut endoderm development, in which the *Alb1* gene has not yet been expressed (Gualdi et al. 1996; Bossard and Zaret 1998; Xu et al. 2012). Upon liver specification, all binding sites are occupied, and the *Alb1* gene becomes active. Thus, the FoxA and GATA factors act as pioneers for liver specification by engaging chromatin in multipotent progenitors and helping to provide the competence for hepatic fate (Table 1). During ES cell differentiation into endoderm in vitro, some FoxA2-binding sites are preoccupied by H2A.Z, and FoxA2 binding at the sites during endoderm induction results in nucleosome depletion and gene activation (Li et al. 2012a).

Hierarchical binding, in which pioneer factors bind first, is not unique to liver development. In pituitary melanotrope development, the initial expression of Pax7, a paired homeodomain transcription factor, slightly precedes the other melanotrope transcription factors, such as Tpit (Budry et al. 2012). Budry et al. (2012) exogenously expressed Pax7 in a pituitary corticotrope cell line in which Tpit is expressed but Pax7 is not. Exogenous Pax7 binding increased chromatin accessibility and induced de novo Tpit recruitment at the Pax7-binding sites. Pax7 is bound to DNA with either or both motifs for the protein's paired domain and its homeodomain. Interestingly, Pax7 binding with both motifs gives more chromatin accessibility than those with a single motif. Although the intrinsic mechanism is not understood, Pax7 opens chromatin structure through the binding to composite motifs and provides competence for melanotrope development (Table 1; Drouin 2014).

PU.1 has been known to play a crucial role in myeloid and lymphoid development, and its expression there precedes other lineage-determining transcription factors (Carotta et al. 2010). PU.1 binding induces local nucleosome remodeling followed by the deposition of the active histone modification H3K4me1 (Heinz et al. 2010). PU.1 also engages other lineage-determining transcription factors (Heinz et al. 2010) and promotes overall nucleosome depletion (Barozzi et al. 2014). Additionally, PU.1 binding contributes to endotoxin-induced transcriptional activation in macrophages (Ghisletti et al. 2010). PU.1 is bound to a substantial fraction of the endotoxin-inducible enhancers prior to endotoxin stimulation to keep the enhancers accessible. Exogenous PU.1 expression in fibroblasts also induces accessible chromatin at de novo PU.1-binding sites in the inducible enhancers. PU.1 uses hydration to recognize target DNA and forms a long-lived complex relative to the other Ets factors, which use electrostatic interactions (Wang et al. 2014). We speculate that the hydration-based recognition imparts the necessary adaptability for PU.1 to bind nucleosomal DNA. Although more mechanistic studies are required to understand how PU.1 exerts its function, PU.1 can act as

a pioneer factor to confer accessibility to the target enhancers during myeloid and lymphoid development (Table 1). Taken together, these studies show that pioneer factor binding occurs prior to lineage commitment and can employ a chromatin-opening step to establish the competence for gene activation.

Extensive computational analyses of changes in DNase cleavages across the genome have been studied during the in vitro differentiation of ES cells to mesendoderm, endoderm, prepancreatic, and intestinal endoderm fates (Sherwood et al. 2014). Sherwood et al. (2014) developed an elegant method called protein interaction quantitation (PIQ) that predicts transcription factor binding based on DNase footprinting and intrinsic binding motifs. PIQ then analyzes genome-wide correlations between a factor's binding and the extent of local chromatin accessibility and thereby may predict pioneer transcription factors.

However, PIQ did not predict FoxA as a pioneer factor. As Sherwood et al. (2014) discussed, it could be because FoxA can recruit corepressors that create closed chromatin (Sekiya and Zaret 2007); such binding events by FoxA would not be scored because they would not be predicted to cause the hypersensitivity that is necessary for PIQ to identify pioneer activity. In addition, the PIQ method predicted only 50% of a factor's actual binding sites detected by ChIP-seq.

Despite these issues, it is interesting that Sherwood et al. (2014) discovered that a subset of pioneer factors discerned by PIQ causes chromatin opening preferentially on one side of their nonpalindromic DNA-binding motif. That a cohort of factors behaved asymmetrically was striking and suggests directionally specific mechanisms to be discovered with regard to "downstream" events that follow pioneer factor binding.

Post-developmentally, there are many examples of where prior binding by FoxA factors enables the subsequent binding by hormone receptors and other DNA-binding proteins (Gao et al. 2003; Carroll et al. 2005; Zhang et al. 2005; Li et al. 2009, 2012b; Pihlajamaa et al. 2014), as previously covered in greater detail (Zaret and Carroll 2011). The mammalian circadian clock is driven by the CLOCK and BMAL1 transcription factors that rhythmically bind to nucleosomal target sites, promote incorporation of the histone variant H2A.Z, and enable the subsequent binding of other transcription factors (Menet et al. 2014). Thus, the mammalian clock appears to employ pioneer factors to activate genes at specific times of the daily cycle.

The basis for pioneer activity: nucleosome binding and chromatin opening

What are the molecular mechanisms underlying pioneer function? The FoxA family of transcription factors has been examined in mechanistic depth by biochemical analyses on mononucleosomal, dinucleosomal, and nucleosomal array substrates in vitro in conjunction with in vivo footprinting (Table 1; Cirillo et al. 2002; Zaret and Carroll 2011). Purified FoxA protein can bind to its target sites on nucleosomal DNA, open a local domain of compacted chromatin, and stabilize nucleosome position

(McPherson et al. 1996; Shim et al. 1998; Cirillo et al. 2002). Chromatin opening by FoxA in vitro does not require ATP or ATP-dependent chromatin remodelers (Cirillo et al. 2002). Notably, the DNA-binding domain of FoxA resembles that of linker histone and binds to one side of a DNA helix along the long axis of DNA, which leaves the other side of DNA to bind core histones (Clark et al. 1993; Ramakrishnan et al. 1993; Cirillo et al. 1998). Thus, the FoxA pioneer factor is structurally suited to recognize its DNA target sites on a nucleosome. Also, the C-terminal domain of FoxA can bind directly to core histone proteins and is required for opening chromatin (Cirillo et al. 2002). We need direct nucleosome-binding studies for other pioneer factors to determine how different DNA-binding domains may adapt to the nucleosome surface and how other domains of the proteins may interact directly with the core histones, perhaps destabilizing them. In addition, we need to understand how pioneer factors may engage chromatin-modifying enzymes to expand the “openness” of a local domain of chromatin, helping to activate an enhancer or a promoter.

Nucleosomes at enhancers: a collaborator for pioneer factors

In higher eukaryotes, gene expression is regulated by the coordinated action of promoters, which determine a gene's transcription start site, and one or more distal enhancers, which modulate promoter activity (Levine et al. 2014). These regulatory sequences occur in the context of nucleosomes, which provide a repressive ground state for gene expression (Struhl 1999; Wang et al. 2011). Nucleosome occupancy can be determined by multiple factors: DNA sequences, transcription factors, chromatin remodeling enzymes, and the transcriptional machinery (Struhl and Segal 2013). DNA sequences can favor or disfavor nucleosome formation. For example, sequences with poly(dA:dT) tracts are known to destabilize nucleosomes (Anderson and Widom 2001), whereas sequences with high GC content or ~10-bp periodicities of AA or TT dinucleotides favor nucleosome formation (Ioshikhes et al. 2006; Segal et al. 2006; Tillo and Hughes 2009). In higher eukaryotes, the DNA sequence at promoters, enhancers, and transcription factor-binding sites generally encode a high intrinsic nucleosome occupancy (Tillo et al. 2010; Gaffney et al. 2012). However, in disagreement with predicted high nucleosome occupancy, genome-wide mapping typically shows that nucleosomes are depleted at active promoters (Schones et al. 2008; Valouev et al. 2011; Teif et al. 2012). DNA accessibility studies further indicate that promoters tend to open ubiquitously among multiple cell types (Thurman et al. 2012), suggesting that generally expressed *trans*-factors keep promoters accessible. However, enhancers tend to become open in a tissue-specific manner (Thurman et al. 2012).

How do enhancer chromatin sites become accessible in a tissue-specific manner against an intrinsic propensity of the underlying DNA for high nucleosome occupancy? Although most transcription factors cannot access nucleosomal DNA by themselves, pioneer transcription factors

(e.g., FoxA) can bind to nucleosomal DNA and recruit other factors to bind (Cirillo et al. 2002). Along these lines, the transcription factors p53 and PU.1 and the progesterone receptor, which apparently operates with FoxA1 (Clarke and Graham 2012), preferentially target their binding motifs amid sequences with high intrinsic nucleosome occupancy rather than high-affinity motifs with low intrinsic nucleosome occupancy (Lidor Nili et al. 2010; Ballare et al. 2013; Barozzi et al. 2014). Thus, contrary to the dogma that nucleosomes would be inherently repressive to gene activity and must be removed for factor binding, pioneer transcription factors positively use the feature of high nucleosome occupancy at enhancers as their functional binding target (Fig. 2A).

Given that pioneer factors can bind their targets on nucleosomal DNA, where other factors cannot, we suggest that, at enhancers involved with cell and tissue identity, nucleosomes act as a filter to prevent binding of non-pioneer factors that prefer free DNA and would otherwise perturb a cell identity network. By this logic, if the target sequence became stably nucleosome-free, the nucleosome could no longer serve as such a filter. We need further studies of nucleosome stability at higher eukaryotic enhancers to assess these issues.

Facilitators and impediments for pioneer factor binding in vivo

Genome-wide mapping studies show that all transcription factors, including pioneer factors, bind to only a subset of their putative binding motifs in the genome and show distinct binding patterns in different cell types. As an example of the requisite analysis, see the application of the computational tool MultiGPS to compare ChIP-seq data sets (Mahony et al. 2014). The observations of Mahony et al. (2014) led to the hypothesis that (1) pioneer factors can initiate targeting of a region and then engage other factors to stabilize a pioneer factor's binding, (2) positive chromatin features can enable pioneer factor binding, and (3) repressive chromatin features can prevent pioneer factor binding. We discuss these possibilities below.

Nucleosome-mediated cooperativity between transcription factors

The binding of multiple transcription factors that function combinatorially is inherent to eukaryotic gene regulation, particularly for cell-specific control (Yamamoto 1985; Carey 1998). As described earlier, transdifferentiation events typically require sets of transcription factors. In liver development, the pioneer factors FoxA and GATAs initially occupy a liver-specific enhancer (Gualdi et al. 1996). In macrophage and B-cell development, PU.1 acts as a pioneer transcription factor but requires a distinct set of lineage-determining transcription factors to enhance PU.1 binding at target sites and the deposition of active histone modifications (Heinz et al. 2010). Furthermore, in the initiation of iPSC reprogramming, Oct3/4, Sox2, and Klf4 act as pioneer transcription factors, and c-Myc facilitates Oct3/4, Sox2, and Klf4 binding (Soufi et al. 2012).

These results suggest that pioneer transcription factors can bind coordinately with other factors. We speculate that the intrinsic nucleosome recognition properties of pioneer factors allows them to scan the closed chromatin for potential target sites and then recruit other factors, which could in turn stabilize the pioneer factor's binding to chromatin (Fig. 2).

The most commonly known cooperative binding is based on protein–protein interaction between transcription factors. However, the nucleosome structure of DNA enables cooperative binding without such protein–protein interactions; that is, nucleosome-mediated cooperativity between transcription factors can be achieved by their simultaneous competition with histones for the underlying DNA (Adams and Workman 1995; Chavez and Beato 1997; Vashee et al. 1998; Miller and Widom 2003; Mirny 2010; Moyle-Heyrman et al. 2011). Such cooperativity typically can be seen when multiple (e.g., four to five) transcription factor-binding events can occur within one or one-half of a nucleosome. In contrast, pioneer factors on their own can recognize nucleosomal target sites and then recruit other transcription factors. Such recruitments could stabilize the binding of a pioneer and a non-pioneer factor to elicit a cell type-specific binding pattern. With or without such stabilization, the enhanced ability of pioneer factors to interact with nucleosomal DNA provides an advantage for cooperative mechanisms to facilitate the binding of multiple transcription factors to closed chromatin at silent genes.

Positive chromatin features for pioneer factor binding

Do pioneer transcription factors positively use chromatin modifications to enhance their interaction with certain target sites? Here the data are mixed. First, recent studies indicate that ~40% of the mammalian and *Drosophila* genomes lack distinctive histone modifications and are considered to be in a “low signal” state (Kharchenko et al. 2011; Ho et al. 2014). Thus, other than being nucleosomal, a large fraction of chromatin appears to be neutral—not marked for either activation or repression. Consistent with such observations, human Oct3/4, Sox2, and Klf4 in initial reprogramming bind to genomic sites that are not enriched, on average, for histone modifications (Soufi et al. 2012), and PU.1 binding does not depend on H3K4me1 but is followed by H3K4me1 deposition (Ghisletti et al. 2010; Heinz et al. 2010).

Two independent studies show that the deposition of active histone modifications (H3K4me2, H3K4me1, and H3K9ac) follow or are concomitant with FoxA binding but are not prior to FoxA binding during differentiation induced by retinoic acid in pluripotent ES or P19 embryonal carcinoma cells (Taube et al. 2010; Serandour et al. 2011). However, another study showed that reduction of H3K4me1 and H3K4me2 impaired FoxA binding in the breast cancer cell line MCF7 (Lupien et al. 2008). Altogether, it appears that active histone modifications may enhance pioneer factor binding to chromatin but are not necessarily required for the factor's initial engagement. Also, DNA hypomethylation is not required for FoxA

binding (Serandour et al. 2011). In terms of histone variants, FoxA2-binding sites can be preoccupied by H2A.Z during ES cell differentiation to endoderm/hepatic progenitor cells, but these preoccupied sites are only a fraction (16%) of the total FoxA2-binding sites (Li et al. 2012a). At this point, the most consistent chromatin feature that predicts engagement of pioneer transcription factors is high intrinsic nucleosome occupancy (Fig. 2).

Repressive chromatin features for pioneer factor binding

Are there chromatin features that prevent a pioneer factor's binding to nonfunctional or alternative lineage sites? Comparing between MCF7 breast cancer and LNCaP prostate cancer cell lines, FoxA1 binds differentially to more than half of its target sites (Lupien et al. 2008). This cell type-specific binding is inversely correlated with the repressive histone modification H3K9me2. Furthermore, megabase-scale heterochromatic domains spanned by H3K9me3 in fibroblasts contain genes required for cell reprogramming to pluripotent cells but are refractory to the initial binding by OSKM during the reprogramming (Fig. 1; Soufi et al. 2012). A reduction of H3K9me3 deposition allows Oct3/4 and Sox2 binding to these domains and enhances reprogramming. Strikingly, various genes that function late in the reprogramming process, causing reprogramming to proceed more deterministically in the cell population, map to the heterochromatic domains that are resistant to the initial OSKM binding (Buganim et al. 2012). It thus appears that the H3K9me3 heterochromatic domains represent a barrier to the conversion from the stochastic phase to the deterministic phase of reprogramming. Altogether, H3K9me2 and H3K9me3 domains seem to block pioneer factor binding.

Heterochromatic blocks to pioneer factor binding may provide a means for cells to stably retain their fate. We suggest that a reason why transdifferentiation typically fails to shut off an initial genetic program may relate to the failure of resetting heterochromatic domains for new cell type (Cahan et al. 2014). Understanding how heterochromatic regions are established and maintained will continue to provide insight into enhancing cell conversions and may help explain changes in cell fate in diseases such as cancer (Cruickshanks et al. 2013).

Pioneer factors as bookmarking factors in mitosis

Pioneer transcription factors play an important role in cellular memory during mitotic growth. During mitosis, the chromosomes markedly condense, RNA polymerase is excluded from the chromatin, and transcription ceases. Recent chromosome conformation capture studies show that the three-dimensional partitioning of genomic domains in interphase is almost completely lost during mitosis (Naumova et al. 2013). During mitotic exit, such domains are re-established, RNA polymerase re-engages the chromatin, and the entire genome becomes transcriptionally activated in a manner that faithfully recapitulates the prereplicated state of the cell (Egli et al. 2008).

Although most transcription factors and RNA polymerase II dissociate from mitotic chromatin, a subset of transcription factors are retained, originally referred to as “bookmarking” factors (Martinez-Balbas et al. 1995; Michelotti et al. 1997; Zaidi et al. 2010). In the time since this initial discovery, it has become appreciated that the mitotic-retaining bookmarking factors are typically those classified as pioneer transcription factors, including GATA1 and FoxA1 (Table 1; Kadauke et al. 2012; Caravaca et al. 2013). Genome-wide binding studies of GATA1 and FoxA1 in mitosis show that only a minority of their interphase binding sites are retained in mitotic chromatin (Kadauke et al. 2012; Caravaca et al. 2013). Interestingly, the mitotic-retained sites of binding in GATA1 and FoxA1 are not associated with a particular histone modification. The one correlate that was observed is that the mitotic-retained FoxA1 sites are those with a higher predicted intrinsic nucleosome occupancy (Caravaca et al. 2013). This is consistent with the above discussion about how predicted nucleosome occupancy is a prominent feature at genomic regulatory sequences and underscores the nucleosome-binding pioneer feature for mitotic bookmarking.

Notably, the genes associated with GATA1 and FoxA1 binding in mitosis were among the first to be reactivated during mitotic exit (Kadauke et al. 2012; Caravaca et al. 2013). A careful experiment in which GATA1 was destroyed specifically in mitosis proved definitively that mitotic retention was crucial for timely gene reactivation during mitotic exit (Kadauke et al. 2012). All of the known bookmarking/pioneer factors function in cell programming during development, leading to the suggestion that the mechanism of gene reactivation during mitotic exit could recapitulate, in part, the hierarchical processes by which different cell types are specified in the first place (Zaret 2014).

Pioneer factors in disease progression

Since pioneer transcription factors so markedly affect cell reprogramming, their misregulation could cause severe effects on human health. *Sox2* is not expressed in the normal epidermis but is up-regulated in mouse and human cancer stem cells in skin squamous cell carcinomas (Boumahdi et al. 2014). There, *Sox2* is involved in the initial stage of tumor formation and its maintenance. In human esophageal and lung squamous cell carcinomas, chromosome segments containing *Sox2* are often genetically amplified (Bass et al. 2009). Thus, ectopic *Sox2* expression or a higher level of its expression could activate previously silent programs to establish competence for tumorigenesis, perhaps analogous to cell reprogramming. Furthermore, FoxA and GATA factors are involved in a variety of hormone-dependent cancers, such as estrogen receptor-positive breast cancer and androgen receptor-positive prostate cancer (Zaret and Carroll 2011; Jozwik and Carroll 2012). In breast cancer, FoxA1 expression is a significant predictor of cancer-specific survival in patients (Badve et al. 2007). Thus, pioneer transcription factors would be potential biomarkers and drug targets for cancers treatment.

Pioneer factors and the future of cellular reprogramming

As seen from the above discussion, pioneer factors are distinguished by their ability to recognize target DNA sequences on nucleosomes under conditions where other factors cannot. Such binding may be stabilized by coordinate actions of other proteins and is typically succeeded by changes in the local chromatin structure, thereby enabling subsequent events required for gene regulation. The ability to target silent genes embedded in closed chromatin makes pioneer factors especially suitable for cellular reprogramming, much as they have evolved for their natural roles in cellular programming in development. Still, important questions remain (Box 2).

Despite their special nucleosome-binding ability, how is it that pioneer factors recognize only a subset of their target sites in native chromatin? As we discussed here, perhaps the subset of sites that are targeted by pioneer factors involve ensembles of positioned nucleosomes in a particular cell type that favor pioneer binding. More detailed mapping of nucleosome positions in cells, along with how they may or may not be targeted by transcription factors, should provide insight.

How do different DNA-binding domain structures enable or inhibit pioneer activity? This question seems most efficiently addressed by detailed biochemical studies, comparing how different pioneer and nonpioneer factors bind (or not) to purified, defined nucleosomes. Are there preferred motif variants for targeting a closed chromatin site in the genome versus open sites, and how does the ability of a DNA-binding domain to adapt to nucleosomal DNA relate to nucleosome-binding capacity? This could be modeled on nucleosome substrates in vitro and the information used to modify a DNA-binding domain in order to enhance its initial targeting to nucleosomal DNA, thereby enhancing cellular reprogramming. This strategy is feasible because the transcription factors that are used to elicit cellular reprogramming appear to be needed only for the initial activation of the appropriate cellular networks. After initiation, the exogenous reprogramming factors can be extinguished, while

Box 2. Important issues to be addressed

- What is the influence of predicted high nucleosome occupancy and nucleosome position on defining the subset of target sites to which pioneer factors will bind in a cell?
- What are the structural features of DNA-binding domains that allow certain factors to have pioneer activity?
- How do domains outside of the DNA-binding domain affect pioneer activity and subsequent chromatin-opening events?
- How does a better understanding of the mechanism of pioneer factor interaction with nucleosomes or chromatin inform how to generate transcription factors with enhanced cellular reprogramming activity?
- How do chromatin features (e.g., H3K9me3 domains) impede pioneer factor binding?
- How are heterochromatic domains established or broken down in order to impede or enhance pioneer factor binding?

the new networks self-sustain and complete the fate conversion in the absence of the initial reprogramming factors (Takahashi and Yamanaka 2006).

How might pioneer protein domains outside of the DNA-binding domain affect nucleosome architecture and stability? Other protein domains could interact with the core histones to stabilize nucleosome binding. Furthermore, it remains to be determined whether pioneer factors “tickle” the nucleosome in a way that destabilizes the core histones or changes nucleosome conformation in a way that modulates how other factors can bind. In addition, it will be important to perform detailed biochemical studies on polynucleosomal templates so that protein domains that affect higher-order folding of the local chromatin can be assessed (Cirillo et al. 2002). Conceivably, such chromatin-opening domains could be transferred to other factors and augment their regulatory function.

Exactly how does highly compacted heterochromatin exclude pioneer factor access? Understanding how tissue-specific heterochromatic domains are made and can be broken down will provide much insight into ways to enhance cellular reprogramming. Assuming that the terminal establishment of a new cell fate after reprogramming will involve the resetting of heterochromatic domains, it seems likely that transient rather than permanent methods for heterochromatin diminution at the onset of cellular reprogramming will be the most useful.

It is presently possible to use small molecules to target chromatin modifications that globally affect most cell networks at once. We anticipate that understanding the mechanisms by which pioneer factors target specific sites in the genome, initiate regulatory events, and maintain them through mitosis and other contexts will provide more specific means for modulating cell networks that control cell fate and function in health and disease.

Acknowledgments

We thank Eileen Hulme for help in preparing the manuscript. M.I.-D. was supported by post-doctoral fellowships from the Japan Society for the Promotion of Science and the Naito, Astellas, and Uehara Foundations. K.S.Z. was supported by National Institutes of Health grants R37GM36477 and P01GM099134.

References

- Adams CC, Workman JL. 1995. Binding of disparate transcriptional activators to nucleosomal DNA is inherently cooperative. *Mol Cell Biol* **15**: 1405–1421.
- Addis RC, Ifkovits JL, Pinto F, Kellam LD, Estes P, Rentschler S, Christoforou N, Epstein JA, Gearhart JD. 2013. Optimization of direct fibroblast reprogramming to cardiomyocytes using calcium activity as a functional measure of success. *J Mol Cell Cardiol* **60**: 97–106.
- Anderson JD, Widom J. 2001. Poly(dA–dT) promoter elements increase the equilibrium accessibility of nucleosomal DNA target sites. *Mol Cell Biol* **21**: 3830–3839.
- Badve S, Turbin D, Thorat MA, Morimiya A, Nielsen TO, Perou CM, Dunn S, Huntsman DG, Nakshatri H. 2007. FOXA1 expression in breast cancer—correlation with luminal subtype A and survival. *Clin Cancer Res* **13**: 4415–4421.
- Ballare C, Castellano G, Gaveglia L, Althammer S, Gonzalez-Vallinas J, Eyraes E, Le Dily F, Zaurin R, Soronellas D, Vicent GP, et al. 2013. Nucleosome-driven transcription factor binding and gene regulation. *Mol Cell* **49**: 67–79.
- Barozzi I, Simonatto M, Bonifacio S, Yang L, Rohs R, Ghisletti S, Natoli G. 2014. Coregulation of transcription factor binding and nucleosome occupancy through DNA features of mammalian enhancers. *Mol Cell* **54**: 844–857.
- Bass AJ, Watanabe H, Mermel CH, Yu S, Perner S, Verhaak RG, Kim SY, Wardwell L, Tamayo P, Gat-Viks I, et al. 2009. SOX2 is an amplified lineage-survival oncogene in lung and esophageal squamous cell carcinomas. *Nat Genet* **41**: 1238–1242.
- Berkes CA, Bergstrom DA, Penn BH, Seaver KJ, Knoepfler PS, Tapscott SJ. 2004. Pbx marks genes for activation by MyoD indicating a role for a homeodomain protein in establishing myogenic potential. *Mol Cell* **14**: 465–477.
- Bertrand N, Castro DS, Guillemot F. 2002. Proneural genes and the specification of neural cell types. *Nat Rev Neurosci* **3**: 517–530.
- Bort R, Signore M, Tremblay K, Barbera JP, Zaret KS. 2006. Hex homeobox gene controls the transition of the endoderm to a pseudostratified, cell emergent epithelium for liver bud development. *Dev Biol* **290**: 44–56.
- Bossard P, Zaret KS. 1998. GATA transcription factors as potentiators of gut endoderm differentiation. *Development* **125**: 4909–4917.
- Boumahdi S, Driessens G, Lapouge G, Rorive S, Nassar D, Le Mercier M, Delatte B, Caauwe A, Lenglez S, Nkusi E, et al. 2014. SOX2 controls tumour initiation and cancer stem-cell functions in squamous-cell carcinoma. *Nature* **511**: 246–250.
- Budry L, Balsalobre A, Gauthier Y, Khetchoumian K, L'Honore A, Vallette S, Brue T, Figarella-Branger D, Meij B, Drouin J. 2012. The selector gene Pax7 dictates alternate pituitary cell fates through its pioneer action on chromatin remodeling. *Genes Dev* **26**: 2299–2310.
- Buganim Y, Faddah DA, Cheng AW, Itskovich E, Markoulaki S, Ganz K, Klemm SL, van Oudenaarden A, Jaenisch R. 2012. Single-cell expression analyses during cellular reprogramming reveal an early stochastic and a late hierarchic phase. *Cell* **150**: 1209–1222.
- Buganim Y, Markoulaki S, van Wietmarschen N, Hoke H, Wu T, Ganz K, Akhtar-Zaidi B, He Y, Abraham BJ, Porubsky D, et al. 2014. The developmental potential of iPSCs is greatly influenced by reprogramming factor selection. *Cell Stem Cell* **15**: 295–309.
- Cahan P, Li H, Morris SA, Lummertz da Rocha E, Daley GQ, Collins JJ. 2014. CellNet: network biology applied to stem cell engineering. *Cell* **158**: 903–915.
- Caiazzo M, Dell'Anno MT, Dvoretzskova E, Lazarevic D, Taverna S, Leo D, Sotnikova TD, Menegon A, Roncaglia P, Colciago G, et al. 2011. Direct generation of functional dopaminergic neurons from mouse and human fibroblasts. *Nature* **476**: 224–227.
- Caravaca JM, Donahue G, Becker JS, He X, Vinson C, Zaret KS. 2013. Bookmarking by specific and nonspecific binding of FoxA1 pioneer factor to mitotic chromosomes. *Genes Dev* **27**: 251–260.
- Carey M. 1998. The enhanceosome and transcriptional synergy. *Cell* **92**: 5–8.
- Carotta S, Wu L, Nutt SL. 2010. Surprising new roles for PU.1 in the adaptive immune response. *Immunol Rev* **238**: 63–75.

- Carroll JS, Liu XS, Brodsky AS, Li W, Meyer CA, Szary AJ, Eeckhoutte J, Shao W, Hestermann EV, Geistlinger TR, et al. 2005. Chromosome-wide mapping of estrogen receptor binding reveals long-range regulation requiring the forkhead protein FoxA1. *Cell* **122**: 33–43.
- Chavez S, Beato M. 1997. Nucleosome-mediated synergism between transcription factors on the mouse mammary tumor virus promoter. *Proc Natl Acad Sci* **94**: 2885–2890.
- Chen J, Zhang Z, Li L, Chen BC, Revyakin A, Hajj B, Legant W, Dahan M, Lionnet T, Betzig E, et al. 2014. Single-molecule dynamics of enhanceosome assembly in embryonic stem cells. *Cell* **156**: 1274–1285.
- Cirillo LA, Zaret KS. 1999. An early developmental transcription factor complex that is more stable on nucleosome core particles than on free DNA. *Mol Cell* **4**: 961–969.
- Cirillo LA, McPherson CE, Bossard P, Stevens K, Cherian S, Shim E-Y, Clark EA, Burley SK, Zaret KS. 1998. Binding of the winged-helix transcription factor HNF3 to a linker histone site on the nucleosome. *EMBO J* **17**: 244–254.
- Cirillo L, Lin FR, Cuesta I, Jarnik M, Friedman D, Zaret K. 2002. Opening of compacted chromatin by early developmental transcription factors HNF3 (FOXA) and GATA-4. *Mol Cell* **9**: 279–289.
- Clark KL, Halay ED, Lai E, Burley SK. 1993. Co-crystal structure of the HNF3/fork head DNA recognition motif resembles histone H5. *Nature* **364**: 412–420.
- Clarke CL, Graham JD. 2012. Non-overlapping progesterone receptor cisomes contribute to cell-specific transcriptional outcomes. *PLoS ONE* **7**: e35859.
- Cruikshanks HA, McBryan T, Nelson DM, Vanderkraats ND, Shah PP, van Tuyn J, Singh Rai T, Brock C, Donahue G, Dunican DS, et al. 2013. Senescent cells harbour features of the cancer epigenome. *Nat Cell Biol* **15**: 1495–1506.
- Davis RL, Weintraub H, Lassar AB. 1987. Expression of a single transfected cDNA converts fibroblasts to myoblasts. *Cell* **51**: 987–1000.
- Drouin J. 2014. Minireview: pioneer transcription factors in cell fate specification. *Mol Endocrinol* **28**: 989–998.
- Egli D, Birkhoff G, Eggan K. 2008. Mediators of reprogramming: transcription factors and transitions through mitosis. *Nat Rev Mol Cell Biol* **9**: 505–516.
- Espinosa JM, Emerson BM. 2001. Transcriptional regulation by p53 through intrinsic DNA/chromatin binding and site-directed cofactor recruitment. *Mol Cell* **8**: 57–69.
- Fakhouri THL, Stevenson J, Chisholm AD, Mango SE. 2010. Dynamic chromatin organization during foregut development mediated by the organ selector gene PHA-4/FoxA. *PLoS Genet* **6**: e1001060.
- Feng R, Desbordes SC, Xie H, Tillo ES, Pixley F, Stanley ER, Graf T. 2008. PU.1 and C/EBP α / β convert fibroblasts into macrophage-like cells. *Proc Natl Acad Sci* **105**: 6057–6062.
- Feng B, Ng JH, Heng JC, Ng HH. 2009. Molecules that promote or enhance reprogramming of somatic cells to induced pluripotent stem cells. *Cell Stem Cell* **4**: 301–312.
- Foo SM, Sun Y, Lim B, Ziukaite R, O'Brien K, Nien CY, Kirov N, Shvartsman SY, Rushlow CA. 2014. Zelda potentiates morphogen activity by increasing chromatin accessibility. *Curr Biol* **24**: 1341–1346.
- Foygel K, Choi B, Jun S, Leong DE, Lee A, Wong CC, Zuo E, Eckart M, Reijo Pera RA, Wong WH, et al. 2008. A novel and critical role for Oct4 as a regulator of the maternal-embryonic transition. *PLoS ONE* **3**: e4109.
- Gaffney DJ, McVicker G, Pai AA, Fondufe-Mittendorf YN, Lewellen N, Michelini K, Widom J, Gilad Y, Pritchard JK. 2012. Controls of nucleosome positioning in the human genome. *PLoS Genet* **8**: e1003036.
- Gao N, Zhang J, Rao MA, Case TC, Mirosevich J, Wang Y, Jin R, Gupta A, Rennie PS, Matusik RJ. 2003. The role of hepatocyte nuclear factor-3 α (Forkhead box A1) and androgen receptor in transcriptional regulation of prostatic genes. *Mol Endocrinol* **17**: 1484–1507.
- Gao Y, Chen J, Li K, Wu T, Huang B, Liu W, Kou X, Zhang Y, Huang H, Jiang Y, et al. 2013. Replacement of Oct4 by Tet1 during iPSC induction reveals an important role of DNA methylation and hydroxymethylation in reprogramming. *Cell Stem Cell* **12**: 453–469.
- Ghisletti S, Barozzi I, Mietton F, Polletti S, De Santa F, Venturini E, Gregory L, Lonie L, Chew A, Wei CL, et al. 2010. Identification and characterization of enhancers controlling the inflammatory gene expression program in macrophages. *Immunity* **32**: 317–328.
- Gualdi R, Bossard P, Zheng M, Hamada Y, Coleman JR, Zaret KS. 1996. Hepatic specification of the gut endoderm in vitro: cell signaling and transcriptional control. *Genes Dev* **10**: 1670–1682.
- Gurdon JB. 1962. The developmental capacity of nuclei taken from intestinal epithelium cells of feeding tadpoles. *J Embryol Exp Morphol* **10**: 622–640.
- Han J, Yuan P, Yang H, Zhang J, Soh BS, Li P, Lim SL, Cao S, Tay J, Orlov YL, et al. 2010. Tbx3 improves the germ-line competency of induced pluripotent stem cells. *Nature* **463**: 1096–1100.
- Harrison MM, Li XY, Kaplan T, Botchan MR, Eisen MB. 2011. Zelda binding in the early *Drosophila melanogaster* embryo marks regions subsequently activated at the maternal-to-zygotic transition. *PLoS Genet* **7**: e1002266.
- Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, Cheng JX, Murre C, Singh H, Glass CK. 2010. Simple combinations of lineage-determining transcription factors prime cis-regulatory elements required for macrophage and B cell identities. *Mol Cell* **38**: 576–589.
- Ho JW, Jung YL, Liu T, Alver BH, Lee S, Ikegami K, Sohn KA, Minoda A, Tolstorukov MY, Appert A, et al. 2014. Comparative analysis of metazoan chromatin organization. *Nature* **512**: 449–452.
- Hochedlinger K, Plath K. 2009. Epigenetic reprogramming and induced pluripotency. *Development* **136**: 509–523.
- Holtzinger A, Evans T. 2005. Gata4 regulates the formation of multiple organs. *Development* **132**: 4005–4014.
- Huang P, He Z, Ji S, Sun H, Xiang D, Liu C, Hu Y, Wang X, Hui L. 2011. Induction of functional hepatocyte-like cells from mouse fibroblasts by defined factors. *Nature* **475**: 386–389.
- Ieda M, Fu JD, Delgado-Olguin P, Vedantham V, Hayashi Y, Bruneau BG, Srivastava D. 2010. Direct reprogramming of fibroblasts into functional cardiomyocytes by defined factors. *Cell* **142**: 375–386.
- Ioshikhes IP, Albert I, Zanton SJ, Pugh BF. 2006. Nucleosome positions predicted through comparative genomics. *Nat Genet* **38**: 1210–1215.
- Jozwik KM, Carroll JS. 2012. Pioneer factors in hormone-dependent cancers. *Nat Rev Cancer* **12**: 381–385.
- Kadaka S, Udugama MI, Pawlicki JM, Achtman JC, Jain DP, Cheng Y, Hardison RC, Blobel GA. 2012. Tissue-specific mitotic bookmarking by hematopoietic transcription factor GATA1. *Cell* **150**: 725–737.
- Kharchenko PV, Alekseyenko AA, Schwartz YB, Minoda A, Riddle NC, Ernst J, Sabo PJ, Larschan E, Gorchakov AA, Gu T, et al. 2011. Comprehensive analysis of the chromatin landscape in *Drosophila melanogaster*. *Nature* **471**: 480–485.
- Kornberg RD, Lorch Y. 1999. Twenty-five years of the nucleosome, fundamental particle of the eukaryote chromosome. *Cell* **98**: 285–294.

- Laptenko O, Beckerman R, Freulich E, Prives C. 2011. p53 binding to nucleosomes within the p21 promoter in vivo leads to nucleosome loss and transcriptional activation. *Proc Natl Acad Sci* **108**: 10385–10390.
- Lee CS, Friedman JR, Fulmer JT, Kaestner KH. 2005. The initiation of liver development is dependent on Foxa transcription factors. *Nature* **435**: 944–947.
- Lee MT, Bonneau AR, Takacs CM, Bazzini AA, DiVito KR, Fleming ES, Giraldez AJ. 2013. Nanog, Pou5f1 and Sox1 activate zygotic gene expression during the maternal-to-zygotic transition. *Nature* **503**: 360–364.
- Leichsenring M, Maes J, Mossner R, Driever W, Onichtchouk D. 2013. Pou5f1 transcription factor controls zygotic gene activation in vertebrates. *Science* **341**: 1005–1009.
- Levine M, Cattoglio C, Tjian R. 2014. Looping back to leap forward: transcription enters a new era. *Cell* **157**: 13–25.
- Li Z, White P, Tuteja G, Rubins N, Sackett S, Kaestner KH. 2009. Foxa1 and Foxa2 regulate bile duct development in mice. *J Clin Invest* **119**: 1537–1545.
- Li Z, Gadue P, Chen K, Jiao Y, Tuteja G, Schug J, Li W, Kaestner KH. 2012a. Foxa2 and H2A.Z mediate nucleosome depletion during embryonic stem cell differentiation. *Cell* **151**: 1608–1616.
- Li Z, Tuteja G, Schug J, Kaestner KH. 2012b. Foxa1 and Foxa2 are essential for sexual dimorphism in liver cancer. *Cell* **148**: 72–83.
- Liang HL, Nien CY, Liu HY, Metzstein MM, Kirov N, Rushlow C. 2008. The zinc-finger protein Zelda is a key activator of the early zygotic genome in *Drosophila*. *Nature* **456**: 400–403.
- Lidor Nili E, Field Y, Lubling Y, Widom J, Oren M, Segal E. 2010. p53 binds preferentially to genomic regions with high DNA-encoded nucleosome occupancy. *Genome Res* **20**: 1361–1368.
- Lupien M, Eeckhoutte J, Meyer CA, Wang Q, Zhang Y, Li W, Carroll JS, Liu XS, Brown M. 2008. FoxA1 translates epigenetic signatures into enhancer-driven lineage-specific transcription. *Cell* **132**: 958–970.
- Mahony S, Edwards MD, Mazzoni EO, Sherwood RJ, Kakumanu A, Morrison CA, Wichterle H, Gifford DK. 2014. An integrated model of multiple-condition ChIP-seq data reveals predeterminants of Cdx2 binding. *PLoS Comput Biol* **10**: e1003501.
- Martinez-Balbas MA, Dey A, Rabindran SK, Ozato K, Wu C. 1995. Displacement of sequence-specific transcription factors from mitotic chromatin. *Cell* **83**: 29–38.
- McPherson CE, Shim E-Y, Friedman DS, Zaret KS. 1993. An active tissue-specific enhancer and bound transcription factors existing in a precisely positioned nucleosomal array. *Cell* **75**: 387–398.
- McPherson CE, Horowitz R, Woodcock CL, Jiang C, Zaret KS. 1996. Nucleosome positioning properties of the albumin transcriptional enhancer. *Nucleic Acids Res* **24**: 397–404.
- Menet JS, Pescatore S, Rosbash M. 2014. CLOCK:BMAL1 is a pioneer-like transcription factor. *Genes Dev* **28**: 8–13.
- Meshorer E, Yellajoshula D, George E, Scambler PJ, Brown DT, Misteli T. 2006. Hyperdynamic plasticity of chromatin proteins in pluripotent embryonic stem cells. *Dev Cell* **10**: 105–116.
- Michelotti EF, Sanford S, Levens D. 1997. Marking of active genes on mitotic chromosomes. *Nature* **388**: 895–899.
- Miller JA, Widom J. 2003. Collaborative competition mechanism for gene activation in vivo. *Mol Cell Biol* **23**: 1623–1632.
- Mirny LA. 2010. Nucleosome-mediated cooperativity between transcription factors. *Proc Natl Acad Sci* **107**: 22534–22539.
- Morris SA, Cahan P, Li H, Zhao AM, San Roman AK, Shivdasani RA, Collins JJ, Daley GQ. 2014. Dissecting engineered cell types and enhancing cell fate conversion via CellNet. *Cell* **158**: 889–902.
- Moyle-Heyman G, Tims HS, Widom J. 2011. Structural constraints in collaborative competition of transcription factors against the nucleosome. *J Mol Biol* **412**: 634–646.
- Naumova N, Imakaev M, Fudenberg G, Zhan Y, Lajoie BR, Mirny LA, Dekker J. 2013. Organization of the mitotic chromosome. *Science* **342**: 948–953.
- Nien CY, Liang HL, Butcher S, Sun Y, Fu S, Gocha T, Kirov N, Manak JR, Rushlow C. 2011. Temporal coordination of gene networks by Zelda in the early *Drosophila* embryo. *PLoS Genet* **7**: e1002339.
- Pan H, Schultz RM. 2011. Sox2 modulates reprogramming of gene expression in two-cell mouse embryos. *Biol Reprod* **85**: 409–416.
- Pihlajamaa P, Sahu B, Lyly L, Aittomäki V, Hautaniemi S, Janne OA. 2014. Tissue-specific pioneer factors associate with androgen receptor cisomes and transcription programs. *EMBO J* **33**: 312–326.
- Ptashne M. 2011. Principles of a switch. *Nat Chem Biol* **7**: 484–487.
- Ramakrishnan V, Finch JT, Graziano V, Lee PL, Sweet RM. 1993. Crystal structure of globular domain of histone H5 and its implications for nucleosome binding. *Nature* **362**: 219–224.
- Schäfer BW, Blakely BT, Darlington GJ, Blau HM. 1990. Effect of cell history on response to helix-loop-helix family of myogenic regulators. *Nature* **344**: 454–459.
- Schones DE, Cui K, Cuddapah S, Roh TY, Barski A, Wang Z, Wei G, Zhao K. 2008. Dynamic regulation of nucleosome positioning in the human genome. *Cell* **132**: 887–898.
- Segal E, Fondufe-Mittendorf Y, Chen L, Thastrom A, Field Y, Moore IK, Wang JP, Widom J. 2006. A genomic code for nucleosome positioning. *Nature* **442**: 772–778.
- Sekiya S, Suzuki A. 2011. Direct conversion of mouse fibroblasts to hepatocyte-like cells by defined factors. *Nature* **475**: 390–393.
- Sekiya T, Zaret KS. 2007. Repression by Groucho/TLE/Grg proteins: genomic site recruitment generates compacted chromatin in vitro and impairs activator binding in vivo. *Mol Cell* **28**: 291–303.
- Serandour AA, Avner S, Percevault F, Demay F, Bizot M, Lucchetti-Miganeh C, Barloy-Hubler F, Brown M, Lupien M, Metivier R, et al. 2011. Epigenetic switch involved in activation of pioneer factor FOXA1-dependent enhancers. *Genome Res* **21**: 555–565.
- Sherwood RJ, Hashimoto T, O'Donnell CW, Lewis S, Barkal AA, van Hoff JP, Karun V, Jaakkola T, Gifford DK. 2014. Discovery of directional and nondirectional pioneer transcription factors by modeling DNase profile magnitude and shape. *Nat Biotechnol* **32**: 171–178.
- Shim EY, Woodcock C, Zaret KS. 1998. Nucleosome positioning by the winged-helix transcription factor HNF3. *Genes Dev* **12**: 5–10.
- Son EY, Ichida JK, Wainger BJ, Toma JS, Rafuse VF, Woolf CJ, Eggan K. 2011. Conversion of mouse and human fibroblasts into functional spinal motor neurons. *Cell Stem Cell* **9**: 205–218.
- Soufi A, Donahue G, Zaret KS. 2012. Facilitators and impediments of the pluripotency reprogramming factors' initial engagement with the genome. *Cell* **151**: 994–1004.
- Struhl K. 1999. Fundamentally different logic of gene regulation in eukaryotes and prokaryotes. *Cell* **98**: 1–4.
- Struhl K, Segal E. 2013. Determinants of nucleosome positioning. *Nat Struct Mol Biol* **20**: 267–273.

- Tadros W, Lipshitz HD. 2009. The maternal-to-zygotic transition: a play in two acts. *Development* **136**: 3033–3042.
- Takahashi K, Yamanaka S. 2006. Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors. *Cell* **126**: 663–676.
- Taube JH, Allton K, Duncan SA, Shen L, Barton MC. 2010. Foxal functions as a pioneer transcription factor at transposable elements to activate *Afp* during differentiation of embryonic stem cells. *J Biol Chem* **285**: 16135–16144.
- Teif VB, Vainshtein Y, Caudron-Herger M, Mallm JP, Marth C, Hofer T, Rippe K. 2012. Genome-wide nucleosome positioning during embryonic stem cell development. *Nat Struct Mol Biol* **19**: 1185–1192.
- Thurman RE, Rynes E, Humbert R, Vierstra J, Maurano MT, Haugen E, Sheffield NC, Stergachis AB, Wang H, Vernot B, et al. 2012. The accessible chromatin landscape of the human genome. *Nature* **489**: 75–82.
- Tillo D, Hughes TR. 2009. G+C content dominates intrinsic nucleosome occupancy. *BMC Bioinformatics* **10**: 442.
- Tillo D, Kaplan N, Moore IK, Fondufe-Mittendorf Y, Gossett AJ, Field Y, Lieb JD, Widom J, Segal E, Hughes TR. 2010. High nucleosome occupancy is encoded at human regulatory sequences. *PLoS ONE* **5**: e9129.
- Updike DL, Mango SE. 2006. Temporal regulation of foregut development by HTZ-1/H2A.Z and PHA-4/FoxA. *PLoS Genet* **2**: e161.
- Valouev A, Johnson SM, Boyd SD, Smith CL, Fire AZ, Sidow A. 2011. Determinants of nucleosome organization in primary human cells. *Nature* **474**: 516–520.
- Vashee S, Melcher K, Ding WV, Johnston SA, Kodadek T. 1998. Evidence for two modes of cooperative DNA binding in vivo that do not involve direct protein–protein interactions. *Curr Biol* **8**: 452–458.
- Vierbuchen T, Ostermeier A, Pang ZP, Kokubu Y, Sudhof TC, Wernig M. 2010. Direct conversion of fibroblasts to functional neurons by defined factors. *Nature* **463**: 1035–1041.
- Wang X, Bai L, Bryant GO, Ptashne M. 2011. Nucleosomes and the accessibility problem. *Trends Genet* **27**: 487–492.
- Wang S, Linde MH, Munde M, Carvalho VD, Wilson WD, Poon GM. 2014. Mechanistic heterogeneity in site recognition by the structurally homologous DNA-binding domains of the ETS family transcription factors Ets-1 and PU.1. *J Biol Chem* **289**: 21605–21616.
- Wapinski OL, Vierbuchen T, Qu K, Lee QY, Chanda S, Fuentes DR, Giresi PG, Ng YH, Marro S, Neff NF, et al. 2013. Hierarchical mechanisms for direct reprogramming of fibroblasts to neurons. *Cell* **155**: 621–635.
- Watt AJ, Zhao R, Li J, Duncan SA. 2007. Development of the mammalian liver and ventral pancreas is dependent on GATA4. *BMC Dev Biol* **7**: 37.
- Weintraub H, Tapscott SJ, Davis RL, Thayer MJ, Adam MA, Lassar AB, Miller AD. 1989. Activation of muscle-specific genes in pigment, nerve, fat, liver, and fibroblast cell lines by forced expression of MyoD. *Proc Natl Acad Sci* **86**: 5434–5438.
- Xu C, Lu X, Chen EZ, He Z, Uyunbilig B, Li G, Ma Y, Hui L, Xie B, Gao Y, et al. 2012. Genome-wide roles of Foxa2 in directing liver specification. *J Mol Cell Biol* **4**: 420–422.
- Xu Z, Chen H, Ling J, Yu D, Struffi P, Small S. 2014. Impacts of the ubiquitous factor Zelda on Bicoid-dependent DNA binding and transcription in *Drosophila*. *Genes Dev* **28**: 608–621.
- Yamamoto KR. 1985. Steroid receptor regulated transcription of specific genes and gene networks. *Annu Rev Genet* **19**: 209–252.
- Yu J, Vodyanik MA, Smuga-Otto K, Antosiewicz-Bourget J, Frane JL, Tian S, Nie J, Jonsdottir GA, Ruotti V, Stewart R, et al. 2007. Induced pluripotent stem cell lines derived from human somatic cells. *Science* **318**: 1917–1920.
- Zaidi SK, Young DW, Montecino MA, Lian JB, van Wijnen AJ, Stein JL, Stein GS. 2010. Mitotic bookmarking of genes: a novel dimension to epigenetic control. *Nat Rev Genet* **11**: 583–589.
- Zaret KS. 2014. Genome reactivation after the silence in mitosis: recapitulating mechanisms of development? *Dev Cell* **29**: 132–134.
- Zaret KS, Carroll JS. 2011. Pioneer transcription factors: establishing competence for gene expression. *Genes Dev* **25**: 2227–2241.
- Zhang L, Rubins NE, Ahima RS, Greenbaum LE, Kaestner KH. 2005. Foxa2 integrates the transcriptional response of the hepatocyte to fasting. *Cell Metab* **2**: 141–148.
- Zhao R, Watt AJ, Li J, Luebke-Wheeler J, Morrissey EE, Duncan SA. 2005. GATA6 is essential for embryonic development of the liver but dispensable for early heart formation. *Mol Cell Biol* **25**: 2622–2631.

Pioneer transcription factors in cell reprogramming

Makiko Iwafuchi-Doi and Kenneth S. Zaret

Genes Dev. 2014, **28**:

Access the most recent version at doi:[10.1101/gad.253443.114](https://doi.org/10.1101/gad.253443.114)

References

This article cites 122 articles, 31 of which can be accessed free at:
<http://genesdev.cshlp.org/content/28/24/2679.full.html#ref-list-1>

Creative Commons License

This article, published in *Genes & Development*, is available under a Creative Commons License (Attribution 4.0 International), as described at <http://creativecommons.org/licenses/by/4.0>.

Email Alerting Service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).