A monoallelic-to-biallelic T-cell transcriptional switch regulates GATA3 abundance

Chia-Jui Ku,¹ Kim-Chew Lim,¹ Sundeep Kalantry,² Ivan Maillard,^{1,3,4} James Douglas Engel,¹ and Tomonori Hosoya¹

¹Department of Cell and Developmental Biology, University of Michigan Medical School, Ann Arbor, Michigan 48109, USA;

²Department of Human Genetics, University of Michigan Medical School, Ann Arbor, Michigan 48109, USA; ³Division of Hematology-Oncology, Department of Internal Medicine, University of Michigan Medical School, Ann Arbor, Michigan 48109, USA; ⁴Life Sciences Institute, University of Michigan Medical School, Ann Arbor, Michigan 48109, USA

Protein abundance must be precisely regulated throughout life, and nowhere is the stringency of this requirement more evident than during T-cell development: A twofold increase in the abundance of transcription factor GATA3 results in thymic lymphoma, while reduced GATA3 leads to diminished T-cell production. GATA3 haploinsufficiency also causes human HDR (hypoparathyroidism, deafness, and renal dysplasia) syndrome, often accompanied by immunodeficiency. Here we show that loss of one *Gata3* allele leads to diminished expansion (and compromised development) of immature T cells as well as aberrant induction of myeloid transcription factor PU.1. This effect is at least in part mediated transcriptionally: We discovered that *Gata3* is monoallelically expressed in a parent of origin-independent manner in hematopoietic stem cells and early T-cell progenitors. Curiously, half of the developing cells switch to biallelic *Gata3* transcription abruptly at midthymopoiesis. We show that the monoallelic-to-biallelic transcriptional switch is stably maintained and therefore is not a stochastic phenomenon. This unique mechanism, if adopted by other regulatory genes, may provide new biological insights into the rather prevalent phenomenon of monoallelic expression of autosomal genes as well as into the variably penetrant pathophysiological spectrum of phenotypes observed in many human syndromes that are due to haploinsufficiency of the affected gene.

[**Keywords:** GATA3; T-cell development; monoallelic transcription; biallelic transcription]

Supplemental material is available for this article.

Received May 4, 2015; revised version accepted August 21, 2015.

Precise control of protein abundance—achieved through balanced transcriptional activation and repression, mRNA generation and stability, translation and protein stability, and localization—is vital for normal development. In metazoan organisms, most genes are encoded by paternal and maternal alleles, and cells are thought to most often express autosomal genes from both alleles. Trisomy can result in miscarriage or physiological abnormalities (e.g., Down syndrome, Edwards syndrome, Patau syndrome, Warkany syndrome 2, and more) (Hassold and Hunt 2001), while haploinsufficiency has been shown to result in numerous autosomal dominant disorders (e.g., Williams syndrome, autosomal dominant Zinsser-Cole-Engman syndrome, Ehlers-Danlos syndrome, Marfan syndrome, and others) (Inoue and Lupski 2002). Thus, the number of active alleles can be a principal determinant used to modulate the level of protein expression within a cell.

Transcription factor GATA3 (originally named NF-E1c) (Yamamoto et al. 1990; Ko et al. 1991) is required throughout T-cell development (Hosoya et al. 2010). T lymphocytes develop in the thymus from multipotential hematopoietic progenitors that reside in the bone marrow. Bone marrow-derived progenitors then home to the thymus, where they undergo cell expansion to become early T-cell progenitors (ETPs) that are specified to the T-cell lineage (Allman et al. 2003; Porritt et al. 2004; Sambandam et al. 2005; Tan et al. 2005; Bell and Bhandoola 2008). ETPs sequentially develop through double-negative (DN) stages (DN2–DN4)—during which cells do not express either the CD4 or CD8 coreceptor (CD4[−] CD8[−])—into a double-positive stage (DP; CD4⁺ CD8⁺) and finally resolve into either CD4 single-positive (SP) helper T cells

Corresponding author: engel@umich.edu

Article is online at <http://www.genesdev.org/cgi/doi/10.1101/gad.265025.115>.

© 2015 Ku et al. This article is distributed exclusively by Cold Spring Harbor Laboratory Press for the first six months after the full-issue publication date (see <http://genesdev.cshlp.org/site/misc/terms.xhtml>). After six months, it is available under a Creative Commons License (Attribution-NonCommercial 4.0 International), as described at <http://creativecommons.org/licenses/by-nc/4.0/>.

or CD8 SP cytotoxic T cells. These mature SP cells then exit the thymus via the circulatory system to seed secondary lymphoid organs, where they acquire immune competence (Fowlkes and Mathieson 1985; Penit and Vasseur 1988).

Patients with *GATA3* haploinsufficiency suffer from HDR (hypoparathyroidism, deafness, and renal dysplasia) syndrome and present variable clinical symptoms, including hypoparathyroidism, heart defects, deafness, renal malformation, and reduced T-cell numbers (Daw et al. 1996; van Esch et al. 2000). These developmental anomalies are also reproduced when *Gata3* is disrupted in mice (Lim et al. 2000; Grigorieva et al. 2010), while a twofold increase in the abundance of GATA3 protein results in thymic lymphoma (Nawijn et al. 2001). Thus, GATA3 abundance must be precisely controlled during T-lymphocyte development as well as in multiple organs. While it has been reported that such stringent control over GATA3 abundance exists (e.g., see Scripture-Adams et al. 2014), how such precision is achieved is currently speculative.

Results

T cells with a single Gata3 allele exhibit diminished GATA3 mRNA, impaired cell expansion, and elevated PU.1 expression

To examine the consequences of *Gata3* haploinsufficiency on T-cell development, we analyzed heterozygous mice bearing one wild-type (*Gata3*⁺) and one null mutant (*Gata3*^z) allele (van Doorninck et al. 1999). The absolute number of CD4 SP thymocytes in *Gata3*^{z/+} mice was 77% of *Gata3*^{+/+} (wild-type) mice, which was statistically significant, while the reduction in the number of DP (85%) and CD8 SP (92%) thymocytes was modest and statistically insignificant (Supplemental Fig. S1A). A similar reduction in the number of DP (79%), CD4 SP (56%), and CD8 SP (72%) thymocytes was also observed in *Gata3*^{flox/+}:Tg^{Cd2Cre} mice (Supplemental Fig. S1B) in which one *Gata3* allele was removed in only B and T lymphocytes, and thus the effect is lymphoid cell-autonomous. Since Tg^{Cd2Cre} alone has no effect on T-cell development (Shi and Petrie 2012), compromised T-cell development in the thymi of *Gata3*^{flox/+}:Tg^{Cd2Cre} mice could be directly attributed to the lymphocyte-specific inactivation of one *Gata3* allele. Both *Gata3*^{z/+} (Supplemental Fig. S2A) and *Gata3*^{g/+} (Supplemental Fig. S2B) heterozygotes had a reduced number (~50%) of ETP in comparison with wild-type mice, while the number of ETP in *Gata3*^{flox/+}:Tg^{Cd2Cre} heterozygous mice was not statistically different from the number in controls (Supplemental Fig. S2C). The difference in ETP number between these different mutant alleles suggests that the 50% reduced number of ETP (in *Gata3*^{z/+} [Supplemental Fig. S2A] or *Gata3*^{g/+} [Supplemental Fig. S2B] heterozygous mutant mice) is because there is only one active *Gata3* allele in either prethymic progenitors or other non-lymphoid cells. However, this reduced number of ETP was compensated, possibly by excess proliferation in

these immature cells, such that no significant reduction in thymocyte number was observed in heterozygous mutant mice through the DN2 to DN4 stages (Supplemental Fig. S2).

To confirm the intrinsic developmental potential of *Gata3* heterozygous mutant cells, we isolated DN4 stage T cells from *Gata3*^{z/+} thymi and cocultured them on OP9-DL1 feeder cells. OP9-DL1 feeder cells promote Notch signaling, thereby supporting further T-cell development (Schmitt and Zuniga-Pflucker 2002). This method has some advantages as an assay for cell yield and viability, as phagocytes may conceal differences by clearing dead cells in the thymus in vivo. Equal numbers of flow-sorted *Gata3*^{z/+} or *Gata3*^{+/+} DN4 stage cells were cocultured on OP9-DL1 in the presence of Flt3 ligand and IL7. T cells were recovered, stained, and analyzed by flow cytometry for four consecutive days following the initiation of the cocultures. The significant reduction in DN and DP T cells in cocultures seeded with *Gata3*^{z/+} DN4 cells was most evident after 3 or 4 d of coculture (Fig. 1A), when compared with cocultures seeded with wild-type DN4 stage thymocytes. The frequency of apoptotic and dead (Annexin V⁺) cells on each day in *Gata3*^{z/+} DN and DP cell populations was higher in comparison with *Gata3*^{+/+} cells (Fig. 1B). Hence, a reduction in *Gata3* gene dosage attenuates DN4 cell proliferation, viability, and differentiation.

To evaluate whether the different outcomes of *Gata3*^{z/+} and *Gata3*^{+/+} DN4 OP9-DL1 cocultures could be attributed to altered kinetics in GATA3 mRNA accumulation during T-lineage progression, cells at different T-cell developmental stages (ETP to SP) were isolated from *Gata3*^{z/+} and *Gata3*^{+/+} thymi and analyzed by quantitative RT-PCR (qRT-PCR). We found that the levels of GATA3 mRNA in *Gata3*^{+/+} and *Gata3*^{z/+} were virtually identical at the ETP and DN2 stages, while its abundance in DN3a and later stage thymocytes was reduced by between 15% and 30% (Fig. 1C). We next tested whether the differences in mRNA levels coincide with protein accumulation at different stages. Thymocytes were isolated from *Gata3*^{z/+} and *Gata3*^{+/+} thymi, and GATA3 protein abundance was quantified by flow cytometry as described (Scripture-Adams et al. 2014). GATA3 protein abundance in *Gata3*^{z/+} was virtually identical to wild type at the ETP and DN2 stages, while its abundance in DN3a and later stage thymocytes was reduced (Fig. 1D), in keeping with the reduction in GATA3 mRNA (Fig. 1C). These data indicate that, during late thymopoiesis, only cells bearing two wild-type *Gata3* alleles can elevate GATA3 to wild-type levels and that the failure to attain this temporally critical GATA3 surge (in *Gata3*^{z/+} DN4 cells) perturbs their development in vitro and in vivo. The data suggested that GATA3 protein abundance is at least in part regulated at the transcriptional level and that GATA3 abundance critically controls T-cell expansion and the development of thymocytes. This conclusion is entirely consistent with the modest deficiency in T-cell number observed in human GATA3 haploinsufficient patients (Daw et al. 1996; Lichtner et al. 2000).

Figure 1. Reduced activity of *Gata3* alleles results in reduced expansion of immature T cells and elevated expression of myeloid transcription factor PU.1. (A,B) DN4 stage thymocytes were isolated from *Gata3*^{+/+} (○) or *Gata3*^{z/+} (●) mice and then cocultured on OP9-DL1 feeder cells. Cells were harvested and stained for CD4 and CD8 coreceptors followed by flow cytometric analysis at the indicated time points (after 1–4 d of coculture initiation) for cell number (A) and apoptosis (by Annexin V staining) (B). The horizontal bar in each genotype panel represents the average value for each analyzed genotype group. Representative data are shown from one experiment examining three mice of each genotype; similar results were collected from at least six mice of each genotype in two independent experiments. (C,E) Quantitative RT-PCR (qRT-PCR) in staged T cells of wild-type (*Gata3*^{+/+}; ●) or heterozygous mutant (*Gata3*^{z/+}; ○) mice. GATA3 (C) and PU.1 (E) mRNA abundance was quantified using the total reverse-transcribed product from 100 live-cell equivalents of RNA normalized to HPRT mRNA. Each circle represents an individual mouse, and the black bars represent the averages. (*) $P < 0.05$; (NS) not significant. The data summarize duplicate measurements of three to eight mice of each genotype from at least two independent experiments. (D) Intracellular GATA3 (filled curve) protein abundance in staged T cells from wild-type (*Gata3*^{+/+}) (top panel) and heterozygous mutant (*Gata3*^{z/+}) (bottom panel) thymi was monitored by flow cytometry. The open curve represents background (IgG) staining in each sample. Representative histograms are shown as characterized in at least three mice of each genotype.

During the transition from the DN2 to the DN3 stage, when a quantitative difference in GATA3 mRNAs and protein between heterozygous and wild-type mice (Fig. 1C,D) was first detected, GATA3 was shown to block apoptosis and repress transcription factor PU.1 (Scripture-Adams et al. 2014), a key protein that promotes myeloid gene expression (Nerlov and Graf 1998). Mice in which the *Gata3* gene was conditionally ablated at the DN3 stage exhibited increased apoptosis and reduced TCR β protein, but not mRNA, expression (Pai et al. 2003). To address the possibility that the differential *Gata3* abundance observed between *Gata3*^{z/+} and *Gata3*^{+/+} animals might account for these observations, *Gata3*^{z/+} heterozygotes were further analyzed.

We first assessed the intracellular abundance of TCR β by flow cytometry and found that, contrary to the study by Pai et al. (2003), there was only very modest variation in intracellular TCR β abundance in *Gata3*^{+/+} compared with *Gata3*^{z/+} DN3 stage thymocytes (Supplemental Fig. S3). In contrast, PU.1 mRNA abundance increased by approximately twofold in *Gata3*^{z/+} T cells at both the DN3a and DN3b stages when compared with the equivalent wild-type *Gata3* stages (Fig. 1E). Hence, these data show that only T cells with two intact *Gata3* alleles can promote normal expansion and development as well as repress PU.1 in DN3a and DN3b thymocytes but also that one active *Gata3* allele is sufficient for normal levels of TCR β protein accumulation.

Gata3 is monoallelically expressed in early DN thymocytes but is biallelically expressed in a subpopulation of late stage thymocytes

The previous data demonstrate that the molecular basis for regulating GATA3 abundance at different developmental stages is at least in part dictated by transcription. To shed light on possible mechanisms by which this reduction in abundance of GATA3 between the DN2 and DN3a stages (Fig. 1C,D) might be elicited, we initially hypothesized that this could be most directly explained by either enhanced transcription from both chromosomes or, although unprecedented in developmental mechanisms, switching from monoallelic to biallelic *Gata3* transcription.

Three mechanisms have been shown to lead to the epigenetic activation of genes on only one of two equivalent chromosomes. X-chromosome inactivation, first described more than half a century ago, generates one randomly inactivated X chromosome in females so that both mammalian sexes express equivalent amounts of X-encoded gene product (Ohno et al. 1959; Lyon 1961). In contrast to the inactivation of a whole chromosome, mammalian genomic imprinting selectively inactivates specific autosomal loci in a male or female parent of origin-dependent manner (Bartolomei et al. 1991; DeChiara et al. 1991). Finally, the discovery that multiple autosomal genes, including *Ilf2* (Hollander et al. 1998; Rhoades et al. 2000) and *Tlr4* (Pereira et al. 2003), are monoallelically expressed provided yet a third class of unequally expressed genes. Even more recently, this category was extended to include widely dispersed genes comprising 8%–24% of all autosomal transcripts that are monoallelically regulated (Gimelbrant et al. 2007; Deng et al. 2014). From these studies, there appeared to emerge two distinct types of autosomal monoallelic expression: (1) a subset that is expressed in an unstable, stochastic pattern (Deng et al. 2014) and (2) a subset in which the pattern is stably maintained (Gimelbrant et al. 2007; Zwemer et al. 2012). The latter observations imply that the number of active alleles is precisely regulated in order to achieve alternative levels of gene product encoded by some unknown number of key genes, although this commonly held maxim has not been directly challenged.

To address whether *Gata3* is monoallelically or biallelically expressed in ETP and DN2 stage thymocytes (Fig. 1C,D), we exploited a silent single nucleotide polymorphism (SNP) in the third *Gata3* exon that differs between inbred mouse strains C57Bl/6J (B6) and FVB/NJ (FVB) F1 progeny using single-cell RT–PCR followed by Sanger sequencing. To distinguish between the paternal and maternal alleles, F1 (B6 × FVB) hybrid mice were generated by crossing B6 males to FVB females in order to determine whether one or both alleles in individual thymocytes were expressed and whether that expression was dependent on parental gender (genomic imprinting). Hematopoietic stem cells (HSCs) in the bone marrow and ETP through SP F1 thymocytes were individually isolated by flow sorting and lysed, total RNA was reverse-transcribed, and the cDNA was amplified by PCR and sequenced (Supplemental Fig. S4A). To test the sensitivity of the method, cDNAs from FVB and B6 strains were mixed in different ratios and analyzed (Supplemental Fig. S4B). The detection limit of the method was ~5%, and therefore, when >95% dominance of either transcript was detected, that was defined as monoallelic expression, while a mixture of both alleles (between 5% and 95%) was defined as biallelic.

As summarized in Table 1, individual cells recovered from adult mice at all early stages exhibited a profound bias for monoallelic *Gata3* expression: HSC (100%), ETP (97%), DN2 (67%), and DN3a (74%). In contrast, from the DN3b stage onward, approximately half of the cells expressed *Gata3* from only one allele, while the other half expressed both mRNAs (Table 1). The observed gender-independent activation of either the paternal (B6) or maternal (FVB) allele (modestly biased in favor of generation of the paternal allele) indicates that *Gata3* monoallelic expression is not sex-dependent or parent of origin-dependent, and thus the monoallelic to biallelic GATA3 expression bias that is exhibited during midthymopoiesis is likely to be regulated by a mechanism distinct from the ones used in X inactivation or genomic imprinting.

To confirm the conclusion from the single-cell mRNA analysis, we also examined transcript localization in developmentally staged T-cell nuclei by nascent transcript fluorescence in situ hybridization (RNA-FISH). Cells

Table 1. *GATA3* mRNA expression in single, staged adult bone marrow cells and thymocytes (from *GATA3* RT–PCR SNP sequence)

Developmental stage	Cells sorted	Monoallelic		Biallelic	Monoallelic
		Paternal	Maternal		
HSC	212	55	28	0	100%
ETP	158	31	27	2	97%
DN2	50	18	9	13	68%
DN3a	50	20	6	9	74%
DN3b	50	12	9	21	50%
DN4	50	11	3	20	43%
DP	122	20	6	22	44%
CD4 SP	92	14	11	26	49%
CD8 SP	92	15	10	18	58%

Data represent the summary of at least six animals analyzed at each stage.

recovered from the bone marrow or thymi of wild-type adult B6 mice were flow-sorted according to developmental stage and then hybridized to FISH probes that detect CD45 (expressed on all hematopoietic progenitor cells) as well as GATA3 (expressed at all stages of T-cell development) (David-Fung et al. 2006; Ku et al. 2012). Cells were then counterstained with 4',6-diamidion-2-phenylindole (DAPI) to define the nucleus and finally visualized by fluorescence microscopy. Only cells with visually complete, intact nuclei were imaged.

The pan-hematopoietic marker CD45 was expressed from two chromosomal alleles in all cells imaged (Fig. 2, pseudo-red spots in each panel). In all HSCs (13 of 13) and ETPs (13 of 13) as well as nearly two-thirds of the DN2 (10 of 15) and DN3a (13 of 20) stage T cells, GATA3 primary transcripts could be detected only at a single chromosomal location (Fig. 2, green spots; quantified in Table 2). However, in DN3b and later stage thymocytes, monoallelic and biallelic expression among the cells was essentially equally distributed (Fig. 2; Table 2).

Genes expressed at very low levels have been theorized to exhibit transcription of only one allele due to completely random events (Hume 2000; Kuznetsov et al. 2002), and experimentally it has been shown that *Il4* and *Il13* are transcribed from either one or two alleles in Th2 cells by probabilistic stochastic regulation (Guo et al. 2005). If the availability of transcriptional machinery is the limiting cause of monoallelic expression at the *Gata3* locus, then one should observe a higher monoallelic cell frequency in CD8 SP thymocytes, which express the lowest levels of GATA3 at all thymocyte stages (Hernández-

Hoyos et al. 2003; Ku et al. 2012). The data shown here suggest the opposite of this mode of monoallelic expression; GATA3 is expressed lower in CD8 SP cells than in ETPs or HSCs, but, nonetheless, CD8 SP cells exhibit a higher frequency of *Gata3* biallelic cells (Tables 1, 2). If the *Gata3* locus were regulated by probabilistic stochastic regulation like *Il4* or *Il13*, a higher frequency of biallelic expression should be observed at stages of development where GATA3 is most abundant. On the contrary, stage-specific and GATA3 transcript abundance-independent monoallelic expression was observed only at the HSC and ETP stages in the *Gata3* locus. These data demonstrate that *Gata3* monoallelic expression and the switch to biallelic expression differ from a monoallelic expression pattern that would be generated in a stochastic manner (Eckersley-Maslin and Spector 2014). Thus, we conclude that *Gata3* is monoallelically expressed in early stages of T-cell development and that half of the cells then switch to biallelic expression at intermediate stages of T-cell thymic development.

The commitment to Gata3 monoallelic or biallelic transcriptional status is regulated by a stable mechanism

GATA3 activity is required for initial T-cell generation as well as several intermediate maturational stages (Hosoya et al. 2010; Yui and Rothenberg 2014). In considering experiments that might shed further light on these phenomena, we reasoned that a hypomorphic *Gata3-eGFP* fusion allele (*Gata3⁸*) (Hosoya et al. 2009) might not generate adequate GATA3 functional activity to allow

Figure 2. *Gata3* is transcribed from only one allele in HSCs and ETPs and then switches to biallelic transcription in half of DN3b and later stage T cells. Adult bone marrow cells and thymocytes were isolated using conventional cell surface markers (Hosoya et al. 2009; Ku et al. 2012), hybridized in situ to fluorescently labeled probes to detect either GATA3 (green spots) or CD45 (red spots), and analyzed by RNA-FISH (Materials and Methods). Bar, 3 μ m. Representative images are shown from at least four mice of each developmental stage from five independent experiments and are summarized in Table 2. The images shown were taken as a single focal plane, but Z-stack images were recorded to confirm that no overlapping dots were misrepresented as a single dot on the single-focal-plane images.

Table 2. *GATA3* mRNA expression in single, staged adult bone marrow cells and thymocytes (from *GATA3* RNA-FISH)

Developmental stage	Cells imaged	Monoallelic	Biallelic	Monoallelic
HSC	40	13	0	100%
ETP	40	13	0	100%
DN2	30	10	5	67%
DN3a	30	13	7	65%
DN3b	30	9	13	41%
DN4	30	8	12	40%
DP	40	11	13	46%
CD4 SP	40	10	13	44%
CD8 SP	40	7	11	39%

Data represent the summary of at least four animals analyzed at each stage. CD45 was biallelically expressed in all imaged cells.

T-cell intermediates to survive and thereby confer a selective survival or developmental advantage in adult T cells that initially express the wild-type allele at the ETP and DN2 stages in *Gata3*^{g/+} mice (Fig. 3A). Again, this hypothesis is consistent with the recent demonstration that progression through early T-cell development is exquisitely sensitive to GATA3 abundance (Scripture-Adams et al. 2014). We discovered support for this hypothesis by determining that eGFP fluorescence was absent in *Gata3*^{g/+} adult ETP and DN2 stage thymocytes (Fig. 3B), but, as cells mature and switch to biallelic transcription at the DN3 stage, eGFP-positive (eGFP⁺) thymocytes began to appear (Fig. 3B). This observation was confirmed by examining a mouse bearing a different *Gata3*-null mutant allele in which the bacterial *lacZ* gene was inserted at the *Gata3* start codon (*Gata3*^{z/+}) (van Doorninck et al. 1999); thymocytes only began to transcribe the *lacZ* allele after the DN2 stage (Supplemental Fig. S5).

To determine whether *Gata3* is transcribed from the wild-type allele in eGFP-negative (eGFP⁻) *Gata3*^{g/+} T cells, eGFP⁻ and eGFP⁺ DN4 stage cells were isolated from *Gata3*^{g/+} thymi by flow cytometry (Supplemental Fig. S6A), fixed and separated into equal portions to determine intracellular GATA3 and IgG (background) staining, and finally analyzed by flow cytometry (Supplemental Fig. S6B). Since eGFP fluorescence is diminished after fixation, we used a two-step method. We initially observed that eGFP⁻ DN4 cells express GATA3 but also that eGFP⁺ cells express much more GATA3 (resulting from the sum of the wild-type and GATA3-eGFP fusion proteins). A similar result was observed in CD4 SP stage cells (Supplemental Fig. S6C,D). We also confirmed that GATA3 mRNA is expressed at all stages in eGFP⁻ immature T cells (Supplemental Fig. S7). Based on these data, we conclude that eGFP⁻ thymocytes in *Gata3*^{g/+} animals express monoallelic *Gata3*, while eGFP⁺ cells that express GATA3 more abundantly transcribe both *Gata3* alleles.

We next asked whether the monoallelic-to-biallelic expression switch at the *Gata3* locus represents a gradual change (Fig. 3C, model A) or a bimodal event (Fig. 3C, model B). One model might posit that, as the demand for *Gata3* mRNA increases and progenitor cells mature, the total transcriptional activity from the *Gata3* gene increases in 100% of immature cells, at which point roughly half of the cells stochastically activate both al-

leles (Fig. 3C, model A). For example, some component of the machinery required for *Gata3* transcription may be limiting, and thus T cells could have an equal probability of activating either one or both *Gata3* alleles as cells traverse DN2 to DN3 maturation to allow increased GATA3 abundance. This would result in a shift of a single peak of eGFP intensity to greater overall fluorescence in the thymocytes of *Gata3*^{g/+} mice. However, the bimodal eGFP expression profile detected at the DN4 stage in *Gata3*^{g/+} thymocytes (Fig. 3B) best supports a model in which *Gata3* is transcribed from only one allele at the beginning of thymopoiesis, and, at some time during the DN2/DN3 transition, half of the cells begin to transcribe the second *Gata3* allele (Fig. 3C, model B).

To remove any possible bias between expression of the wild-type and mutant alleles, we also analyzed thymocytes recovered from homozygous *Gata3*^{g/g} fetal liver HSC-reconstituted animals; these T cells again exhibited bimodal GATA3 expression at the DN2 stage and beyond (Fig. 3D) and never exhibited a single shifted peak from lower to greater fluorescence intensity as the thymocytes matured and as would be predicted from the first model. The fact that a more robust eGFP peak in *Gata3*^{g/g} mice is first detected at the DN2 stage (Fig. 3D) is entirely consistent with the observation that the *Gata3* monoallelic-to-biallelic switch initiates around the DN2 stage in approximately one-third of all thymocytes (Fig. 2; Tables 1, 2). Thus, in *Gata3*^{g/+} thymi, we could identify cells that express biallelic *Gata3* (eGFP⁺) and monoallelic *Gata3* (eGFP⁻) at the DN4 and later stages. These data validate the conclusions that, first, *Gata3* is monoallelically transcribed in ETPs, and, second, the expression pattern switches to biallelic transcription in about half of the T cells during late thymocyte development.

To test the hypothesis that the commitment to *Gata3* monoallelic or biallelic transcriptional status is regulated by a stable mechanism, we next asked whether, after the switch in transcription occurs at DN2/DN3, late stage monoallelic cells continue to express from only one chromosome or can switch to biallelic transcription and, conversely, whether cells that express both alleles can revert to single-allele expression. To address this question, we isolated an equal number of *Gata3* biallelic (eGFP⁺) and *Gata3* monoallelic (eGFP⁻) DN4 stage thymocytes from *Gata3*^{g/+} animals (Fig. 3E, left panel) and cocultured

Figure 3. The commitment to *Gata3* monoallelic or biallelic transcriptional status is mechanistically stable. (A) A hypothetical model. The GATA3-eGFP fusion (*Gata3^g*; red) allele-expressing adult *Gata3^{g/+}* ETP cells are not viable because the reduced hypomorphic GATA3 monoallelic activity is not sufficient to allow ETPs to survive and they are eliminated, since GATA3 is vital for early T-cell generation, while wild-type (*Gata3⁺*; blue) monoallelic-expressing cells survive and continue to develop. Later during T cell development, after the DN3b stage, the second allele is activated in about half of the surviving cells and expresses both hypomorphic (*Gata3^g*) and wild-type (*Gata3⁺*) alleles. (B) Expression of a GATA3-eGFP fusion protein. eGFP expression from *Gata3^{+/+}* (open curve) and *Gata3^{g/+}* (filled curve) heterozygotes was monitored by flow cytometry in immature thymocytes to track transcription from the *Gata3^g* allele (Hosoya et al. 2009). Numbers above the bars indicate the percentage of eGFP-positive cells compared with wild-type controls from the same ancestral population. These histograms are representative of data recovered for at least five mice of each genotype from three independent experiments. (C) Hypothetical gradual (left) versus bimodal (right) switching models. (D) *Gata3* monoallelic cells expressing the *Gata3^g* allele survive and develop in the absence of the *Gata3⁺* allele. *Gata3^{g/g}* homozygous hypomorphic mutant thymocytes express eGFP in a cell-autonomous manner. *Gata3^{g/g}* or *Gata3^{+/+}* HSCs were recovered from fetal livers and adoptively transferred into sublethally irradiated adult mice. Sixteen weeks after HSC transplantation, thymocytes were analyzed for eGFP expression. The diagram presents representative histograms in which seven to eight recipients were examined for each genotype in two independent experiments. (E) DN4 stage thymocytes were sorted into eGFP-negative (eGFP⁻; open curve) or eGFP⁺ (filled curve) populations from *Gata3^{g/+}* mice and then cocultured on OP9-DL1 feeder cells. The eGFP expression profiles depicted immediately after sorting (left panels) or after 4 d of coculture and development into DP stage cells (right panels) are shown. The diagrams are representative of a total of eight mice examined in two independent experiments.

them separately on OP9-DL1 feeder cells to induce T-cell development to the DP stage. After 4 d of coculture, a majority of DN4 stage cells develop to the DP stage. DP cells that had developed from biallelic (eGFP⁺) DN4 cells did not exhibit a bimodal expression pattern (i.e., none of the cells reverted to monoallelic transcription), although eGFP intensity was significantly reduced (Fig. 3E, bottom right panel), as was also observed *in vivo* (Hosoya et al. 2009). This reduction in intensity of fluorescence is probably attributable to dilution of the fluorochrome that results from the extensive proliferation during culture for 4 d (see Fig. 1A for the remarkable increase in cell number after 4 d of coculture; Supplemental Fig. S8). Similarly, DP cells that were differentiated from monoallelic (eGFP⁻) DN4 stage cells did not develop a bimodal expression pattern (i.e., they failed to switch to biallelic transcription) (Fig. 3E, top right panel). Both patterns were consistent

over all 4 d of coculture (data not shown). These data demonstrate that once cells are set to express either one or both *Gata3* alleles at the DN2/DN3 stage, they stably maintain their monoallelic or biallelic *Gata3* transcriptional identity during T-cell development at least to the DP stage.

Maintenance of the *Gata3* monoallelic-to-biallelic switch

To explore one aspect of the mechanism that might regulate the early thymocyte *Gata3* monoallelic-to-biallelic switch, we performed chromatin immunoprecipitation (ChIP)-qPCR in the *Gata3* locus to determine the methylation status of two key epigenetic marks: H3K27me3 and H3K4me3. H3K27me3 is a histone mark for gene repression (Plath et al. 2003; Umlauf et al. 2004) and is often found near monoallelically expressed genes (Nag et al. 2013), while H3K4me3 modifications are commonly

coincident with actively transcribed genes (Bernstein et al. 2005; Kim et al. 2005). Therefore, we hypothesized that H3K27me3 could be observed in or near the *Gata3* locus in monoallelic (eGFP⁻) *Gata3*^{g/+} DN4 stage cells but not biallelic eGFP⁺ cells (since both alleles would be active). Similarly, we speculated that the H3K4me3-activating mark would be higher in the neighborhood of the *Gata3* locus in biallelic eGFP⁺ cells (from *Gata3*^{g/+} DN4 stage) when compared with monoallelic eGFP⁻ cells. Since H3K27me3 ChIP-seq (ChIP combined with deep sequencing) data for DN3 and DP stage cells have been reported (Zhang et al. 2012), PCR primer sets for ChIP-qPCR were designed that would detect the H3K27me3 mark that was previously observed 5' to the *Gata3* gene in DN3 and DP cells (Supplemental Fig. S9). Unexpectedly, we found that there was no enrichment of the H3K27me3 mark near the *Gata3* locus in monoallelic eGFP⁻ cells when compared with (biallelic) eGFP⁺ cells (Supplemental Fig. S9). We also found no difference in the activating H3K4me3 mark in these two cell populations (Supplemental Fig. S9). These data suggest that these two specific histone modifications do not play an active role in maintaining the silencing of the inactive locus in lymphocytes that transcribe only one *Gata3* allele. However, the data do not address the possibility that other epigenetic modifications may be involved in the switch.

Monoallelic or biallelic expression in Th2 or nonhematopoietic cells

Circulating Th2 cells express the highest levels of GATA3 in T lymphocytes, much more abundantly than in any thymocyte. We wondered whether those cells all express both alleles of *Gata3*, or perhaps the increased GATA3 levels were generated simply by increased transcription of a single allele. The bimodal eGFP profile in *Gata3*^{g/+} Th2 cells (Hosoya-Ohmura et al. 2011) suggests that both *Gata3* monoallelic expression and biallelic expression are observed in Th2 stage lymphocytes and thus that any increase in GATA3 expression at the Th2 stage is achieved by augmented transcription from either one or both *Gata3* loci but additionally that the monoallelic-to-biallelic switch remains stable and intact.

Gata3 is vital for not only T-cell development but also the generation or maintenance of multiple other tissues; for example, the sympathetic nervous system (Lim et al. 2000; Moriguchi et al. 2006), kidney development (Hasegawa et al. 2007), breast epithelial development (Kouros-Mehret et al. 2006), and a growing list of other organs. This provoked the question of whether *Gata3* might also be monoallelically transcribed in tissues other than T cells. To examine this hypothesis, we interrogated *Gata3* expression in embryos in tissues where prominent *Gata3* expression has been reported previously (Hasegawa et al. 2007).

When we originally generated the *Gata3*^g allele, a full-length mouse GATA3 cDNA was fused in-frame to the eGFP cDNA (Hosoya et al. 2009). In *Gata3*^{g/+} tissues, we could easily visualize individual cells expressing the chimeric GATA3-gfp fusion protein using an anti-GFP antibody. However, this would not allow distinction of cells

that solely expressed the wild-type *Gata3* allele, since an anti-GATA3 antibody would recognize both the wild-type and chimeric GATA3 proteins. Given this technical constraint, we therefore attempted to address whether monoallelic or biallelic GATA3 expression occurred in tissues other than T cells by breeding together two mutant alleles: one a *lacZ* knock-in null allele (directed by normal *Gata3* regulatory elements) (van Doorninck et al. 1999) and the second the hypomorphic *Gata3*^g allele. *Gata3*^{g/z} embryos did not survive to late gestation (K-C Lim and C-J Ku, unpubl.).

Cells lining the nephric duct (Supplemental Fig. S10, top panels, dashed circle) of a wild-type embryonic day 10.5 (E10.5) *Gata3* embryo were devoid of fluorescence (except for autofluorescing erythrocytes) (Supplemental Fig. S10, bottom right), demonstrating the specificity of both the anti-GFP and anti-β-galactosidase (anti-β-gal) antibodies used in these experiments. In an E10.5 *Gata3*^{g/z} littermate (Supplemental Fig. S10, middle panels), all nephric duct cells displayed immunoreactivity with both the anti-GFP and anti-β-gal antibodies, indicating exclusive biallelic GATA3 expression in the E10.5 nephric duct. However, we also observed that, while most cells coexpressed both the *Gata3*^g and *Gata3*^z alleles (for example, in E10.5 branchial arches [BAs] [data not shown] and the V2 interneurons of the E13.5 developing spinal cord [SC] [Supplemental Fig. S10, bottom panels]), a few cells appeared to be very faintly stained for either GFP or β-gal; however, upon close inspection, every cell that labeled with the GFP antibody also stained with the anti-β-gal antibody. This was also true when we immunostained *Gata3*^{z/+} adult bladders and kidneys (data not shown) with anti-GATA3 and anti-β-gal antibodies. We tentatively concluded that, if *Gata3* is monoallelically expressed in the embryonic urothelium, sympathetic ganglia, or BAs or adult bladders or kidneys, such monoallelic *Gata3*-expressing cells are, at most, very rare.

Discussion

We showed previously that cells can be quite sensitive to modest changes in transcription factor abundance (Motohashi et al. 2000; Shimizu et al. 2004; Hasegawa et al. 2007). Here, we showed that an increase in GATA3 protein and transcript abundance accompanies the monoallelic-to-biallelic genetic switch, which appears to be biologically required for both the repression of PU.1 and apoptosis during thymic T-cell development. Since progression through early T-cell development is exquisitely sensitive to GATA3 abundance (Scripture-Adams et al. 2014) and since forced expression of GATA3 in ETP to DN2 stage thymocytes leads to thymic lymphoma (Nawijn et al. 2001), diploid organisms appear to have evolved a mechanism by which they can exquisitely control gene expression by regulating monoallelic versus biallelic transcription. Although B-cell transcription factor *Pax5* was originally reported by RNA-FISH to be monoallelically expressed (Nutt et al. 1999), this was later disproven using a more sensitive single-cell RT-PCR assay (Rhoades et al. 2000). Here, we used both methods to ensure the veracity

of the results. We propose that both *Gata3* T-cell-specific enhancer-mediated regulation (Hosoya-Ohmura et al. 2011) and the monoallelic-to-biallelic switch must collaborate to achieve precise control of transcription factor GATA3 abundance during T-cell development.

Why is the switch to biallelic expression observed in only half of the cells while the other half remain stably monoallelic? The most intriguing hypothesis concerns the correlation with allelic exclusion at the *Tcr β* locus, which happens during the DN3 stage within the same differentiation window in which cells must surmount β selection. Another viable hypothesis is that DN3 stage thymocytes already “know” whether they are committed to a CD4 SP versus CD8 SP cell fate, since GATA3 is more abundantly expressed in CD4 SP cells than in CD8 SP cells and is required for CD4 SP cell development. We are in the process of testing these and several alternative hypotheses at the present time.

With the knowledge that the monoallelic-to-biallelic switch of a tissue-restricted transcription factor, GATA3, is intimately involved in several different aspects of thymic T-cell development, this raises the question of whether *Gata3* monoallelic or biallelic transcriptional status mediates processes in other tissues. While *Gata3* monoallelic or biallelic transcription in the many other cell types and organ systems where it is normally expressed has not yet been analyzed (e.g., in the CNS, peripheral nervous system [PNS], and kidneys) (Yamamoto et al. 1990; George et al. 1994; Lim et al. 2000; Moriguchi et al. 2006), *Gata3* haploinsufficiency is known to lead to variably penetrant HDR syndrome in humans as well as other partially penetrant pathophysiological effects (Van Esch et al. 2000; Usary et al. 2004; Grigorieva et al. 2010). We therefore speculate that *Gata3* monoallelic expression and/or a monoallelic-to-biallelic switch in tissues other than T cells could play critical roles in their normal development or homeostatic maintenance. It will be extremely interesting to determine, for example, whether the quite variable kidney dysplasia observed in human HDR patients might arise as a consequence of early monoallelic expression of GATA3 followed by apoptotic failure of a variable number of cells in this developing organ, since this aspect of the disease can be readily modeled in the kidneys of mice (Hasegawa et al. 2007).

It remains to be determined how frequently similar genetic switch mechanisms are employed among the growing number of monoallelic autosomal genes that have been documented in diploid organisms. Although *Nanog* was originally reported to be monoallelic in early pre-implantation embryos, with the proportion of biallelism increasing progressively from 2% at the eight-cell stage to 70% of the inner cells in the late blastocyst (Miyanari and Torres-Padilla 2012), that conclusion has been challenged recently (Faddah et al. 2013; Filipczyk et al. 2013). Furthermore, the active allelic selection appears to be reversible during culture of embryonic stem cells for 6 d (Chambers et al. 2007). In contrast, *Gata3* monoallelic (eGFP⁺) cells stably maintain expression from the active allele during 4 d of in vitro culture as the cells developed from the DN4 stage to the DP stage. The imprinted

gene *Kcnq1* (also known as *Kvlqt1*) is expressed from only the maternal allele early during embryogenesis, while the paternal allele becomes active as this gene acquires tissue-specific expression (Lee et al. 1997; Gould and Pfeifer 1998); in contrast, *Gata3* monoallelic expression is not parent of origin-dependent. These differences suggest that the mechanism that regulates the *Gata3* monoallelic-to-biallelic expression switch is fundamentally different from the mechanism exploited by these genes. If this phenomenon is prevalent, the number of genes controlled by monoallelic regulation could be even greater than previously estimated (8%–24%) simply because all of the genes that are expressed in a monoallelic fashion might not have been detected under the specific experimental conditions used (Gimelbrant et al. 2007; Deng et al. 2014). Given the present findings coupled with the fact that there are a large number of autosomal genes that are monoallelically expressed, it does not seem unreasonable to suggest that defects in monoallelic-to-biallelic conversion in some subset of these could contribute to many variably penetrant phenotypes that result from human haploinsufficiency.

Materials and methods

Mice

The *Gata3*^z allele (van Doorninck et al. 1999) and *Gata3*^g allele (Hosoya et al. 2009) have been described previously and were maintained on a B6 genetic background. Wild-type B6 mice and FVB mice were purchased from Jackson Laboratory. All mice were analyzed at 6–8 wk of age after backcrossing onto the B6 background for at least seven generations. All mice used in this study were housed in the Unit for Laboratory Animal Medicine at the University of Michigan, and all experiments were approved by the University Committee on Use and Care of Animals.

Cell preparation and flow cytometry

Cells were analyzed and sorted by flow cytometry (LSRFortessa and FACSAria III, BD Biosciences) as previously described (Hosoya et al. 2009; Ku et al. 2012). Sorted cells were reanalyzed to confirm that they were of >95% purity before subsequent study. Thymocytes were resuspended in propidium iodide (Calbiochem) or DAPI (Roche) solution to discriminate between live and dead cells. Depending on the specific analysis, the following antibodies were used: B220 (RA3-6B2), CD2 (RM2-5), CD3 (17A2), CD4 (RM4-5), CD5 (53-7.3), CD8 (53-6.7), CD11c (N418), CD19 (6D5), CD25 (PC61.5), CD27 (LG.7F9), CD41 (MWReg30), CD48 (HM48.1), CD150 (TC15-12F12.2), c-Kit (2B8), GATA3 (16E10A23), Gr1 (RB6-8C5), Mac-1 (M1/70), NK1.1 (PK136), Sca-1 (D7), TCR β (H57-597), TCR $\gamma\delta$ (GL3), and TER119 (TER-119). All antibodies were purchased from BD Biosciences, BioLegend, or eBioscience. After cell surface staining, apoptosis was analyzed using Annexin V and DAPI staining in Annexin V-binding buffer, per the manufacturer's instructions (BD Pharmingen). Data were analyzed using FlowJo software (Tree Star) or FACSDiva software (BD Biosciences).

OP9-DL1 coculture

Confluent OP9-DL1 stromal cells were prepared prior to coculture initiation. DN4 stage thymocytes (2×10^5) were cultured

on a feeder layer of OP9-DL1 cells (Schmitt and Zuniga-Pflucker 2002) in the presence of 5 ng/mL IL7 and 5 ng/mL Flt3 ligand (R&D Systems). Suspension and semiadherent cells were harvested by pipetting and stained for flow cytometry for four consecutive days after coculture initiation.

qRT-PCR

Total RNA was isolated from sorted cells by RNeasy micro kit (Qiagen) with DNase treatment and used to synthesize cDNA with SuperScript III first strand synthesis kit (Invitrogen). qRT-PCR was performed with SYBR Green dye using the StepOne real-time PCR system (Applied Biosystems). The primers used were previously described as follows: *GATA3* (Yu et al. 2009), *HPRT* (Maillard et al. 2008), *PU.1* (Sarrazin et al. 2009), and β -actin (Esteghamat et al. 2013). All results of qRT-PCR were performed using the total RT product from 100 cell equivalents and quantified relative to the mRNA expression of endogenous reference genes *Hprt* and β -actin. Data normalized to *HPRT* as the control are shown in the figures; data normalized to β -actin (data not shown) exhibited a similar tendency.

Allele-specific single-cell RT-PCR

Allele-specific single-cell RT-PCR is presented diagrammatically in Supplemental Figure S3. cDNA synthesis was carried out by reverse transcription using the iScript cDNA synthesis kit (Bio-Rad). A SNP was identified in the *Gata3* third exon (rs27150232), which allows discrimination between alleles of the *Mus musculus* inbred strains B6 and FVB. First, flow-sorted cell populations from F1 hybrid (B6 $\times$ FVB) mice were resorted as single cells into 1.8 μ L of 0.14% Triton X-100 (Sigma-Aldrich) followed by the addition of 0.5 μ L of 5 $\times$ iScript reaction mix and 0.2 μ L of iScript reverse transcriptase, resulting in a 2.5- μ L RT reaction volume. This was then incubated for 5 min at 25°C, 30 min at 42°C, and 5 min at 85°C per the manufacturer's instructions. cDNA was amplified by polymerase chain reaction in a 25- μ L reaction mixture consisting of 2.5 μ L of cDNA, 1 $\times$ PCR buffer (Qiagen), 200 μ M dNTPs, 200 nmol of each primer, and 0.75 U of Taq DNA polymerase (Qiagen). The PCR condition was 5 min at 95°C followed by 42 cycles of 20 sec at 95°C, 30 sec at 55°C, and 45 sec at 72°C and then a final extension for 7 min at 72°C. Primers used were 5'-GAGGTGGACGTACTTTTAAACATCG-3' and 5'-CGGGAAGGTGAAGAGATGAGG-3'. β -Actin (Esteghamat et al. 2013) was used as a control to confirm for the presence of a cell in each well and the efficiency of single-cell sorting, which was 65% $\pm$ 15% overall. Ten percent to 20% of the PCR products were visualized on a 2% agarose gel, and positives were then purified using QIAquick gel kit (Qiagen) or GeneJET gel extraction kit (Fisher Scientific) and sequenced on an Applied Biosystems 3730XL DNA sequencer at the University of Michigan Sequencing Core facility. The primer used for sequencing was 5'-ATGG CGGCAAAGCCCTGAGC-3'.

RNA-FISH

RNA-FISH was performed as described previously (Maclary et al. 2014). In detail, RNA probes were prepared by randomly priming DNA templates using the BioPrime DNA labeling system (Invitrogen). Probes were labeled with FITC-dUTP (Agilent) or Cy3-dCTP (GE Healthcare). Labeled probes were column-purified (GE Healthcare) and then precipitated in 0.3 M sodium acetate with 300 μ g of yeast transfer RNA (Invitrogen) and 150 μ g of sheared, boiled salmon sperm DNA (Invitrogen). The probe solution was spun down at 4°C and then washed in 70% ethanol followed by 100% ethanol. The pellet was dried and resuspended

in deionized formamide (Amresco). The resuspended probe was denatured for 10 min at 95°C followed immediately by a 5-min incubation on ice. A 2 $\times$ hybridization solution (4 $\times$ SSC, 20% Dextran sulfate [Millipore]) was added to the denatured probe solution and stored at -20°C until use. The *GATA3* (WIBR1-2318H06) and receptor-type tyrosine protein phosphatase C (WIBR1-2452A19; CD45) clones used to prepare probes were purchased from Children's Hospital Oakland Research Institute BACPAC Resource Center. Cells deposited on coverslips were fixed in freshly prepared cold 4% paraformaldehyde (Sigma-Aldrich) for 10 min, washed in PBS followed by permeabilization on ice in PBS with 0.5% Triton X-100 and 10 mM Ribonucleoside Vanadyl Complex (New England Biolabs) for 8 min, and then stored in 70% ethanol for up to 2 wk at 4°C before use. Samples were dehydrated through 2-min incubations in 70%, 85%, 95%, and 100% ethanol, subsequently air-dried, and then hybridized to the probe overnight at 37°C in a humid chamber (humidity provided by 2 $\times$ SSC/50% formamide). After hybridization, samples were washed three times with 2 $\times$ SSC/50% formamide for 7 min each at 39°C while shaking, twice with 2 $\times$ SSC, and twice with 1 $\times$ SSC. A 1 $\times$ DAPI solution was added to the first 1 $\times$ SSC wash. Cells were then mounted in ProLong Gold anti-fade (Invitrogen) and imaged using a 60 $\times$ (air) or 100 $\times$ (oil) objective on Nikon Eclipse TiE inverted microscope with a Photometrics CCD camera. Neuroblastoma cell line C1300 (George et al. 1994) was used as a positive control for *GATA3* expression and as a negative control for CD45 expression. All images were deconvolved and processed using NIS-Elements software.

Immunohistochemistry

Embryos and adult tissues were fixed in 4% paraformaldehyde for 1–2 h at 4°C and then equilibrated in 30% sucrose prior to cryosectioning. Sagittal sections (10 μ m) of *Gata3*^{+/+} and *Gata3*^{8/z} embryos (E10.5 and E13.5) were first stained with rabbit anti- β -gal antibody (1:2000) (Moriguchi et al. 2006) for 2 h at room temperature followed by goat anti-rabbit Cy3-conjugated secondary antibody (1:400; Invitrogen) for 30 min at room temperature. These sections were then incubated overnight at 4°C with rabbit anti-GFP Alexa fluor 488 antibody (1:1500; Invitrogen). Sagittal sections (10 μ m) of adult *Gata3*^{z/+} bladders and kidneys were similarly treated sequentially with rabbit anti- β -gal antibody followed by goat anti-rabbit Alexa fluor 488-conjugated secondary antibody (1:400; Invitrogen) and then incubated overnight at 4°C with mouse anti-GATA3 Alexa fluor 647 antibody (1:3; BD Bioscience). All tissue sections were counterstained with DAPI and then imaged using a Nikon A1 confocal microscope in the University of Michigan Microscopy and Image Analysis Laboratory.

Statistical analysis

Statistical significance was determined by Student *t*-test. Data were considered statistically significant when *P* < 0.05.

Acknowledgments

We are grateful to Stephen J. Weiss for numerous thoughtful clarifying comments, Thomas Graf for posing a significant question addressed here, and Maria Figueroa and the University of Michigan Epigenetics Core for assistance with the histone ChIP assays. This work was supported by National Institute of Health grants AI094642 (to T.H. and J.D.E.) and AI091627 (to I.M.), the Leukemia and Lymphoma Society (I.M.), and the Lymphoma Research Foundation (C.-J.K.). The research was also supported in part by the National Cancer Institute through the University of

Michigan Comprehensive Cancer Center Support Grant (P30 CA046592) and the use of the Flow Cytometry, Morphology, Microscopy and Image Analysis Laboratory, Experimental Irradiation, and Sequencing cores at University of Michigan. C.-J.K. designed the study, performed experiments, analyzed the data, and wrote the manuscript. K.-C.L. designed the study, performed experiments, and edited the manuscript. S.K. designed the FISH study and edited the manuscript. I.M. designed the study and edited the manuscript. J.D.E. designed the study, analyzed the data, and wrote the paper. T.H. designed the study, performed experiments, analyzed the data, and wrote the paper.

References

- Allman D, Sambandam A, Kim S, Miller JP, Pagan A, Well D, Meraz A, Bhandoola A. 2003. Thymopoiesis independent of common lymphoid progenitors. *Nat Immunol* **4**: 168–174.
- Bartolomei MS, Zemel S, Tilghman SM. 1991. Parental imprinting of the mouse H19 gene. *Nature* **351**: 153–155.
- Bell JJ, Bhandoola A. 2008. The earliest thymic progenitors for T cells possess myeloid lineage potential. *Nature* **452**: 764–767.
- Bernstein BE, Kamal M, Lindblad-Toh K, Bekiranov S, Bailey DK, Huebert DJ, McMahon S, Karlsson EK, Kulbokas EJ III, Gingeras TR, et al. 2005. Genomic maps and comparative analysis of histone modifications in human and mouse. *Cell* **120**: 169–181.
- Chambers I, Silva J, Colby D, Nichols J, Nijmeijer B, Robertson M, Vrana J, Jones K, Grothowold L, Smith A. 2007. Nanog safeguards pluripotency and mediates germline development. *Nature* **450**: 1230–1234.
- David-Fung E-S, Yui MA, Morales M, Wang H, Taghon T, Diamond RA, Rothenberg EV. 2006. Progression of regulatory gene expression states in fetal and adult pro-T-cell development. *Immunol Rev* **209**: 212–236.
- Daw SC, Taylor C, Kraman M, Call K, Mao J, Schuffenhauer S, Meitinger T, Lipson T, Goodship J, Scambler P. 1996. A common region of 10p deleted in DiGeorge and velocardiofacial syndromes. *Nat Genet* **13**: 458–460.
- DeChiara TM, Robertson EJ, Efstratiadis A. 1991. Parental imprinting of the mouse insulin-like growth factor II gene. *Cell* **64**: 849–859.
- Deng Q, Ramskold D, Reinius B, Sandberg R. 2014. Single-cell RNA-seq reveals dynamic, random monoallelic gene expression in mammalian cells. *Science* **343**: 193–196.
- Eckersley-Maslin MA, Spector DL. 2014. Random monoallelic expression: regulating gene expression one allele at a time. *Trends Genet* **30**: 237–244.
- Esteghamat F, Gillemans N, Bilic I, van den Akker E, Cantu I, van Gent T, Klingmuller U, van Lom K, von Lindern M, Grosveld F, et al. 2013. Erythropoiesis and globin switching in compound Klf1::Bcl11a mutant mice. *Blood* **121**: 2553–2562.
- Faddah DA, Wang H, Cheng AW, Katz Y, Buganim Y, Jaenisch R. 2013. Single-cell analysis reveals that expression of nanog is biallelic and equally variable as that of other pluripotency factors in mouse ESCs. *Cell Stem Cell* **13**: 23–29.
- Filipczyk A, Gkatzis K, Fu J, Hoppe PS, Lickert H, Anastassiadis K, Schroeder T. 2013. Biallelic expression of nanog protein in mouse embryonic stem cells. *Cell Stem Cell* **13**: 12–13.
- Fowlkes BJ, Mathieson BJ. 1985. Intrathymic differentiation: thymocyte heterogeneity and the characterization of early T-cell precursors. *Surv Immunol Res* **4**: 96–109.
- George KM, Leonard MW, Roth ME, Lieu KH, Kioussis D, Grosveld F, Engel JD. 1994. Embryonic expression and cloning of the murine GATA-3 gene. *Development* **120**: 2673–2686.
- Gimelbrant A, Hutchinson JN, Thompson BR, Chess A. 2007. Widespread monoallelic expression on human autosomes. *Science* **318**: 1136–1140.
- Gould TD, Pfeifer K. 1998. Imprinting of mouse Kvlqt1 is developmentally regulated. *Hum Mol Genet* **7**: 483–487.
- Grigorieva IV, Mirczuk S, Gaynor KU, Nesbit MA, Grigorieva EF, Wei Q, Ali A, Fairclough RJ, Stacey JM, Stechman MJ, et al. 2010. Gata3-deficient mice develop parathyroid abnormalities due to dysregulation of the parathyroid-specific transcription factor Gcm2. *J Clin Invest* **120**: 2144–2155.
- Guo L, Hu-Li J, Paul WE. 2005. Probabilistic regulation in TH2 cells accounts for monoallelic expression of IL-4 and IL-13. *Immunity* **23**: 89–99.
- Hasegawa SL, Moriguchi T, Rao A, Kuroha T, Engel JD, Lim K-C. 2007. Dosage-dependent rescue of definitive nephrogenesis by a distant Gata3 enhancer. *Dev Biol* **301**: 568–577.
- Hassold T, Hunt P. 2001. To err (meiotically) is human: the genesis of human aneuploidy. *Nat Rev Genet* **2**: 280–291.
- Hernández-Hoyos G, Anderson MK, Wang C, Rothenberg EV, Alberola-Ila J. 2003. GATA-3 expression is controlled by TCR signals and regulates CD4/CD8 differentiation. *Immunity* **19**: 83–94.
- Hollander GA, Zuklys S, Morel C, Mizoguchi E, Mobisson K, Simpson S, Terhorst C, Wishart W, Golan DE, Bhan AK, et al. 1998. Monoallelic expression of the interleukin-2 locus. *Science* **279**: 2118–2121.
- Hosoya T, Kuroha T, Moriguchi T, Cummings D, Maillard I, Lim KC, Engel JD. 2009. GATA-3 is required for early T lineage progenitor development. *J Exp Med* **206**: 2987–3000.
- Hosoya T, Maillard I, Engel JD. 2010. From the cradle to the grave: activities of GATA-3 throughout T-cell development and differentiation. *Immunol Rev* **238**: 110–125.
- Hosoya-Ohmura S, Lin YH, Herrmann M, Kuroha T, Rao A, Moriguchi T, Lim KC, Hosoya T, Engel JD. 2011. An NK and T cell enhancer lies 280 kilobase pairs 3' to the gata3 structural gene. *Mol Cell Biol* **31**: 1894–1904.
- Hume DA. 2000. Probability in transcriptional regulation and its implications for leukocyte differentiation and inducible gene expression. *Blood* **96**: 2323–2328.
- Inoue K, Lupski JR. 2002. Molecular mechanisms for genomic disorders. *Annu Rev Genomics Hum Genet* **3**: 199–242.
- Kim TH, Barrera LO, Zheng M, Qu C, Singer MA, Richmond TA, Wu Y, Green RD, Ren B. 2005. A high-resolution map of active promoters in the human genome. *Nature* **436**: 876–880.
- Ko LJ, Yamamoto M, Leonard MW, George KM, Ting P, Engel JD. 1991. Murine and human T-lymphocyte GATA-3 factors mediate transcription through a cis-regulatory element within the human T-cell receptor δ gene enhancer. *Mol Cell Biol* **11**: 2778–2784.
- Kouros-Mehr H, Slorach EM, Sternlicht MD, Werb Z. 2006. GATA-3 maintains the differentiation of the luminal cell fate in the mammary gland. *Cell* **127**: 1041–1055.
- Ku CJ, Hosoya T, Maillard I, Engel JD. 2012. GATA-3 regulates hematopoietic stem cell maintenance and cell-cycle entry. *Blood* **119**: 2242–2251.
- Kuznetsov VA, Knott GD, Bonner RF. 2002. General statistics of stochastic process of gene expression in eukaryotic cells. *Genetics* **161**: 1321–1332.
- Lee MP, Hu RJ, Johnson LA, Feinberg AP. 1997. Human KVLQT1 gene shows tissue-specific imprinting and encompasses Beckwith-Wiedemann syndrome chromosomal rearrangements. *Nat Genet* **15**: 181–185.
- Lichtner P, König R, Hasegawa T, Van Esch H, Meitinger T, Schuffenhauer S. 2000. An HDR (hypoparathyroidism, deafness, renal dysplasia) syndrome locus maps distal to the

- DiGeorge syndrome region on 10p13/14. *J Med Genet* **37**: 33–37.
- Lim KC, Lakshmanan G, Crawford SE, Gu Y, Grosveld F, Engel JD. 2000. *Gata3* loss leads to embryonic lethality due to nor-adrenaline deficiency of the sympathetic nervous system. *Nat Genet* **25**: 209–212.
- Lyon MF. 1961. Gene action in the X-chromosome of the mouse (*Mus musculus* L.). *Nature* **190**: 372–373.
- Maclary E, Buttigieg E, Hinten M, Gayen S, Harris C, Sarkar MK, Purushothaman S, Kalantry S. 2014. Differentiation-dependent requirement of Tsix long non-coding RNA in imprinted X-chromosome inactivation. *Nat Commun* **5**: 4209.
- Maillard I, Koch U, Dumortier A, Shestova O, Xu L, Sai H, Pross SE, Aster JC, Bhandoola A, Radtke F, et al. 2008. Canonical notch signaling is dispensable for the maintenance of adult hematopoietic stem cells. *Cell Stem Cell* **2**: 356–366.
- Miyanari Y, Torres-Padilla ME. 2012. Control of ground-state pluripotency by allelic regulation of Nanog. *Nature* **483**: 470–473.
- Moriguchi T, Takako N, Hamada M, Maeda A, Fujioka Y, Kuroha T, Huber RE, Hasegawa SL, Rao A, Yamamoto M, et al. 2006. *Gata3* participates in a complex transcriptional feedback network to regulate sympathoadrenal differentiation. *Development* **133**: 3871–3881.
- Motohashi H, Katsuoka F, Shavit JA, Engel JD, Yamamoto M. 2000. Positive or negative MARE-dependent transcriptional regulation is determined by the abundance of small Maf proteins. *Cell* **103**: 865–875.
- Nag A, Savova V, Fung HL, Miron A, Yuan GC, Zhang K, Gimelbrant AA. 2013. Chromatin signature of widespread monoallelic expression. *eLife* **2**: e01256.
- Nawijn MC, Ferreira R, Dingjan GM, Kahre O, Drabek D, Karis A, Grosveld F, Hendriks RW. 2001. Enforced expression of *GATA-3* during T cell development inhibits maturation of CD8 single-positive cells and induces thymic lymphoma in transgenic mice. *J Immunol* **167**: 715–723.
- Nerlov C, Graf T. 1998. PU.1 induces myeloid lineage commitment in multipotent hematopoietic progenitors. *Genes Dev* **12**: 2403–2412.
- Nutt SL, Vambrie S, Steinlein P, Kozmik Z, Rolink A, Weith A, Busslinger M. 1999. Independent regulation of the two Pax5 alleles during B-cell development. *Nat Genet* **21**: 390–395.
- Ohno S, Kaplan WD, Kinoshita R. 1959. Formation of the sex chromatin by a single X-chromosome in liver cells of *Rattus norvegicus*. *Exp Cell Res* **18**: 415–418.
- Pai S-Y, Truitt ML, Ting C-N, Leiden JM, Glimcher LH, Ho I-C. 2003. Critical roles for transcription factor *GATA-3* in thymocyte development. *Immunity* **19**: 863–875.
- Penit C, Vasseur F. 1988. Sequential events in thymocyte differentiation and thymus regeneration revealed by a combination of bromodeoxyuridine DNA labeling and antimitotic drug treatment. *J Immunol* **140**: 3315–3323.
- Pereira JP, Girard R, Chaby R, Cumano A, Vieira P. 2003. Monoallelic expression of the murine gene encoding Toll-like receptor 4. *Nat Immunol* **4**: 464–470.
- Plath K, Fang J, Mlynarczyk-Evans SK, Cao R, Worringer KA, Wang H, de la Cruz CC, Otte AP, Panning B, Zhang Y. 2003. Role of histone H3 lysine 27 methylation in X inactivation. *Science* **300**: 131–135.
- Porritt HE, Rummelt LL, Tabrizifard S, Schmitt TM, Zuniga-Pflucker JC, Petrie HT. 2004. Heterogeneity among DN1 prothymocytes reveals multiple progenitors with different capacities to generate T cell and non-T cell lineages. *Immunity* **20**: 735–745.
- Rhoades KL, Singh N, Simon I, Glidden B, Cedar H, Chess A. 2000. Allele-specific expression patterns of interleukin-2 and Pax-5 revealed by a sensitive single-cell RT-PCR analysis. *Curr Biol* **10**: 789–792.
- Sambandam A, Maillard I, Zediak VP, Xu L, Gerstein RM, Aster JC, Pear WS, Bhandoola A. 2005. Notch signaling controls the generation and differentiation of early T lineage progenitors. *Nat Immunol* **6**: 663–670.
- Sarrazin S, Mossadegh-Keller N, Fukao T, Aziz A, Mourcin F, Vanhille L, Kelly Modis L, Kastner P, Chan S, Duprez E, et al. 2009. MafB restricts M-CSF-dependent myeloid commitment divisions of hematopoietic stem cells. *Cell* **138**: 300–313.
- Schmitt TM, Zuniga-Pflucker JC. 2002. Induction of T cell development from hematopoietic progenitor cells by Delta-like-1 in vitro. *Immunity* **17**: 749–756.
- Scripture-Adams DD, Damle SS, Li L, Elihu KJ, Qin S, Arias AM, Butler RR III, Champhekar A, Zhang JA, Rothenberg EV. 2014. *GATA-3* dose-dependent checkpoints in early T cell commitment. *J Immunol* **193**: 3470–3491.
- Shi J, Petrie HT. 2012. Activation kinetics and off-target effects of thymus-initiated cre transgenes. *PLoS One* **7**: e46590.
- Shimizu R, Ohneda K, Engel JD, Trainor CD, Yamamoto M. 2004. Transgenic rescue of *GATA-1*-deficient mice with *GATA-1* lacking a FOG-1 association site phenocopies patients with X-linked thrombocytopenia. *Blood* **103**: 2560–2567.
- Tan JB, Visan I, Yuan JS, Guidos CJ. 2005. Requirement for Notch1 signals at sequential early stages of intrathymic T cell development. *Nat Immunol* **6**: 671–679.
- Umlauf D, Goto Y, Cao R, Cerqueira F, Wagschal A, Zhang Y, Feil R. 2004. Imprinting along the Kcnq1 domain on mouse chromosome 7 involves repressive histone methylation and recruitment of Polycomb group complexes. *Nat Genet* **36**: 1296–1300.
- Usary J, Llaca V, Karaca G, Presswala S, Karaca M, He X, Langerod A, Karsen R, Oh DS, Dressler LG, et al. 2004. Mutation of *GATA3* in human breast tumors. *Oncogene* **23**: 7669–7678.
- van Doorninck JH, van Der Wees J, Karis A, Goedknecht E, Engel JD, Coesmans M, Rutteman M, Grosveld F, De Zeeuw CI. 1999. *GATA-3* is involved in the development of serotonergic neurons in the caudal raphe nuclei. *J Neurosci* **19**: RC12.
- Van Esch H, Groenen P, Nesbit MA, Schuffenhauer S, Lichtner P, Vanderlinden G, Harding B, Beetz R, Bilous RW, Holdaway I, et al. 2000. *GATA3* haplo-insufficiency causes human HDR syndrome. *Nature* **406**: 419–422.
- Yamamoto M, Ko LJ, Leonard MW, Beug H, Orkin SH, Engel JD. 1990. Activity and tissue-specific expression of the transcription factor NF-E1 multigene family. *Genes Dev* **4**: 1650–1662.
- Yu Q, Sharma A, Oh SY, Moon H-G, Hossain MZ, Salay TM, Leeds KE, Du H, Wu B, Waterman ML, et al. 2009. T cell factor 1 initiates the T helper type 2 fate by inducing the transcription factor *GATA-3* and repressing interferon- γ . *Nat Immunol* **10**: 992–999.
- Yui MA, Rothenberg EV. 2014. Developmental gene networks: a triathlon on the course to T cell identity. *Nat Rev Immunol* **14**: 529–545.
- Zhang JA, Mortazavi A, Williams BA, Wold BJ, Rothenberg EV. 2012. Dynamic transformations of genome-wide epigenetic marking and transcriptional control establish T cell identity. *Cell* **149**: 467–482.
- Zwemer LM, Zak A, Thompson BR, Kirby A, Daly MJ, Chess A, Gimelbrant AA. 2012. Autosomal monoallelic expression in the mouse. *Genome Biol* **13**: R10.

A monoallelic-to-biallelic T-cell transcriptional switch regulates GATA3 abundance

Chia-Jui Ku, Kim-Chew Lim, Sundeep Kalantry, et al.

Genes Dev. 2015, **29**:

Access the most recent version at doi:[10.1101/gad.265025.115](https://doi.org/10.1101/gad.265025.115)

Supplemental Material

<http://genesdev.cshlp.org/content/suppl/2015/09/18/29.18.1930.DC1>

References

This article cites 69 articles, 21 of which can be accessed free at:
<http://genesdev.cshlp.org/content/29/18/1930.full.html#ref-list-1>

Creative Commons License

This article is distributed exclusively by Cold Spring Harbor Laboratory Press for the first six months after the full-issue publication date (see <http://genesdev.cshlp.org/site/misc/terms.xhtml>). After six months, it is available under a Creative Commons License (Attribution-NonCommercial 4.0 International), as described at <http://creativecommons.org/licenses/by-nc/4.0/>.

Email Alerting Service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).