Neuroectoderm in *Drosophila* embryos is dependent on the mesoderm for positioning but not for formation

Yi Rao,¹ Harald Vaessin, Lily Y. Jan, and Yuh-Nung Jan

Howard Hughes Medical Institute, Departments of Physiology and Biochemistry, University of California at San Francisco, San Francisco, California 94143-0724 USA

By studying neuroectoderm formation in the absence of mesoderm and mesectoderm in mutants of the zygotic genes *snail* and *twist*, we have found that the number of neuroblasts is not reduced in these mutants, suggesting that mesoderm and mesectoderm are not essential for the initiation of neural development. The position of the neuroectoderm, however, is ventrally shifted: Neuroectoderm takes over the presumptive peripheral mesoderm domain in single mutants, whereas the entire presumptive mesoderm domain in double mutants takes on the neuroectodermal fate. The shifted neuroectoderm still requires the proneural genes and the neurogenic genes. This shift is unlikely to be due to any shift in the nuclear localization gradient of the maternally supplied dorsal protein. A model for cell fate determination of the neuroectoderm, mesectoderm, and mesoderm will be discussed.

[Key Words: Neurogenesis; induction; mesoderm; neuroectoderm; *snail*; *twist*]

Received June 11, 1991; revised version accepted July 19, 1991.

The central nervous system of *Drosophila* originates from the ventrolateral neurogenic regions (the neuroectoderm) of the embryo (Poulson 1950; Campos-Ortega and Hartenstein 1985). In the blastoderm, neurogenic regions are separated from the mesoderm in the ventral region by one row of mesectodermal cells. During gastrulation, the mesodermal cells invaginate, eventually lying under the neurogenic, as well as the more dorsal, ectoderm, whereas the mesectodermal cells on the two sides of the embryo are brought to the ventral midline. The determination of neuroblasts requires the activity of at least two groups of genes in the neuroectoderm: the proneural genes, including *daughterless* (*da*) and genes of the *achaete-scute complex* (AS-C), which give groups of cells, the proneural clusters, the potential to become neuroblasts (Campuzano et al. 1985; Dambly-Chaudiere and Ghysen 1987; Villares and Cabrera 1987; Caudy et al. 1988; Ghysen and Dambly-Chaudiere 1988), and the neurogenic genes, such as *Notch* (*N*) or *Delta* (*Dl*), which allow only subsets of cells of the proneural clusters to become neuroblasts (Lehmann et al. 1983; Wharton et al. 1985; Kidd et al. 1986; Vaessin et al. 1987; Kopczynski et al. 1988; for reviews, see Artavanis-Tsakonas 1988; and Campos-Ortega and Jan 1991).

Whether genes expressed in non-neuroectodermal cells can affect the formation or patterning of the neuroblasts is not known. The physical proximity of the neuroectoderm to the mesoderm raises the question as

to whether the neural tissue could be induced by the mesoderm, as has been found in vertebrates (Spemann 1938; Gurdon 1987; Hamburger 1988). The presence of mesectodermal cells in the ventral midline after mesoderm invagination has further raised the question as to whether these cells function in a way analogous to those of the vertebrate floor plate cells in the patterning of the nerve cord (Klamt et al. 1991; Yamada et al. 1991). To assess the influence of mesodermal and mesectodermal cells in the formation of the embryonic nervous system of *Drosophila*, we examined the number and position of neuroblasts in mutants without mesoderm and mesectoderm.

Formation of the mesoderm requires the zygotic gene activities of *snail* (*sna*) and *twist* (*twi*), whose expression in the ventral region is controlled by the maternal gene, *dorsal* (*dl*), the morphogen for the dorsal-ventral axis (Simpson 1983; Anderson 1987). Through the action of a group of maternal-effect genes, the *dl* protein becomes localized to the nuclei in the ventral region of the embryo; the concentration of nuclear *dl* protein is high in the presumptive mesoderm and drops sharply in the region flanking the mesodermal region (Roth et al. 1989; Rushlow et al. 1989; Steward 1989). The *twi* protein is found in cells that form the mesoderm and the adjacent mesectoderm (Thisse et al. 1988; Leptin and Grunewald 1990). Expression of *sna*, on the other hand, appears to be restricted to the presumptive mesoderm (Boulay et al. 1987; Leptin and Grunewald 1990). No mesodermal tissue forms in *sna* or *twi* mutants. Both *sna*, which has zinc-finger motifs, and *twi*, which contains a helix-

¹Present address: Department of Biochemistry and Molecular Biology, Harvard University, Cambridge, Massachusetts 02138 USA.

loop-helix domain, are putative transcription factors (Boulay et al. 1987; Thiesse et al. 1988); they probably initiate mesodermal development by controlling gene expression.

We have determined both the late and early phenotypes of *sna* and *twi* mutants in the neuroectoderm and the mesectoderm. We found that the presence of neither the mesoderm nor the mesectoderm is necessary for the formation of neuroblasts, although the position of the neuroectoderm is shifted ventrally in the absence of mesoderm and mesectoderm. Cells that normally would form the mesoderm or the mesectoderm can take on the fate of neuroectodermal cells in these mutants. Our observations indicate that the positioning of the neuroectoderm depends on zygotic genes functioning in the mesoderm and mesectoderm; most of the mutant phenotypes can be accounted for by a testable hypothesis that involves cell-autonomous and cell-nonautonomous actions of these genes.

Results

Neuronal phenotype

In wild-type embryos, the neuroectoderm gives rise to both neuroblasts and epidermoblasts. Neuroblasts delaminate from the neuroectoderm, move into the space between the rest of the neuroectoderm and the invagi-

nated mesodermal cells, and divide to produce neurons. Neurons from both sides of the embryo form the ventral nerve cord, which can be visualized immunocytochemically with mAb44C11, a monoclonal antibody that recognizes specifically the product of the *elav* gene in neuronal nuclei, or anti-HRP, a polyclonal antiserum that stains the neuronal membrane (Jan and Jan 1982; Bier et al. 1988). As shown in Figure 1, A-H, these antibodies revealed the formation of nerve cord in *twi* mutants and in *sna twi* double mutants, whereas the nervous system in *sna* mutants appeared to be deficient in the formation of the commissures and contained a gap between neurons on either side of the embryo. It is possible that some of the abnormalities revealed by these neuron-specific markers late during embryogenesis arose secondarily in a twisted embryo that failed to gastrulate properly. Nonetheless, the presence of ventral nerve cords in *twi* and *sna twi* mutants indicates that mesoderm is not required for the formation of the nerve cord.

To examine the effects of *sna* and *twi* mutations on early stages of neural development, prior to the appearance of neuron-specific markers, we looked at the pattern of neuroblasts that have just segregated from the epidermal layer by using an antiserum to the product of a gene in the 44C region of the second chromosome and an antiserum to the T8 product of the AS-C (E. Bier, H. Vaessin, L.Y. Jan, and Y.N. Jan, in prep.; M. Brand, un-

Figure 1. Neural phenotype of *sna* and *twi* mutant embryos. (A–D) Anti-*elav* staining of neuronal nuclei in stage 14 wild-type, *sna*^{II^{G05}}, *twi*^{S60}, and *sna*^{II^{G05}} *twi*^{S60} double mutants; (E–H) anti-HRP staining of neuronal membranes in stage 14 embryos; (I–L) anti-44C staining of neuroblasts in stage 9 embryos. Ventral views of embryos are shown. The anterior end is up.

publ.). The number of neuroblasts per hemisegment in the cephalic and thoracic region of the stage 9 embryo was similar in the wild type (14.4 ± 1.6 ; mean $\pm$ S.D., average of 12 embryos), *sna* mutant (14.8 ± 1.8 , 9 embryos), *twi* mutant (14.2 ± 2 , 8 embryos), and the *sna twi* double mutant (14.7 ± 1.5 , 10 embryos) (see Materials and methods). There were, however, a few extra neuroblasts around the midline in *twi* or *sna twi* mutant embryos (Fig. 1I–L). Most of the neuroblasts were arranged roughly in three rows on either side of the embryo at stage 9, similar to the positions observed in the wild-type embryo (Fig. 1I–L). A ventral gap was evident between the neuroblast-containing regions on the two sides of the embryo in single mutants of *sna* or *twi* but not in *sna twi* double mutants, even though in both single mutants and double mutants cells in the ventral region fail to invaginate and form the mesoderm (Leptin and Grunewald 1990). Thus, although the number of neuroblasts was approximately normal in various mutants, the position of cells giving rise to neuroblasts appeared abnormal. This is examined more closely using markers for the proneural clusters, which appear early when the embryo still contains only a single layer of somatic cells.

Ventral displacement of the neuroectoderm

There are ~100 cells in the circumference of a cellular blastoderm. In normal embryo, the ventral-most 15–17 cells become mesodermal and invaginate, bringing the mesectodermal cells (one on either side) to the midline. At the end of ventral invagination, the mesectodermal cells also delaminate from the ectoderm to lie between the mesoderm and ectoderm. In *sna* and *twi* mutants, mesoderm is not formed and ventral cells fail to undergo normal invagination and often remain on the surface (Leptin and Grunewald 1990). A number of observations suggest that cells of the presumptive mesoderm do not die. First, the total number of cells in the mutants is the same as in the wild type (Leptin and Grunewald 1990). Second, the ventral cells show normal morphology and nuclear staining with antibodies (Leptin and Grunewald 1990), as well as expression of specific cytoplasmic RNAs (see below). Finally, mutant embryos stained with a dye for dying cells were indistinguishable from wild-type embryos at the early stages of development (B. Hay and Y. Rao, unpubl.). As shown below, the uninvaginated presumptive mesoderm domain in the mutants has been taken over partially or completely by the neuroectoderm.

T5, T4, and T3 transcripts of AS-C are expressed initially in the proneural clusters that acquire the potential to become neural precursors (Cabrera et al. 1987; Romani et al. 1987). One or more cells in a proneural cluster will maintain the expression and go on to become a neural precursor. In the wild type, the distance between T5-staining cells on either side of the embryo is 17–20 cells. In *sna* or *twi* mutants, the distance between the neurogenic domains is reduced to ~8–12 cells (Fig. 2). In *sna twi* double mutants (Fig. 2D,H), there was no gap separating the neuroectoderm on either side of the em-

bryo. Thus, the neuroectoderm shifted ventrally to take over part—or all, in the case of *sna twi* double mutant—of the presumptive mesoderm domain. In the single mutants the ventral region that did not become neuroectoderm corresponded to the domain that would have formed the central mesoderm in wild-type embryos (Leptin and Grunewald 1990), whereas the region that would have become the peripheral mesoderm in wild-type embryos became the neuroectoderm.

Ventrally displaced neuroectoderm requires proneural and neurogenic gene function

Normal development of the nervous system is initiated with the expression of proneural genes such as those of the AS-C, which endows clusters of cells with the potential to become neuroblasts. In the absence of proneural gene function, some of the neuroblasts fail to appear (Dambly-Chaudiere and Ghysen 1987; Jimenez and Campos-Ortega 1990). Once a neuroblast has formed, it is thought to inhibit its neighboring cells from taking on the neuroblast fate (Doe and Goodman 1985). This process of lateral inhibition probably involves the function of zygotic neurogenic genes including *N*, *Dl*, *Enhancer of split* [*E(spl)*], *neuralized* (*neu*), *big brain* (*bib*), and *mastermind* (*mam*) (Lehmann et al. 1983; Wharton et al. 1985; Kidd et al. 1986; Knust et al. 1987; Vaessin et al. 1987; Artavanis-Tsakonas 1988; Kopczynski et al. 1988; Preiss et al. 1988; Yedbovnick et al. 1988; Rao et al. 1990; Boulianne et al. 1991; Heitzler and Simpson 1991). Loss of neurogenic gene activities leads to an overproduction of neuroblasts and neurons.

To determine whether the proneural genes and neurogenic genes also specify the formation of neuroblasts in *sna* and *twi* mutant embryos, we studied the expression patterns and the mutant phenotypes. As described above, the expression of proneural genes in AS-C was shifted ventrally. In addition, we found that *Dl* and *bib* were also expressed in more ventral regions in the mutants; these two neurogenic genes were normally expressed in mesectoderm, neuroectoderm, and more dorsal ectoderm but not in the mesoderm (Vaessin et al. 1987; Kopczynski et al. 1988; Haenlin et al. 1990; Rao et al. 1990). To test for the functional involvement of these genes, we looked at double mutants of *sna* and *sc^{B57}*, *twi* and *sc^{B57}* or triple mutants of *sna*, *twi*, and *sc^{B57}* (*sc^{B57}* mutation is a deletion of the entire AS-C) and found that the nervous system was reduced (Fig. 3A–D). Loss of *Dl* activity, on the other hand, led to overgrowth of the nervous system in *sna*, *twi*, or *sna twi* double mutants (Fig. 3E–H). Combinations of the *sna* or *twi* mutation with other neurogenic mutations such as *N*, *E(spl)*, or *neu* also gave neurogenic phenotype (data not shown). Thus, both the proneural and neurogenic genes function in the ventrally displaced neuroectoderm.

Mesectoderm and trachea formation in *sna* and *twi* mutants

To examine the boundary of the neuroectoderm, we looked at the mesectoderm and tracheal pits, which are

Figure 2. T5 expression in proneural clusters and neural precursors. (A) Proneural clusters in a stage 6 wild-type embryo that has just begun ventral invagination; (B–D) proneural clusters in stage 7 mutant embryos without any ventral invagination, showing that the proneural clusters on either side of the embryo are separated by a gap in *sna* (B) or *twi* (C) mutants, whereas they are contiguous in the double mutant (D). Mutant embryos can be detected at stage 7 because ventral invagination would have been completed by this stage in the wild-type embryo. To examine the extent of separation of the proneural clusters on either side of the embryo, one may compare these mutant embryos with stage 6 wild-type embryos (A), which have not yet undergone ventral invagination. (E–H) T5 expression is restricted to one or a few neural precursors in late stage 8 embryos. In the wild-type embryo (E), mesodermal and mesectodermal cells have invaginated and are thus no longer on the surface. In the double mutant (H), the figure shown appears to reveal extra neuroblasts at the ventral midline, though the number of neuroblasts that express the gene at 44C is not much increased in the mutants (see text). (I–L) Side views of late stage 8 embryos; the arrow points to invaginated mesoderm in the wild-type, which is absent in the mutant embryos.

the immediate ventral and dorsal neighbors of the neuroectoderm, respectively (Campos-Ortega and Hartenstein 1985). The mesectodermal cells can be examined with the expression pattern of the *rhomboid* (*rho*) gene and the m7 gene of the *E(spl)* locus (Knust et al. 1987; Bier et al. 1990). *rho* is initially expressed in stripes of eight cells flanking the mesoderm in a cellularizing embryo. This pattern is then modulated segmentally so that the medial-most row is *rho* positive in all the segments, whereas there are additional ectodermal cells expressing *rho* in every other segment (Fig. 4D). The expression in all segments becomes restricted to this medial-most row of cells, the mesectodermal cells, on either side of the embryo, just before the beginning of gastrulation and remains in these cells thereafter. In addition, *rho* is expressed in the tracheal pits during germ-band extension (Bier et al. 1990). m7, on the other hand, is initially expressed in stripes of about four cells on either side of the embryo and then becomes restricted to the medial-most row on either side of the embryo before the completion of cellularization (Fig. 4A) (Knust et al. 1987). The medial-most rows of *rho*-expressing cells and m7-expressing cells were identified as the mesectodermal cells because these cells were brought to the ventral midline as the mesoderm invaginated, and they subsequently delaminated to lie between the ectoderm and the mesoderm.

The maintained expression of *rho* in mesectodermal cells in germ-band-extended embryos revealed further that the *rho*-expressing ventral midline cells are the same cells that would express another mesectodermal marker, *single-minded* (*sim*) (Nambu et al. 1990).

Previous studies have shown that the number of cells expressing *sim* was increased in *sna* mutants in germ-band-extended embryos (Leptin and Grunewald 1990; Nambu et al. 1990). Consistent with these findings in late embryos, we found an increased number of *rho*-expressing cells in *sna* mutants (Fig. 4G). In *twi* mutants, there were several *rho*-expressing cells flanking a stripe of nonexpressing cells (Fig. 4H). In the double mutant, no *rho*-expressing cells were detected at the ventral midline, although *rho* was still expressed in the tracheal pits (Fig. 4I). Thus, both the *sim* expression pattern shown in previous studies and the *rho* pattern in germ-band-extended embryos revealed an expansion of mesectodermal cells in *sna* mutants, whereas no mesectodermal cells were found in the double mutant.

To examine the early steps in mesectodermal formation, we looked at m7 and *rho* expression in cellularizing (stage 5) and gastrulating (stage 6) embryos. In *sna* mutants, m7 expression was detected initially in the ventral region of the cellular blastoderm (~17–19 cells); the central 10–12 cells showed stronger expression than the pe-

Figure 3. Dependence of the ventrally shifted neuroectoderm on the function of the proneural and neurogenic genes. (A–D) Reduced nervous systems in stage 14 embryos as revealed by anti-HRP staining of *sc*^{B57} single mutant (A); *sc*^{B57} *sna*^{11G05} (B) and *sc*^{B57} *twi*^{S60} (C) double mutants; or *sc*^{B57} *sna*^{11G05} *twi*^{S60} triple mutants (D). (E–H) Hyperplasia of neural tissue as revealed by anti-HRP staining of *Dl*^{9P39} (E), *Dl*^{9P39} *sna*^{11G05} (F) and *Dl*^{9P39} *twi*^{S60} (G) double mutants; or *Dl*^{9P39} *sna*^{11G05} *twi*^{S60} triple mutants (H).

ripheral cells (Fig. 4B). The expression was then limited to the central 10–12 cells around the beginning of stage 6. In *twi* mutants, flanking the ventral region of cells that did not express m7 (6–10 cells), there were 2–4 m7-expressing cells on either side of a stage 5 embryo, as compared to 1–3 m7-expressing cells on either side of a stage 6 embryo (Fig. 4C). Although this 6- to 10-cell-wide ventral stripe showed no m7 expression at stage 5, some of the cells did express m7 at stage 6 (Fig. 4C). In *sna twi* double mutants, no mesectodermal cells were detected in stage 5 or stage 6 embryos; the two longitudinal rows of *rho*-expressing cells were absent in a late stage 5 wild-type embryo even though *rho* expression was detectable in ectodermal cells that extended from one side of the embryo to the other side (Fig. 4E). The absence of m7 staining was also confirmed by in situ hybridization with both m7 and T5 probes, which showed an expression pattern of only T5 but not m7 in *sna twi* double mutants (data not shown).

The dorsal limit of the neuroectoderm, the tracheal pits, could be identified by their *rho* expression (Bier et al. 1990). The relative positions of the neuroectoderm and the tracheal pits as revealed by double staining with the anti-44C antibody for neuroblasts and the *rho* cDNA probe for the tracheal pits were not altered in the mutants (Fig. 5A–D). Thus, the dorsal limit of the neuroectoderm was also shifted ventrally. In other words, the mutant phenotype was characterized mainly by a ventral shift of the neuroectoderm because of changes of cell fate. How the dorsal-most region of the embryo is altered remains to be determined.

Summary of cell fate transformation in *sna* and *twi* mutant embryos

The observed changes of cell fate are summarized in Figure 6 and schematically illustrated in Figure 7. In either single or double mutants of *sna* and *twi*, the ventral-

most cells fail to form mesoderm. In *sna* mutants, cells in the presumptive central mesoderm domain take on the fate of mesectodermal cells, whereas cells in the presumptive peripheral mesoderm domain take on the fate of neuroectoderm. In *twi* mutants, cells in the presumptive peripheral mesoderm domain again become neuroectodermal cells, whereas cells in the presumptive central mesoderm domain are divided into three types: Some have become mesectodermal cells, some have become neuroblasts (Fig. 8D), and the others are neither. In *sna twi* double mutants, neither mesoderm nor mesectoderm is formed; rather, the neuroectoderm has moved into the ventral most region.

Neuroectoderm positioning is relatively independent of the nuclear concentration of *dl* protein

The ventral shift of neuroectoderm and mesectoderm in *sna* and *twi* mutants could either result directly from the impairment of zygotic gene activities of *sna* and *twi* or be a secondary effect because of the effects of *sna* and *twi* mutations on *dl* concentration. Because *dl* nuclear localization is controlled by a group of maternal-effect genes prior to the expression of *sna* and *twi*, the establishment of a *dl* gradient does not require *sna* or *twi*. To examine the possible roles of *sna* and *twi* in the maintenance of *dl* gradient, we looked at the distribution of *dl* in *sna* and *twi* mutants. In wild-type embryos, *dl* is localized to the nuclei of the mesodermal and mesectodermal cells but is not detectable in the nuclei of the neuroectodermal cells (Roth et al. 1989; Steward 1989). The level of *dl* protein is higher at the cellular blastoderm stage than at the gastrulation stage (Fig. 8A,B). Nuclear *dl* protein was also found in the mutants at the earliest stage (gastrulation) when mutant embryos could be identified (Fig. 8C). Although it was difficult to determine quantitatively whether the level of *dl* was different between the wild type and the mutants, there was *dl* protein in the nuclei

Figure 4. Mesectoderm formation. (A–C) m7 expression in stage 5 wild-type (A) and *sna* (B) embryos and a stage 6 *twi* embryo (C); the weaker staining of the more peripheral cells in *sna* mutants disappears by stage 6 leaving 8–12 m7-expressing cells; the m7 expression pattern in *twi* mutants at stage 5 is similar to that in the stage 6 embryo shown here except that there are no m7-expressing cells in the ventral-most region between the two longitudinal rows. The ventral side of embryo is shown; anterior is to the top. (D–E) *rho* expression in cellular blastoderms of wild-type and *sna twi* double mutant. In the wild type (D), *rho* is expressed in certain ectodermal cells in every other segment as well as the mesectodermal cells in all segments. In the *sna twi* double mutant (E), there is no mesectodermal expression of *rho* even though *rho* is still expressed in the ectodermal cells that extend from one side of the embryo to the other. Anterior is to the left; dorsal is up. (F–I) *rho* expression in germ band-extended embryos. Both the midline cells (mesectodermal derivatives) and the tracheal pits express *rho*.

of cells in the ventral region of the *sna twi* double mutants (Fig. 8C). The fact that these cells took on the neuroectodermal fate indicates that this switch of cell fate did not arise because the nuclear dl concentration in *sna twi* double mutants was reduced to the nondetectable level normally found in neuroectodermal cells. Moreover, in *twi* mutants, the presence of several ventral neuroblasts surrounded by *rho*-expressing mesectodermal cells (Fig. 8D; note also a few T5-staining cells in the ventral midline in Fig. 2C and one neuroblast surrounded by mesectodermal cells in Fig. 5C) presents a pattern that is unlikely to arise if the effect of the *twi* mutation is simply to change the dl gradient or the threshold concentration of nuclear dl required to suppress neuroectoderm formation. Taken together, these observations suggest that loss of zygotic gene activity of *sna* and *twi* alters the fate that a cell adopts at blasto-

derm without significantly changing the dorsoventral positional value of that cell.

Discussion

Our analyses of cell fate specification in mutants lacking the zygotic gene activities of *sna* and *twi* have implications concerning both neuroectoderm and mesectoderm formation. In the case of mesectoderm, no transcriptional regulators have been shown previously to be essential in the early steps of mesectoderm determination (Bier et al. 1990; Nambu et al. 1990; Klambt et al. 1991), although putative transcription factors have been identified for the determination of the other domains around the circumference of the embryo (Doyle et al. 1986, 1989; Boulay et al. 1987; Villares and Cabrera 1987; Thisse et al. 1988; Gonzalez et al. 1989; Jimenez and

Figure 5. Position of the neuroectoderm relative to tracheal pits and the ventral midline cells. (A–D) Double-labeling with an anti-44C antibody specific for neuroblasts and a *rho* DNA probe that labels the tracheal pits as well as midline cells. The three rows of brown dots are neuroblasts; the blue lines at the ventral midline are mesectodermal cells. The blue circles marked by arrows at the lateral sides of the embryos are the tracheal pits. The relative positions of tracheal pits and the lateral row of neuroblasts appear similar in the mutants (B–D) and the wild type (A). Note that there is a neuroblast surrounded by *rho*-expressing cells in the *twi* mutant embryo (C).

Campos-Ortega 1990). The finding of the absence of the earliest signs of mesectodermal formation in *sna twi* double mutants therefore suggests that these two genes are involved in mesectoderm determination.

Specification and positioning of the neuroectoderm

By following the formation of the nervous system in mutants without mesoderm, we reached the following conclusions concerning neural development in *Drosophila* embryos.

We have shown that neither mesodermal nor mesec-

todermal cells are essential for neuroblast formation, because neuroblasts were found in embryos lacking mesoderm or both mesoderm and mesectoderm, as in *sna twi* double mutants. Although neural induction in vertebrates has been studied for decades (Spemann 1938; Hamburger 1988), the limited accessibility of *Drosophila* embryos to surgical manipulation has made it difficult to assess the potential role of the mesoderm in neural development. In an earlier experiment, in which the dorsal–ventral axis was changed by altering the activity of the *dl* morphogen, neural tissue could be detected in certain *dl* mutants in the apparent absence of mesodermal derivatives (Campos-Ortega 1983). At that time, though,

Figure 6. Summary of the estimated numbers of cells in the mesoderm, mesectoderm, and neuroectoderm domains. Mesoderm, which corresponds to the ventral-most 15–17 cells in the wild-type embryo, is absent in single or double mutants of *sna* and *twi* (Simpson 1983; Leptin and Grunewald 1990). Mesectoderm formation takes place at late stage 5 and can be recognized by the expression of m7. The neuroectoderm is identified by the early expression of T5 in proneural clusters. Numbers in the boxes are the number of cells along the circumference of an embryo staining for a certain marker. (+) Staining; (–) no staining; (v) ventral midline. (*) The width of m7-staining stripes changes from about four cells in early stage 5 to one cell in late stage 5; (**) the width of neuroectoderm is ~15 cells at stage 5 and stage 7; T5 is expressed in a ventral proneural cluster of 3–5 cells wide, which is separated from a lateral T5-expressing proneural cluster of ~5 cells that do not express T5. (***) In *twi* mutants, some of the ventral-most cells express m7, whereas others do not (see Fig. 4C).

Gene Expression	Stage	Wildtype	<i>sna</i>	<i>twi</i>	<i>sna twi</i>
m7	5	<div style="display: flex; justify-content: space-around;"><div style="text-align: center;">v 1-4* 15-17 1-4* + - +</div><div style="text-align: center;">v 3-4 10-12 3-4 + + +</div><div style="text-align: center;">v 2-4 6-8 2-4 + - +</div></div>	-		
m7	6	<div style="display: flex; justify-content: space-around;"><div style="text-align: center;">v 1 15-17 1 + - +</div><div style="text-align: center;">v - 10-12 - - + -</div><div style="text-align: center;">v 1-3 6-8 1-3 + (<u>+</u>)* +</div></div>	-		
T5	7	<div style="display: flex; justify-content: space-around;"><div style="text-align: center;">v 15** 15-17 15** + - +</div><div style="text-align: center;">v 15** 10-12 15** + - +</div><div style="text-align: center;">v 15** 8-10 15** + - +</div><div style="text-align: center;">v 15** 15** + +</div></div>	-		

the absence of early markers precluded the possibility of detecting early mesoderm and neuroectoderm, making it difficult to rule out the possibility that there was a small amount of mesoderm or a transient existence of mesoderm at the early stages of embryogenesis or to assess the overall effect of changing the dorsal–ventral axis. In this study, the dorsal–ventral gradient, as defined by the dl morphogen, is probably not altered by mutations of the zygotic genes required for mesoderm formation, and the use of multiple early markers has made it possible to show that the mesoderm and mesectoderm are not required for neural induction. It remains possible, however, that mesodermal or mesectodermal derivatives are involved in later processes of neural development, such as axonal guidance (Klamt et al. 1991). It is worth noting that in the vertebrates, although the mesoderm is shown to be able to induce the formation of neural tissue on transplantation (Spemann 1938; Hamburger 1988), most of the experiments did not resolve the question of the normal role of mesoderm in neural development. Recent experiments in *Xenopus* embryos have shown that some neural-specific markers could be induced even when mesoderm invagination was prevented (Kintner and Melton 1987; Dixon and Kintner 1989; Jones and

Woodland 1989; Ruiz i Altaba 1990; see also Sharpe et al. 1987; Hemmati-Brivanlou and Harland 1989). It has also been observed in some cases that neural tissue could form without detectable mesodermal structures (Green and Smith 1990; Godsave and Slack 1991). In none of these situations, however, could the role of mesoderm in neural induction in vertebrates be excluded.

With regard to the positioning of the neuroectoderm in *Drosophila* embryos, we found that the neuroectoderm is shifted ventrally in *sna*, *twi*, and *sna twi* double mutants, even though these mutants do not show any obvious alterations in the distribution of the morphogen dl or any cell death. Thus, ventral cells that have positional values of mesodermal cells in wild-type embryos can adopt neuroectodermal cell fates in the absence of zygotic gene activity of *twi*, *sna*, or both.

Concerning the size of the neuroectoderm, we found that the number of neuroectodermal cells appeared fairly constant even though these cells were at different locations along the dorsal–ventral axis and in rather different surroundings. Thus, the control of the size of the neuroectoderm appears to be intrinsic to the developmental program of the neuroectodermal cells and rather independent of their ventral neighbors or their dorsal–ventral

Figure 7. Summary of cell fate changes in *sna* and *twi* mutants. The relative positions of mesoderm, mesectoderm, neuroectoderm, and tracheal pits are shown in the diagram. The tracheal anlagen are indicated by broken lines (in small circles) because it is not known whether the tracheal pits are determined in the cellular blastoderm or at later stages. In the fate map the tracheal anlagen is positioned just dorsal to the neuroectoderm (Campos-Ortega and Hartenstein 1985). “Unspecified” in *twi* mutants refers to cells that do not express several transcripts or proteins specific for mesectodermal or neuroectodermal cells.

Figure 8. Positioning of neuroectoderm is not directly correlated with dl nuclear localization. (A) Anti-dl staining in a wild-type cellularizing embryo; (B) anti-dl staining in a stage 6 wild-type gastrulating embryo showing a decreased level of nuclear staining in the invaginating cells; (C) anti-dl staining of ventral nuclei in a stage 6 *sna twi* double mutant embryo that shows no ventral invagination. (A–C) Anterior is to the top; ventral is to the right. (D) The ventral view of a *twi* mutant embryo labeled with the anti-44C antiserum for neuroblasts and the *rho* DNA probe showing that neuroblasts around the ventral midline are flanked on either side by *rho*-expressing mesectodermal cells.

positional values. It remains to be determined whether the dorsal neighbors of neuroectoderm exert any influence in defining the dorsal limit of the neuroectoderm.

A model for cell fate specification in the ventral region of Drosophila embryos

The removal of mesoderm in *sna* or *twi* mutants results in a ventral shift of both neuroectoderm and mesectoderm, whereas in the double mutant no mesectoderm forms and the neuroectoderm is located in the ventral-most region. Moreover, the ventral-most cells take on mesectodermal cell fate in *sna* mutants, but some of them appear unspecified in *twi* mutants. To explain the results obtained by us and others thus far (for summary, see Fig. 7; see also Leptin and Grunewald 1990; Nambu et al. 1990), we propose a testable model: (1) Mesoderm formation requires both *sna* and *twi* function within mesodermal cells, and these two genes act in combination, not strictly in a sequential manner, as shown previously (Simpson 1983; Leptin and Grunewald 1990); (2) mesectodermal formation requires either *twi* expression in the mesectodermal cells or *sna* activity in adjacent cells, but is suppressed by *sna* activity within the cell; and (3) the neuroectodermal cell fate is suppressed by *sna* activity within the cell or by the commitment of a cell to the mesectodermal fate.

This model is consistent with the observed expression patterns of *sna* and *twi* and with the mutant phenotypes: The reported distribution of *twi* is wider than that of *sna*, which is restricted to the mesoderm (Thisse et al. 1988;

Leptin and Grunewald 1990). We confirmed with double labeling that *twi* is expressed in mesectodermal cells (data not shown). Thus, in the wild-type embryo, ventral cells that express both *sna* and *twi* form the mesoderm, whereas their immediate neighbors that express *twi* but not *sna* form the mesectoderm; neither of these two groups of cells form the neuroectoderm. In *sna* mutants, the ventral cells that express *twi* become mesectodermal cells and their neighbors form the neuroectoderm. In *twi* mutants, the ventral cells that express *sna* do not take on the fate of mesoderm, mesectoderm, or neuroectoderm, whereas their immediate neighbors form the mesectoderm. In *sna twi* double mutants, the ventral cells do not express either gene and therefore will not form mesoderm or mesectoderm; they take on the default fate of neuroectoderm.

A number of predictions may be tested to determine the validity of the proposed model: (1) A variety of double-labeling experiments should determine whether the basic assumption of the correlation between the expression patterns of *sna* and *twi* and the various cell fates is correct; (2) ectopic expression of *sna* and *twi* may test further the causal relation between *sna* and *twi* expression and cell fate; and (3) our proposal that mesectodermal formation in *twi* mutants results from *sna* activity in neighboring cells predicts that *sna* distribution is limited to the presumptive central mesoderm in *twi* mutants, which is flanked by mesectodermal cells. We note that these predictions are all qualitative and apply only to the early expression patterns at a stage relevant for these cell fate determination events. It is quite possible

that future studies will reveal deviations from the predictions of this simple model. Experiments such as those outlined here should nevertheless help further our understanding of cell fate specification during early embryogenesis.

Since the submission of this paper, Alberga et al. (1991) have reported the expression pattern of *sna* RNA and protein. The *sna* protein is expressed first in the primordium of the anterior midgut and in the mesoderm; expression of *sna* in mesectodermal cells and neuroectodermal cells appears much later, probably after the stages when they express specific early markers. Moreover, *sna* expression in *twi* mutant embryos at the blastoderm stage is more restricted, in a ventral stripe that is approximately the width of the presumptive central mesoderm. These observations are in accord with our model, although double-labeling experiments are necessary to examine its predictions with better spatial and temporal resolution.

Materials and methods

Drosophila strains

The alleles used, *sna*^{11G05}, *twi*^{S60}, and *sna*^{11G05} *twi*^{S60} recombinant, are the same amorphic alleles that were characterized in detail for their mesoderm formation defects (Leptin and Grunewald 1990). The *sc*^{B57} mutation is a deletion of the entire AS-C (isolated by E. Grell, Gonzalez et al. 1989). *Dl*^{9P39}, *N*^{55e11}, and *E[sp1]*^{8D06} are amorphic alleles of the neurogenic genes (Lehmann et al. 1983; Ziemer et al. 1988). Double mutants of a mesodermal gene and a proneural gene or a neurogenic gene are easily recognizable because the *sna* or *twi* mutation causes characteristic twisting of the embryo, whereas the other mutations cause reduction or overproduction of the nervous tissue. Staging of mutant embryos was done with the same criteria (except that mutants do not form the ventral furrow) as for the wild-type embryos (Campos-Ortega and Hartenstein 1985); stage 6 and 7 embryos are staged according to the appearance of pole cells and the cephalic furrow, and stage 8 and 9 embryos are staged according to the extent of the germ-band extension.

Immunocytochemistry

Embryos of appropriate stages were collected on grape plates. They were dechorionated with 50% bleach for several minutes and rinsed in water before fixation for 20 min in 4% formaldehyde in PBS (covered with heptane) and removal of the vitelline membrane in methanol. Embryos can be stored in ethanol at -20°C for months before antibody staining (Bodmer et al. 1987) or in situ hybridization.

To obtain a quantitative measure of the number of neuroblasts per segment, we first made sure that the number of transverse rows of neuroblasts was the same in the cephalic and thoracic region of the wild-type and mutant embryos and then measured the total number of neuroblasts that expressed the gene at 44°C so as to obtain an average number per segment (four transverse rows; see Campos-Ortega and Hartenstein 1985). In *twi* mutants, the isolated neuroblasts in the presumptive central mesoderm region were not included in the neuroblast count. In the *sna twi* double mutants, however, the neuroblasts in the ventral midline were included in the cell count.

In situ hybridization

The in situ hybridization protocol that we used was modified from Tautz and Pfeifle (1989). Probes were prepared with a Boehringer Mannheim kit according to manufacturer's protocol with the modifications that 5 μg of random primers was added in the labeling reaction and that reaction was allowed to go on overnight. Products from one labeling reaction with <1 μg of template DNA were stored in 30 μl of water at -20°C , from which 3 μl was used for each hybridization.

Double staining with an antibody and a DNA probe

The embryos were first stained with an antibody and then processed for in situ hybridization. Immunocytochemistry was performed according to the protocol described previously (Bodmer et al. 1987) to the step after diaminobenzidine (DAB) reaction. Stained embryos dehydrated in ethanol were then used for in situ hybridization in the same way as described above. Color development was monitored closely to maximize the detection of both signals.

Acknowledgments

We thank S. Barbel for excellent technical assistance; K. Anderson for *sna* and *twi* mutants; E. Grell for *sc*^{B57}; E. Bier for the *rho* probe; J. Modolell for AS-C probes; E. Knust for the m7 probe; R. Steward, D. Stein, and C. Nüsslein-Volhard for anti-*dl* antisera; L. Ackerman for help with photography; R. Carretto for help with embryo collections; K. Prewitt for manuscript preparation; J.Y. Wu for helpful discussions; and V. Siegel for helpful comments on the manuscript. Y.R. was supported by the Neuroscience Program at the University of California at San Francisco. L.Y.J. and Y.N.J. are Howard Hughes Medical Institute investigators.

The publication costs of this article were defrayed in part by payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 USC section 1734 solely to indicate this fact.

References

- Alberga, A., J.L. Boulay, E. Kempe, C. Dennefeld, and M. Haenlin. 1991. The *snail* gene required for mesoderm formation in *Drosophila* is expressed dynamically in derivatives of all three germ layers. *Development* **111**: 983–992.
- Anderson, K. 1987. Dorsal-ventral embryonic pattern genes of *Drosophila*. *Trends Genet.* **3**: 91–97.
- Artavanis-Tsakonas, S. 1988. The molecular biology of the *Notch* locus and the fine tuning of differentiation in *Drosophila*. *Trends Genet.* **4**: 95–100.
- Bier, E., L. Ackerman, S. Barbel, L. Jan, and Y.N. Jan. 1988. Identification and characterization of a neuron-specific nuclear antigen in *Drosophila*. *Science* **240**: 913–916.
- Bier, E., L.Y. Jan, and Y.N. Jan. 1990. *rhomboid*, a gene required for dorso-ventral axis establishment and peripheral nervous system development in *Drosophila melanogaster*. *Genes & Dev.* **4**: 190–203.
- Bodmer, R., S. Barbel, S. Shepherd, J.W. Jack, L.Y. Jan, and Y.N. Jan. 1987. Transformation of sensory organs by mutations of the *cut* locus of *D. melanogaster*. *Cell* **51**: 293–307.
- Boulay, J.L., C. Dennefeld, and A. Alberga. 1987. The *Drosophila* developmental gene *snail* encodes a protein with nucleic acid binding fingers. *Nature* **330**: 395–398.
- Boulianne, G.L., A. de la Concha, J.A. Campos-Ortega, L.Y. Jan, and Y.N. Jan. 1991. The *Drosophila* neurogenic gene *neural-*

- ized encodes a novel protein which is expressed in precursors of larval and adult neurons. *EMBO J.* (in press).
- Cabrera, C.V., A. Martinez-Arias, and M. Bate. 1987. The expression of three members of the *achaete-scute* gene complex correlates with neuroblast segregation in *Drosophila*. *Cell* **50**: 425–433.
- Campos-Ortega, J.A. 1983. Topological specificity of phenotype expression of neurogenic mutations in *Drosophila*. *Wilhelm Roux's Arch. Dev. Biol.* **192**: 317–326.
- Campos-Ortega, J.A. and V. Hartenstein. 1985. *The embryonic development of Drosophila melanogaster*. Springer Verlag, Heidelberg.
- Campos-Ortega, J.A. and Y.N. Jan. 1991. Genetic and molecular bases of neurogenesis in *Drosophila melanogaster*. *Annu. Rev. Neurosci.* **14**: 399–420.
- Campuzano, S., L. Carramolina, C.V. Cabrera, M. Ruiz-Gomez, R. Villares, A. Boronat, and J. Modolell. 1985. Molecular genetics of the *achaete-scute* gene complex of *Drosophila melanogaster*. *Cell* **40**: 327–338.
- Caudy, M., H. Vaessin, M. Brand, R. Tuma, L.Y. Jan, and Y.N. Jan. 1988. *daughterless*, a *Drosophila* gene essential for both neurogenesis and sex determination, has sequence similarities to *myc* and the *achaete-scute* complex. *Cell* **55**: 1061–1067.
- Dambly-Chaudiere, C. and A. Ghysen. 1987. Independent sub-patterns of sense organs require independent genes of the *achaete-scute* complex in *Drosophila* larvae. *Genes & Dev.* **1**: 297–306.
- Dixon, J.E. and C.R. Kintner. 1989. Cellular contacts required for neural induction in *Xenopus* embryos: Evidence for two signals. *Development* **106**: 749–757.
- Doe, C.Q. and C.S. Goodman. 1985. Early events in insect neurogenesis. II. The role of cell interactions and cell lineages in the determination of neuronal precursor cells. *Dev. Biol.* **111**: 206–219.
- Doyle, H.J., K. Harding, T. Hoey, and M. Levine. 1986. Transcripts encoded by a homeobox gene are restricted to dorsal tissues of *Drosophila* embryos. *Nature* **323**: 76–79.
- Doyle, H.J., R. Kraut, and M. Levine. 1989. Spatial regulation of *zerknüllt*: A dorsal–ventral patterning gene in *Drosophila*. *Genes & Dev.* **3**: 1515–1533.
- Ghysen, A. and C. Dambly-Chaudiere. 1988. From DNA to form: The *achaete-scute* complex. *Genes & Dev.* **2**: 495–501.
- Godsave, S.F. and J.M.W. Slack. 1991. Single cell analysis of mesoderm formation in the *Xenopus* embryo. *Development* **111**: 523–530.
- Gonzalez, F., S. Romani, P. Cubas, J. Modolell, and S. Campuzano. 1989. Molecular analysis of the *asense* gene, a member of the *achaete-scute* complex of *Drosophila melanogaster* and its novel role in optic lobe development. *EMBO J.* **8**: 3553–3562.
- Green, J.B.A. and J.C. Smith. 1990. Graded changes in dose of a *Xenopus* activin A homologue elicit stepwise transitions in embryonic cell fate. *Nature* **347**: 391–394.
- Gurdon, J.B. 1987. Embryonic induction-molecular prospects. *Development* **99**: 285–306.
- Haenlin, M., B. Kramatschek, and J.A. Campos-Ortega. 1990. The pattern of transcription of the neurogenic gene *Delta* of *Drosophila melanogaster*. *Development* **110**: 905–914.
- Hamburger, V. 1988. *The heritage of experimental embryology: Hans Spemann and the organizer*. Oxford University Press, Cambridge.
- Heitzler, P. and P. Simpson. 1991. The choice of cell fate in the epidermis of *Drosophila*. *Cell* **64**: 1083–1092.
- Hemmati-Brivanlou, A.H. and R.M. Harland. 1989. Expression of an engrailed-related protein is induced in the anterior neural ectoderm of early *Xenopus* embryos. *Development* **106**: 611–617.
- Jan, L.Y. and Y.N. Jan. 1982. Antibodies to horseradish peroxidase as specific neuronal markers in *Drosophila* and in grasshopper embryos. *Proc. Natl. Acad. Sci.* **72**: 2700–2704.
- Jimenez, J. and J. Campos-Ortega. 1990. Defective neuroblast commitment in mutants of the *achaete-scute* complex and adjacent genes of *Drosophila melanogaster*. *Neuron* **5**: 81–89.
- Jones, E.A. and H.R. Woodland. 1989. Spatial aspects of neural induction in *Xenopus laevis*. *Development* **107**: 785–792.
- Kidd, S., M.R. Kelley, and M.W. Young. 1986. Sequence of the *Notch* locus of *Drosophila melanogaster*: Relationship of the encoded protein to mammalian clotting and growth factors. *Mol. Cell. Biol.* **6**: 3094–3108.
- Kintner, C.R. and D.A. Melton. 1987. Expression of *Xenopus* N-CAM in ectoderm is an early response to neural induction. *Development* **99**: 311–325.
- Klambt, C., J.R. Jacobs, and C.S. Goodman. 1991. The midline of the *Drosophila* central nervous system: A model for the genetic analysis of cell fate, cell migration and growth cone guidance. *Cell* **64**: 801–815.
- Knust, E., K. Tietze, and J.A. Campos-Ortega. 1987. Molecular analysis of the neurogenic locus *Enhancer of split* of *Drosophila melanogaster*. *EMBO J.* **6**: 4113–4123.
- Kopczynski, C.C., A.K. Alton, K. Fechtel, P.J. Kooh, and M.A.T. Muskavitch. 1988. *Delta*, a *Drosophila* neurogenic gene, is transcriptionally complex and encodes a protein related to blood coagulation factors and epidermal growth factor of vertebrates. *Genes & Dev.* **2**: 1723–1735.
- Lehmann, R., U. Dietrich, F. Jimenez, and J.A. Campos-Ortega. 1983. On the phenotype and development of mutants of early neurogenesis in *Drosophila melanogaster*. *Wilhelm Roux's Arch. Dev. Biol.* **192**: 62–74.
- Leptin, M. and B. Grunewald. 1990. Cell shape changes during gastrulation in *Drosophila*. *Development* **110**: 73–84.
- Nambu, J.R., R.G. Franks, S. Hu, and S.T. Crews. 1990. The *single-minded* gene of *Drosophila* is required for the expression of genes important for the development of CNS midline cells. *Cell* **63**: 63–75.
- Poulson, D.F. 1950. Histogenesis, organogenesis and differentiation in the embryo of *Drosophila melanogaster*. In *Biology of Drosophila* (ed. M. Demerec), pp. 168–274. Wiley, New York.
- Preiss, A., D.A. Hartley, and S. Artavanis-Tsakonas. 1988. The molecular genetics of *Enhancer of split*, a gene required for embryonic neural development in *Drosophila*. *EMBO J.* **7**: 3917–3928.
- Rao, Y., L. Jan, and Y.N. Jan. 1990. Similarity of the product of the *Drosophila* neurogenic gene *big brain* to transmembrane channel proteins. *Nature* **345**: 163–167.
- Romani, S., S. Campuzano, and J. Modolell. 1987. The *achaete-scute* complex is expressed in the neurogenic regions of *Drosophila* embryos. *EMBO J.* **6**: 2085–2092.
- Roth, S., D. Stein, and C. Nüsslein-Volhard. 1989. A gradient of nuclear localization of the *dorsal* protein determines dorsoventral pattern in *Drosophila* embryo. *Cell* **59**: 1189–1202.
- Ruiz i Altaba, A. 1990. Neural expression of the *Xenopus* homeobox gene *Xhox3*: Evidence for a patterning neural signal that spreads through the ectoderm. *Development* **108**: 595–604.
- Rushlow, C.A., K. Han, J.L. Manley, and M. Levine. 1987. The graded distribution of the *dorsal* morphogen is initiated by selective nuclear transport in *Drosophila*. *Cell* **59**: 1165–1177.

- Simpson, P. 1983. Maternal-zygotic gene interactions during formation of the dorsoventral pattern in *Drosophila* embryos. *Genetics* **105**: 615–632.
- Sharpe, C.R. and J.B. Gurdon. 1990. The induction of anterior and posterior neural genes in *X. laevis*. *Development* **109**: 765–774.
- Sharpe, C.R., A. Fritz, E.M. De Robertis, and J.B. Gurdon. 1987. A homeobox containing marker of posterior neural differentiation shows the importance of predetermination in neural induction. *Cell* **50**: 749–758.
- Spemann, H. 1938. *Embryonic development and induction*. Yale University Press, New Haven.
- Spemann, H. and H. Mangold. 1924. Induction of embryonic primordia by implantation of organizers from a different species. *Arch. Mikrosk. Anat. Entwicklungsmech* **100**: 599–638. Translated by V. Hamburger. 1964. *Foundations of experimental embryology* (ed. B.H. Willier and J.M. Oppenheimer). Prentice-Hall, Englewood Cliffs, New Jersey.
- Steward, R. 1989. Relocalization of the dorsal protein from cytoplasm to the nucleus correlates with its function. *Cell* **59**: 1179–1188.
- Tautz, D. and D. Pfeifle. 1989. A nonradioactive *in situ* hybridization method for the localization of specific RNAs in *Drosophila* embryos reveals a translational control of the segmentation gene *hunchback*. *Chromosoma* **98**: 81–85.
- Thisse, B., C. Stoetzel, T.C. Gorostiza, and F. Perin-Schmitt. 1988. Sequence of the *twist* gene and nuclear localization of its protein in endomesodermal cells of early *Drosophila* embryos. *EMBO J.* **7**: 2175–2183.
- Vaessin, H., K.A. Bremer, E. Knust, and J.A. Campos-Ortega. 1987. The neurogenic locus *Delta* of *Drosophila melanogaster* is expressed in neurogenic territories and encodes a putative transmembrane protein with EGF-like repeats. *EMBO J.* **6**: 3431–3440.
- Villares, R. and C.V. Cabrera. 1987. The *achaete-scute* gene complex of *D. melanogaster*: Conserved domains in a subset of genes required for neurogenesis and their homology to *myc*. *Cell* **50**: 415–424.
- Wharton, K.A., K.M. Hohansen, T. Xu, and S. Artavanis-Tsakonas. 1985. Nucleotide sequence from the neurogenic locus *Notch* implies a gene product that shares homology with proteins containing EGF-like repeats. *Cell* **43**: 567–581.
- Yamada, T., M. Placzek, H. Tanaka, J. Dodd, and T.M. Jessell. 1991. Control of cell pattern in the developing nervous system: Polarizing activity of the floor plate and notochord. *Cell* **64**: 635–647.
- Yedbovnick, B., D. Smoller, P. Young, and D. Mills. 1988. Molecular analysis of the neurogenic locus *mastermind* of *Drosophila melanogaster*. *Genetics* **118**: 483–497.
- Ziemer, A., K. Tietze, E. Knust, and J.A. Campos-Ortega. 1988. Genetic analysis of *Enhancer of split*, a locus involved in neurogenesis in *Drosophila melanogaster*. *Genetics* **119**: 63–74.

Neuroectoderm in *Drosophila* embryos is dependent on the mesoderm for positioning but not for formation.

Y Rao, H Vaessin, L Y Jan, et al.

Genes Dev. 1991, **5**:

Access the most recent version at doi:[10.1101/gad.5.9.1577](https://doi.org/10.1101/gad.5.9.1577)

References

This article cites 54 articles, 19 of which can be accessed free at:
<http://genesdev.cshlp.org/content/5/9/1577.full.html#ref-list-1>

License

Email Alerting Service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).