The protein kinase C-activated MAP kinase pathway of *Saccharomyces cerevisiae* mediates a novel aspect of the heat shock response

Yoshiaki Kamada, Un Sung Jung, Julia Piotrowski, and David E. Levin¹

Department of Biochemistry, The Johns Hopkins University School of Public Health, Baltimore, Maryland 21205 USA

The *PKC1* gene of budding yeast encodes a homolog of the α , β , and γ isoforms of mammalian PKC that is proposed to regulate a MAPK-activation pathway. Mutants in this pathway undergo cell lysis resulting from a deficiency in cell wall construction when they attempt to grow at elevated temperatures. We show that the *PKC1*-regulated pathway is important for induced thermotolerance and that the MPK1 protein kinase (the MAPK of this pathway) is strongly activated by mild heat shock. This activation is sustained during growth at high temperature and is dependent on the function of pathway components proposed to function upstream of *MPK1*, including *PKC1*. Expression of genes under the control of known heat shock-inducible promoter elements (HSEs and STREs) was not compromised in *PKC1* pathway mutants, indicating that this pathway mediates a novel aspect of the yeast heat shock response. We propose that the heat-induced signal for pathway activation is generated in response to weakness in the cell wall created during growth under thermal stress, perhaps as a result of increased membrane fluidity. Evidence is presented that the mechanism by which the cell detects this weakness is by measuring stretch of the plasma membrane.

[Key Words: PKC; MAP kinase; heat shock; yeast; cell integrity]

Received March 14, 1995; revised version accepted May 23, 1995.

Members of the family of phospholipid-dependent, serine/threonine-specific protein kinases, known collectively as protein kinase C (PKC), respond to extracellular signals that act through receptor-mediated hydrolysis of phosphatidylinositol-4,5-bisphosphate to diacylglycerol (DAG) and inositol-1,4,5-trisphosphate (IP_3) (Hokin 1985). DAG serves as a second messenger to activate PKC (Takai et al. 1979; Kishimoto et al. 1980; Nishizuka 1986, 1988), and IP_3 functions to mobilize Ca^{2+} from intracellular stores (Berridge and Irvine 1984). A dozen distinct subtypes of mammalian PKC have been reported to date (Dekker and Parker 1994). The four initially identified isozymes, α , βI , βII , and γ , are structurally closely related and display similar catalytic properties. The activity of these enzymes can be stimulated by Ca^{2+} , but at physiological Ca^{2+} concentrations DAG is also required for activation. Five additional isozymes, δ , ϵ , θ , $\eta(L)$, and μ , are related more distantly to the α , β , and γ subtypes and lack a conserved region thought to be important for Ca^{2+} binding. Consistent with this structural feature, these subtypes display Ca^{2+} -independent protein kinase activity (Nishizuka 1992). Three recently identified subtypes, ζ , λ , and ι also lack the putative

Ca^{2+} -binding domain, but are unusual in that they lack one of the pair of DAG-binding zinc fingers.

Mammalian PKC is thought to play a pivotal role in the regulation of a host of cellular functions through its activation by growth factors and other agonists. These functions include cell growth, proliferation, and differentiation (Rosengurt et al. 1984; Kaibuchi et al. 1985; Persons et al. 1988; Michak et al. 1993), release of various hormones (Negro-Vilar and Lapetina 1985; Ohmura and Friesen 1985), and control of ion conductance channels (Farley and Auerbach 1986; Madison et al. 1986; Hardy et al. 1995). PKC induces the transcription of a wide array of genes, including the proto-oncogenes *c-myc*, *c-fos*, *c-sis*, and *c-jun* (Kelly et al. 1983; Greenberg and Ziff 1984; Kruger et al. 1984; Colamonici et al. 1986; Angel et al. 1988), human collagenase (Angel et al. 1987), metallothionein II_A, and the SV40 early genes (Imbra and Karin 1986). Several transcription factors have been implicated in this response, including components of the AP-1 complex, AP-2, AP-3, and NF- κ B (Chiu et al. 1987; Imagawa et al. 1987; Lee et al. 1987; Baeurle and Baltimore 1988).

Although substantial progress has been made toward elucidating the pathways leading to PKC activation, the steps between this activation and subsequent nuclear events are not well understood. However, members of a

¹Corresponding author.

family of enzymes called mitogen-activated protein kinases (MAPKs) have been implicated in PKC-dependent signaling. These enzymes are thought to function as intermediaries between membrane-associated signaling molecules and the nucleus (Pelech and Sanghera 1992; Thomas 1992). The observation that at least some members of the MAPK family translocate between the cytoplasm and the nucleus (Chen et al. 1992; Sanghera et al. 1992; Lenormand et al. 1993) makes them excellent candidates for messengers from the membrane. In support of this notion is the finding that a number of nuclear transcription factors are targets for MAPKs (Hill and Treisman 1995). MAPKs are activated through protein kinase cascades that are comprised of three highly conserved core components. The MAPKs themselves are activated in response to dual phosphorylation on a threonine and tyrosine residue by MAPK kinases (or MEKs) (Ahn et al. 1992). MEKs are phosphorylated and activated by members of a third family of protein kinases called MEK kinases (MEKKs; Herskowitz 1995).

The initially described p42/p44 isoforms of MAPK (ERKs) are activated in response to a wide array of extracellular agonists, including those that stimulate PKC, tyrosine kinases, and G proteins (Cobb et al. 1991). Additional subtypes of mammalian MAPKs reported recently include the p46 and p54 c-Jun amino-terminal kinases (JNKs) (Hibi et al. 1993; Derijard et al. 1994; Galcheva-Gargova et al. 1994; Kyriakis et al. 1994) and p38 MAPKs (Han et al. 1994; Rouse et al. 1994). Apparently distinct in function from the p42/p44 isoforms, these enzymes are activated in response to cytokines and a variety of stresses including heat shock, high osmolarity, hydrogen peroxide, UV light, sodium arsenite, and inhibition of protein glycosylation or synthesis. The mechanisms by which such varied stress signals activate these novel MAPK isotypes are not yet clear. However, JNK activation requires costimulation of both Ca^{2+} - and PKC-dependent pathways in T cells (Su et al. 1994).

We are studying signal transduction from PKC in the budding yeast *Saccharomyces cerevisiae*. The *PKC1* gene of *S. cerevisiae* encodes a homolog of the Ca^{2+} -stimulated subtypes of mammalian PKC that is essential for cell growth (Levin et al. 1990). Loss of *PKC1* function results in a cell lysis defect that is attributable to a deficiency in cell wall construction (Levin and Bartlett-Heubusch 1992; Paravicini et al. 1992; Levin et al. 1994). Several additional components of the signaling pathway regulated by *PKC1* were identified by isolating genes whose mutational activation, or expression at high levels, could suppress the cell lysis defect resulting from pathway inactivation. Among these are four genes that encode protein kinases proposed to constitute a MAPK activation cascade that is under the control of *PKC1*. These include a MEKK homolog (*BCK1*; Lee and Levin 1992), a redundant pair of MEK homologs (*MKK1*/*MKK2*; Irie et al. 1993), and a MAPK homolog (*MPK1*; Lee et al. 1993; initially designated *SLT2*; Torres et al. 1991). Deletion of any of these components results in cell lysis only when cells are cultivated at 37°C. Because loss of *PKC1* function results in a more severe defect

(lysis at any temperature) than does loss of function of the downstream components, we have proposed that *PKC1* regulates a bifurcated pathway, on one branch of which functions this MAPK activation cascade (Lee and Levin 1992). Genetic interactions among these protein kinase-encoding genes suggest that they function in a linear pathway from *PKC1* to *BCK1* to *MKK1*/*MKK2* to *MPK1* (Irie et al. 1993; Lee et al. 1993). This pathway is one of five physiologically distinct MAPK activation cascades known to exist in budding yeast (Herskowitz 1995; Levin and Errede 1995).

To identify the signals that activate the *PKC1*-regulated cell integrity pathway, we have developed an immune-complex protein kinase assay for the *MPK1*-encoded MAPK. We show here that this signaling pathway is strongly activated by growth under mild heat shock conditions. We also present evidence that heat-induced activation of this pathway is in response to the development of weakness in the cell wall that is detected by stretch of the plasma membrane.

Results

The PKC1-regulated signaling pathway is important for induced thermotolerance

Wild-type *S. cerevisiae* cells exhibit the ability to survive conditions of severe heat shock if they are first exposed to a mild heat shock (Sanchez et al. 1992). This behavior, known as induced thermotolerance, is attributable in part to the induced expression of a set of so-called heat shock genes (Parsell and Lindquist 1994). Because deletion mutants in the *PKC1*-regulated MAPK-activation pathway (*bck1Δ*, *mkk1/mkk2Δ*, and *mpk1Δ*) grow normally at 23°C, but undergo cell lysis when cultivated at 37°C (Torres et al. 1991; Lee and Levin 1992; Irie et al. 1993; Lee et al. 1993), we explored the possibility that this pathway plays a role in induced thermotolerance. Figure 1A shows that a *bck1Δ* mutant was no more sensitive than an isogenic *BCK1*⁺ strain to killing by shift from logarithmic growth at 23–50°C. However, the *bck1Δ* mutant was partially deficient in the acquisition of thermotolerance induced by brief pretreatment at 37°C (30 min; Fig. 1B). Similar results were obtained using an *mpk1Δ* mutant (data not shown). In addition, a constitutive allele of *BCK1* (*BCK1-20*), which alleviates the requirement for *PKC1* (Lee and Levin 1992), conferred partial resistance to heat shock (Fig. 1A). These results indicate that the *PKC1*-regulated pathway plays a role in induced thermotolerance and suggest that this pathway might be activated in response to heat shock.

The MPK1-encoded MAPK is activated by heat shock in a PKC1-dependent manner

To test directly the effect of growth temperature on *PKC1*-regulated pathway activity, we developed an immune complex protein kinase assay for the *MPK1*-encoded MAPK. The *MPK1* gene was fused to a sequence encoding an epitope from the influenza virus hemagglu-

Figure 1. *BCK1* is important for induced thermotolerance. Cells growing in YEPD at 23°C were either shifted directly to 50°C for the indicated times (A) or pretreated at 37°C for 30 min before the challenge temperature (B). After heat shock, cells were diluted and plated for viability. For B, percent survival is relative to value after pretreatment. Strains were wild-type (1788), *BCK1-20* bearing YEp352[*MPK1::HA*] (DL1183), and *bck1Δ* (DL251). Overproduction of *MPK1* from YEp352[*MPK1::HA*] enhanced thermal resistance conferred by *BCK1-20*. Each value is the mean and standard deviation of at least three experiments.

tinin (HA) protein (Wilson et al. 1984) to allow detection and immunoprecipitation of MPK1. The resulting fusion product, designated MPK1^{HA}, differed from MPK1 by the addition of 40 amino acids to its carboxyl terminus. MPK1^{HA} expressed in yeast cells appeared as a 68-kD band by SDS-PAGE (Fig. 2A), somewhat larger than predicted by its molecular mass (~60 kD). An *mpk1Δ* mutant was complemented by low-level expression of MPK1^{HA} (not shown), indicating that this construction is biologically functional.

MPK1^{HA} was immunoprecipitated from extracts of cells growing at various temperatures. The protein kinase activity associated with MPK1^{HA} was measured in immune complex using bovine myelin basic protein (MBP) as substrate, an excellent substrate for mammalian MAPKs. MPK1^{HA} from cells growing at 23°C was very weakly active; enzyme from cells at 30°C was moderately active; and enzyme from cells at 37°C was strongly active (Fig. 2B). Because extracts were made after long-term cultivation of cells in log phase (12–15 hr), we conclude that MPK1^{HA} activity remains persistently high in cells growing at elevated temperature. The range of activity at different temperatures reflects differences in MPK1^{HA} specific activity because the amount of this protein in the immunoprecipitates did not vary appreciably in cells grown at different temperatures (Fig. 2B, bottom). In addition, the observed differences in protein kinase activity are not a function of growth rate because the optimal growth temperature for this species is 30°C. This protein kinase activity is an intrinsic property of the MPK1^{HA} protein and is not attributable to a contaminating protein kinase, because a biologically inactive mutant form of this protein was biochemically inactive (see below).

To examine the kinetics of MPK1^{HA} activation in response to temperature change, growing cells were shifted rapidly from 23°C to 39°C by dilution into prewarmed medium. Figure 2C shows that activity remained low until ~20 min after temperature shift and peaked by 30 min. Later time points showed no further increase in activity (not shown). This activation represents a 170-fold increase in MPK1^{HA} specific activity.

The delay in MPK1^{HA} activation in response to elevated temperature is considerably longer than that observed for activation of two other yeast MAPKs in response to extracellular signals. The FUS3 and HOG1 protein kinases become fully active within 1 or 2 min after

Figure 2. Activation of MPK1 by growth at elevated temperatures. (A) Immunoblot detection of MPK1^{HA}. Immunoprecipitations using the anti-HA antibody (12CA5) were done with cell extracts (50 μg protein) prepared from a wild-type strain expressing either untagged MPK1 (DL744), or MPK1^{HA} (DL922) in the presence (lanes 2,4) or absence (lanes 1,3) of antibody. Immune complexes were subjected to SDS-PAGE on a 7.5% acrylamide gel. Immunodetection was performed as described in Materials and methods. The molecular masses of the standards are indicated (in kD). The asterisk indicates coimmunoprecipitated 12CA5 IgG. (B) MPK1 activity is dependent on growth temperature. MPK1^{HA} was immunoprecipitated from extracts of cultures (DL1101; expresses MPK1^{HA}) growing at 23°C (lanes 1,2), 30°C (lanes 3,4), or 37°C (lanes 5,6). In vitro protein kinase assays were conducted using MBP as a substrate, and the reactions were subjected to SDS-PAGE on a 12.5% acrylamide gel, as described in Materials and methods (top). Autophosphorylated MPK1^{HA} and phosphorylated 12CA5 IgG (asterisk) are also indicated. All other bands derive from the MBP preparation. A parallel set of immune complexes were subjected to immunodetection of MPK1^{HA} (bottom). (C) MPK1 is activated by heat shock. MPK1^{HA} was immunoprecipitated from extracts of DL1101 cultures shifted from 23°C to 39°C for the indicated times. Protein kinase assays (top) and immunodetection of MPK1^{HA} (bottom) were carried out on parallel sets of immune complexes.

exposure of cells to their respective activating signals (Gartner et al. 1992; Brewster et al. 1993). This suggests that elevated growth temperature does not induce a direct signal for MPK1^{HA} activation but that this activation is the result of a physiological change arising from growth under the new environmental condition. To determine whether MPK1^{HA} activation depends on the synthesis of a new protein, cells growing at 23°C were pretreated with the protein synthesis inhibitor cycloheximide before a 30-min shift to 39°C. Initial MPK1^{HA} activation was not impaired by the absence of protein synthesis (Fig. 3A, cf. lanes 2 and 5), indicating that the delay in thermally induced activation is not attributable to a requirement for the synthesis of new proteins. Cycloheximide treatment alone had a slight activating effect on MPK1^{HA} (threefold; Fig. 3A, cf. lanes 1 and 4).

To confirm that the protein kinase activity detected was an intrinsic property of MPK1^{HA}, rather than a contaminating activity in the immune precipitates, a mutant form of this protein that lacks the two phosphorylatable residues predicted to be required for activation was tested for protein kinase activity. Mutation of either (or both) Thr-190 to Ala, or Tyr-192 to Phe, severely diminishes the biological activity of MPK1 [Lee et al. 1993]. An MPK1⁺ strain expressing the *mpk1-T190A*, *Y192F::HA* double mutant allele was shifted from

growth at 23°C to 39°C for 30 min. Immunoprecipitated MPK1-TAYF^{HA} displayed no detectable protein kinase activity (Fig. 3A, lane 7).

To test the veracity of the model proposed for the architecture of the *PKC1*-regulated pathway (Lee et al. 1993), we examined the requirement for *BCK1* function in MPK1^{HA} activation. MPK1^{HA} isolated from a temperature-shifted strain bearing a *bck1Δ* mutation possessed no detectable activity (Fig. 3A, lane 9), confirming the dependence on BCK1 for MPK1^{HA} activity. As an additional test of the proposed model, we examined the requirement for *PKC1* function in the activation of MPK1^{HA}. The MPK1^{HA} protein was expressed in a strain bearing a temperature-sensitive allele of *PKC1* (*pkc1-2*; Levin and Bartlett-Heubusch 1992). The mutant strain was subjected to a 30-min shift from 25°C to 37°C, a treatment that had the dual effect of creating the appropriate conditions for MPK1 activation while inactivating the thermolabile PKC1. In addition, this treatment does not cause appreciable killing of the mutant over the time course of the experiment (Levin and Bartlett-Heubusch 1992). Figure 3B shows that this temperature shift failed to activate MPK1^{HA} in the *pkc1* mutant, confirming the requirement for PKC1 in signal transmission to this MAPK.

Heat shock gene expression is not regulated by the PKC1 pathway

Mild heat shock induces the expression of an array of genes under the control of the heat shock transcription factor (HSF; Sorger and Pelham 1988; Weiderrecht et al. 1988). Many of these genes encode molecular chaperones that aid in the refolding or proteolytic degradation of unfolded proteins under thermal stress conditions (Morimoto et al. 1994). HSF binds to a promoter element in these genes known as the heat shock element (HSE; Munro and Pelham 1985). To determine whether heat shock-induced gene expression is dependent on the *PKC1*-regulated pathway, we measured the expression of a reporter gene (*Escherichia coli lacZ*) driven by either a synthetic- or natural HSE-associated promoter in an *mpk1Δ* mutant and a *pkc1Δ* mutant. Table 1 shows that heat shock-induced expression from these promoters was normal in both mutants, although the *pkc1Δ* mutant displayed an elevated constitutive level of expression. This indicates that the *PKC1*-regulated pathway is not responsible for transcriptional activation of genes under HSE control. But because the expression of some heat shock genes is also regulated post-transcriptionally (e.g., *HSP104*; S. Lindquist, pers. comm.), we examined the heat shock-induced accumulation of the HSP104 protein in a *bck1Δ* mutant. Here again, no difference was observed between the mutant and wild-type strain (not shown).

A second promoter element that is activated in response to heat shock is the stress-responsive element (STRE; Marchler et al. 1993). To determine whether STRE-driven gene expression is dependent on the *PKC1*-mediated pathway, we measured expression of the *CTT1*

Figure 3. Activation of MPK1 does not require protein synthesis, but it does require signals from PKC1 and BCK1. (A) Stimulation of MPK1 protein kinase activity in response to shift from growth at 23°C–39°C for 30 min was detected as described in Fig. 2 (top). Strains were wild-type (DL922; expresses MPK1^{HA}), *mpk1-TAYF::HA* (DL1182; expresses MPK1-T190A, Y192F^{HA}), and *bck1Δ::URA3* (DL923; expresses MPK1^{HA}). (Lanes 4,5) Cycloheximide (100 μg/ml) was added to the culture 30 min before heat shock. A parallel set of immune complexes were subjected to immunodetection of MPK1^{HA} (bottom). (B) Stimulation of MPK1 protein kinase activity in response to shift from growth at 25°C to 37°C for 30 min in a temperature-sensitive *pkc1* mutant. Strains were wild type (DL1101; expresses MPK1^{HA}) and *pkc1-2* (DL1102; expresses MPK1^{HA}). Protein kinase assays (top) and immunodetection of MPK1^{HA} (bottom) were carried out on parallel sets of immune complexes.

Table 1. Heat shock gene expression is normal in *mpk1* and *pkc1* mutants

Genotype	Temperature (°C)	HSE-driven β -galactosidase activity (relative units)		STRE-driven catalase T activity (relative units)
		SSA1/HSP26	synthetic	
Wild type	23	1.0	3.2	1.0
	39	19.0	20.6	23.9
<i>mpk1</i> Δ	23	1.2	4.2	1.1
	39	17.8	21.8	22.0
Wild type + salt	23	1.2	N.T.	2.1
	39	14.4	N.T.	25.8
<i>pkc1</i> Δ + salt	23	4.6	N.T.	5.7
	39	30.8	N.T.	36.7

Strains were wild-type (DL1161 and DL1162), *mpk1* Δ (DL1157 and DL1158), and *pkc1* Δ (DL1186). Plasmid pB11 (Susek and Lindquist 1990) expresses *E. coli lacZ* under the control of the yeast *CYC1* promoter driven by upstream activating sequences from *HSP26* and *SSA1*. Plasmid CS11 is a similar construct with four synthetic HSEs fused to the *CYC1* promoter. The elevated basal level of catalase T activity in the presence of salt (0.5 M potassium acetate) results from transcriptional activation of *CTT1* by high osmolarity (Schuller et al. 1994). Each value for β -galactosidase activity is the mean of at least two experiments. (N.T.) Not tested.

gene in an *mpk1* Δ mutant and a *pkc1* Δ mutant. The *CTT1* gene encodes the major yeast catalase (catalase T) and is induced transcriptionally by stresses such as heat shock, hydrogen peroxide, and high osmolarity (Marchler et al. 1993; Schuller et al. 1994). Both mutants displayed normal induction of catalase activity in response to heat shock (Table 1), indicating that the *PKC1* pathway is not responsible for heat shock-induced expression of genes under STRE control. Therefore, thermal activation of this pathway must serve some other purpose.

Stresses other than heat shock that cause proteins to unfold also induce the expression of heat shock genes. Ectopic expression of denatured proteins in cells is sufficient to stimulate heat shock gene expression (Parsell and Lindquist 1994), suggesting that unfolded proteins constitute the primary signal for activation of HSF. We examined the activation of MPK1^{HA} in response to several protein denaturing stresses. Treatment of cells growing at 23°C with 400 μ M hydrogen peroxide, 6% ethanol, or 750 μ M sodium arsenite, all of which induce expression of HSE-driven genes (Sanchez et al. 1992), failed to activate MPK1^{HA} (less than twice basal activity). Therefore, it is not likely that unfolded proteins resulting from heat shock are the signal that stimulates the *PKC1* pathway.

The PKC1-regulated signaling pathway is activated in response to plasma membrane stretch

If the *PKC1* pathway is not activated by the same signal that induces heat shock gene expression, then to what aspect of thermal stress might this pathway respond? In addition to unfolding proteins, heat shock also increases the fluidity of membranes. Because mutants in the *PKC1*-regulated pathway undergo cell lysis resulting from a deficiency in cell wall construction, we speculated that the activating signal might be generated as a consequence of weakness in the cell wall created during

growth at high temperature, possibly attributable to increased membrane fluidity. The observed delay in MPK1^{HA} activation in response to elevated temperature can be explained by the development of weakness in the cell wall during growth under thermal stress. One mechanism by which the cell might detect weakness in the wall is by measuring stretch of the membrane. The yeast plasma membrane possesses mechanosensitive ion channels that are activated by stretch (Gustin et al. 1988).

As a first test of the model that the *PKC1*-regulated pathway is activated by plasma membrane stretch, we examined the effect of altering the extracellular osmolarity on MPK1^{HA} activity. One prediction of this model is that growth in the presence of high extracellular osmolarity, which prevents outward stretch of the plasma membrane, should prevent activation of MPK1^{HA}. Cells growing in the presence or absence of 0.5 M potassium acetate were subjected to a 30-min temperature shift from 24°C to 39°C. Figure 4 shows that activation of MPK1^{HA} by elevated temperature was prevented to a large degree in cells growing in the high osmolarity medium (cf. lanes 2 and 5). This inhibition of MPK1^{HA} activation was not specific to potassium acetate but was also caused by 10% sorbitol (not shown), indicating that this is a general effect of increased osmolarity. Moreover, prevention of heat-induced MPK1^{HA} activation by high osmolarity is not a consequence of activating the high osmolarity glycerol (HOG) response pathway, because potassium acetate prevented activation of MPK1^{HA} in a mutant in this pathway (*pbs2* Δ ; Fig. 4, cf. lanes 7 and 9). Although mechanisms other than the HOG pathway may exist to signal high extracellular osmolarity, these results suggest that elevated osmolarity prevents MPK1^{HA} activation as a direct consequence of the increased external pressure on the plasma membrane. This conclusion is supported by the observation that increasing the osmolarity at the time of temperature shift also prevented MPK1^{HA} activation effectively (not shown).

Figure 4. Growth in high extracellular osmolarity prevents temperature-induced activation of MPK1. Cultures of wild-type cells (DL807; expresses MPK1^{HA}) and a *pbs2Δ* mutant (DL1049; expresses MPK1^{HA}), growing at 24°C in YEPD or YEPD supplemented with 0.5 M potassium acetate (KOAc), were subjected to a 30-min heat shock (at 39°C) before extraction and immunoprecipitation of MPK1^{HA}. Protein kinase assays (*top*) and immunodetection of MPK1^{HA} (*bottom*) were carried out on parallel sets of immune complexes.

Another prediction of this model is that MPK1^{HA} should be activated by environmental changes that induce plasma membrane stretch. We used two mechanisms for inducing stretch. First, we tested the ability of a rapid reduction in extracellular osmolarity to activate MPK1^{HA}. Cells growing at 23°C in high osmolarity medium (YEPD + 1 M sorbitol) were diluted with YEPD (to 0.2 M sorbitol). MPK1^{HA} was activated transiently by this treatment (Fig. 5A). Although activation was weak as compared with heat shock activation, the kinetics were strikingly rapid, with activity peaking 1 min after osmotic shock. This is the expected result if MPK1^{HA} is activated in response to plasma membrane stretch induced by the instantaneous reduction in external osmolarity. MPK1^{HA} activity returned to the preactivation level within 5–10 min after this shock. This too is expected, because as the cell opens membrane ion channels to collapse the osmotic gradient, the outward pressure on the membrane is relieved.

The second mechanism used to induce stretch of the plasma membrane was treatment of cells with chlorpromazine (CPZ), a cationic amphipath, which preferentially inserts into the cytoplasmic leaflet of the plasma membrane lipid bilayer. This agent is very effective at inducing inward membrane stretch in vivo (Sheetz and Singer 1974), and activating mechanosensitive ion channels in vitro (Martinac et al. 1990). Although these ion channels can be activated by either inward or outward membrane stretch, we chose an agent that induces in-

ward stretch to avoid constraint by the cell wall. Treatment of cells growing at 25°C with 250 μM CPZ strongly activated MPK1^{HA} (Fig. 5B, cf. lanes 1 and 4). The level of MPK1^{HA} activity in the presence of CPZ was nearly as high as that induced by heat shock (cf. lanes 2 and 4). CPZ has also been reported to inhibit calmodulin in a Ca²⁺-dependent manner (Babu et al. 1988). To exclude the possibility that CPZ exerts its activating effect on MPK1^{HA} through an indirect mechanism involving calmodulin, we tested a strain with a mutant calmodulin that is unable to bind Ca²⁺ (*cmd1-6*; Geiser et al. 1991), and therefore, should not interact with CPZ. This mutant behaved identically to the isogenic *CMD1*⁺ strain with respect to MPK1^{HA} activation by CPZ (Fig. 5B; lanes 5,7). Although it is possible that CPZ has additional effects on cells that could result in the indirect activation of MPK1^{HA}, it seems most likely that the observed activation results from CPZ-induced membrane stretch.

Intracellular localization of MPK1

Because some MAPKs have been reported to translocate between the cytoplasm and the nucleus (Chen et al. 1992; Sanghera et al. 1992; Lenormand et al. 1993), we examined the intracellular localization of MPK1^{HA} under various thermal conditions. Indirect immunofluorescence of MPK1^{HA} revealed that this MAPK resides in

Figure 5. Membrane stretch activates MPK1. (A) Activation of MPK1 by hypo-osmotic stress. Wild-type cells (DL922; expresses MPK1^{HA}) growing at 23°C in YEPD supplemented with 1 M sorbitol were diluted with YEPD to a final concentration of 0.2 M sorbitol. MPK1^{HA} was immunoprecipitated from extracts of samples taken at the indicated times. Protein kinase assays (*top*) and immunodetection of MPK1^{HA} (*bottom*) were carried out on parallel sets of immune complexes. (B) Activation of MPK1 by chlorpromazine (CPZ). Cultures of wild type (DL1210; expresses MPK1^{HA}) or *cmd1-6* mutant (DL1211; expresses MPK1^{HA}) cells growing at 25°C in YEPD were either shifted to 39°C for 30 min (lanes 2,3,6) or treated with 250 μM CPZ for 2 hr (lanes 4,7). Protein kinase assays (*top*) and immunodetection of MPK1^{HA} (*bottom*) were carried out on parallel sets of immune complexes.

both the cytoplasm and the nucleus in growing cells. MPK1^{HA} appeared to be concentrated in the nucleus in cells growing at 24°C (Fig. 6A). In contrast, in cells growing at 37°C, staining was distributed more uniformly between the two compartments (Fig. 6C). The pattern of localization in cells shifted from 24°C to 39°C for 30 min (Fig. 6B) resembled the uniform staining observed after long-term cultivation at high temperature, except that MPK1^{HA} appeared to be excluded from the enlarged vacuoles that develop in response to heat shock. Because expression of MPK1^{HA} is not altered by heat shock or prolonged growth at elevated temperature (data not shown), the observed staining patterns reflect a redistribution of nuclear MPK1 to the cytoplasm at high temperature.

Discussion

The PKC1-regulated signaling pathway is important for induced thermotolerance

Heat shock disrupts a variety of cellular structures and processes by causing protein unfolding and increased membrane fluidity (for review, see Parsell and Lindquist 1994). The yeast heat shock response is a complex process that allows cells to grow at elevated temperatures and survive brief exposure to extreme temperatures. The best studied aspect of the heat shock response is the phenomenon of induced thermotolerance. This refers to

the resistance to killing by high temperatures (e.g., 50°C) that is acquired by brief pretreatment at a moderately elevated temperature (e.g., 37°C). The thermally induced synthesis of a set of proteins, known collectively as heat shock proteins (HSPs), plays an essential role in induced thermotolerance (Parsell and Lindquist 1994). HSPs function in two ways to prevent the accumulation and aggregation of denatured proteins. Some act as molecular chaperones that promote protein refolding, whereas others are involved in the degradation of denatured proteins (for review, see Morimoto et al. 1994).

In this study we demonstrate that the PKC (encoded by *PKC1*)-regulated cell integrity signaling pathway mediates a novel aspect of the yeast heat shock response. Mutants in the protein kinases under the control of PKC1 (*BCK1*, *MKK1/MKK2*, and *MPK1*) undergo cell lysis because of a deficiency in cell wall construction when they are induced to grow at high temperatures (e.g., 37°C) (for review, see Levin and Errede 1995). We have demonstrated here that (1) such mutants are deficient in the ability to survive a severe heat shock after brief pretreatment with a mild heat shock; (2) a constitutive mutant in the *PKC1*-regulated pathway (*BCK1-20*) is partially resistant to severe heat shock, indicating that one aspect of induced thermotolerance is activated constitutively in this mutant; (3) the activity of the MAPK (encoded by *MPK1*) under the control of PKC1 is stimulated 170-fold in growing cells in response to a mild heat shock (i.e., 23°C to 39°C). This increased activity is sustained indefinitely during growth at elevated temperature; and (4) mutants in the *PKC1* pathway are not compromised for the ability to induce expression of genes under the transcriptional control of known heat shock-activated promoter elements (i.e., HSEs or STREs), indicating that this pathway plays a novel role in induced thermotolerance. To our knowledge, this is the first report linking PKC to a heat shock response.

The PKC1-regulated pathway is activated by plasma membrane stretch

Mutants in the *PKC1* pathway are deficient in remodeling their cell walls during growth. We propose that the heat-induced signal for pathway activation is generated as a consequence of weakness in the cell wall created during growth under conditions of thermal stress, perhaps resulting from increased fluidity of the plasma membrane (Fig. 7). An increase in membrane fluidity could disrupt the interaction between the plasma membrane and the cell wall, thereby adversely affecting wall biosynthesis. This model is consistent with the observation that heat shock-induced activation of MPK1 followed a delay of ~20 min, suggesting that the signal develops as a consequence of growth under the new environmental condition.

We propose further that the *PKC1*-regulated signaling pathway detects and responds to weakness in the cell wall by measuring plasma membrane stretch (Fig. 7). Three lines of evidence support this model. First, growth in high osmolarity medium greatly inhibited the heat

Figure 6. Intracellular localization of MPK1^{HA} by indirect immunofluorescence. Diploid cells, expressing MPK1^{HA} (DL1232) or untagged MPK1 (1788), were grown in log phase for 12–15 hr at 24°C (A) or 37°C (C), or shifted from growth at 24°C to 39°C for 30 min (B). Cells were stained with anti-HA antibodies (12CA5) and DAPI (to visualize DNA).

Figure 7. Model for heat shock-induced activation of the *PKC1* signaling pathway. Elevated growth temperature results in a weakened cell wall at the site of wall remodeling, possibly attributable to increased membrane fluidity. Weakness in the wall allows the plasma membrane to stretch, triggering activation of mechanosensitive (MS) ion channels. Open ion channels allow influx of cations, including Ca^{2+} , which activates directly or indirectly the *PKC1* pathway. Alternatively, pathway activation may be coupled directly to ion channel activation independently of ion flux. Activated *PKC1* stimulates a protein kinase cascade that culminates with the activation of the MPK1 MAPK.

shock-induced activation of MPK1. This effect appears to be a direct consequence of the increased external pressure exerted on the plasma membrane, because it did not require the function of the HOG pathway. Second, a reduction in extracellular osmolarity induced a very rapid, but transient, activation of MPK1, consistent with the instantaneous membrane stretch such treatment would induce. Tyrosine phosphorylation of MPK1 (as an indirect measure of MPK1 activation) has also been detected in response to reduced osmolarity (M.C. Gustin, pers. comm.). Third, treatment of cells with CPZ, an amphipathic drug that induces membrane stretch by preferentially inserting into the cytoplasmic leaflet of the lipid bilayer strongly activated MPK1. Taken in the aggregate, these results indicate that membrane stretch is sufficient to activate the *PKC1* signaling pathway.

Yeast cells possess mechanosensitive ion channels that are activated by membrane stretch and are capable of importing Ca^{2+} (Gustin et al. 1988). Yeast cells experience a rapid influx of Ca^{2+} in response to mild heat shock (H. Iida, pers. comm.) that may result from the opening of such ion channels. Because *PKC1* is related to the Ca^{2+} -activated family of PKC isoforms (Levin et al.

1990), it may be activated directly in response to Ca^{2+} influx. Alternatively, *PKC1* may be activated indirectly through the stimulation of a Ca^{2+} -dependent phospholipase C (PLC). Although such a phospholipase has been described in yeast (the product of the *PLC1* gene) (Flick and Thorner 1993; Payne and Fitzgerald-Hayes 1993; Yoko-o et al. 1993), this enzyme does not appear to function within the *PKC1*-regulated signaling pathway. It is also possible that activation of a PLC is coupled directly to ion channel activation.

It is not yet clear what types of pathway output are engaged in response to activation of the *PKC1*-regulated pathway. Persistent pathway activation in response to elevated temperature might induce the expression of genes whose products can reduce membrane fluidity, perhaps by increasing the level of saturation of the fatty acids in the membrane. Expression of genes involved in cell wall biosynthesis might also be regulated by this pathway. However, if the *PKC1* pathway regulates gene expression in response to heat shock, it must do so through a mechanism other than activation of the known heat-inducible promoter elements. Alternatively, it is possible that this pathway regulates functions other than gene expression. For example, the actin cytoskeleton, which controls polarized cell growth (Drubin 1991), may be reorganized in response to pathway activation. The observed translocation of MPK1 from the nucleus to the cytoplasm at elevated temperature suggests that this MAPK serves a function outside of the nucleus.

Transient pathway activation caused by a reduction in extracellular osmolarity may induce a qualitatively different output than that resulting from sustained activation in response to heat shock. For example, ion channel conductance may be regulated by pathway activity during the equilibration of internal and external osmolarity. This may be relevant to the observation that mammalian PKC regulates the function of some ion channels (Farley and Auerbach 1986; Madison et al. 1986; Hardy et al. 1995). Precedents exist for different responses to transient versus sustained activation of mammalian MAPKs, which may be explained by translocation of MAPKs only under conditions of sustained activation (Marshall 1995).

The *PKC1* pathway may be an appropriate model for mammalian signaling pathways that regulate stress-activated MAPKs. The JNK and p38 MAPKs are regarded as functional homologs of the yeast MAPK encoded by the *HOG1* gene because all of these enzymes are activated in response to high extracellular osmolarity (Galcheva-Gargova et al. 1994; Han et al. 1994). However, unlike *HOG1*, the mammalian enzymes are activated by a variety of other stresses, including heat shock (Hibi et al. 1993; Kyriakis et al. 1994; Rouse et al. 1994). The requirement for PKC activation in the stimulation of JNK in T cells (Su et al. 1994) suggests that a pathway similar to that regulated by *PKC1* might function in mammalian cells to signal membrane stretch. Mammalian cells possess mechanosensitive ion channels (Martinac et al. 1990), which presumably allow them to detect plasma membrane stretch. These cells may monitor the state of

their extracellular matrices by a mechanism similar to that by which we propose yeast cells to monitor the state of their cell walls.

Other signals that lead to activation of PKC1

The *PKC1*-regulated pathway is also required for cell integrity during mating factor-induced morphogenesis (Errede et al. 1995). Mutants in *BCK1* or *MPK1* lyse when treated with mating factor at temperatures that are permissive for vegetative growth. The kinetics of cell lysis indicate that mutant cells die at the onset of morphogenesis. We have found that *MPK1* is activated in response to treatment of cells with mating factor and that this activation is also coincident with the onset of morphogenesis (Errede et al. 1995). These observations may be explained by the finding that during mating factor-induced morphogenesis, a large region of the cell wall undergoes remodeling in a relatively short period, potentially creating weakened areas that would result in stimulation of the *PKC1* pathway.

Cells depleted of *PKC1* lyse at a uniform point in the cell cycle, with small-to-medium sized buds (Levin et al. 1990). We have proposed that this is because a deficiency in cell wall remodeling is manifested most severely in the early stages of bud formation (Levin and Bartlett-Heubusch 1992), during which growth is polarized to the bud tip (Farkas et al. 1974; Lew and Reed 1993). This is supported by the observation that *PKC1*-depleted cells

lyse specifically at their bud tips (Levin et al. 1994). Therefore, it might be expected that *PKC1*-regulated pathway activity would fluctuate with the cell cycle. In addition, a mutation in *MPK1* was isolated in a search for mutations that enhance the growth defect of a temperature-sensitive allele of the *CDC28* cell cycle gene (Mazoni et al. 1993), further suggesting that the cell cycle engine plays a role in cell wall construction. However, measurements of *MPK1* activity in synchronized cells have failed to reveal fluctuations through the cell cycle (Y. Kamada and D.E. Levin, unpubl.). This observation seems to preclude any model in which *PKC1* activity is regulated by the cell cycle engine.

Materials and methods

Strains, growth conditions, transformations, and nucleic acid manipulation

All yeast strains used in this study (Table 2) were derivatives of EG123 (*MAT α leu2-3,112 ura3-52 trp1-1 his4 can1^r*) (Siliciano and Tatchell 1984), except CRY1 and JGY149, which were the gift of Trisha Davis (University of Washington, Seattle). Yeast cultures were grown in YEPD medium (1% Bacto-yeast extract, 2% Bacto-peptone, 2% glucose). Synthetic minimal medium (SD; Rose et al. 1990) supplemented with the appropriate nutrients was used to select for plasmid maintenance and gene replacements. The *pbs2 Δ ::LEU2* plasmid (pMA11) used to delete the *PBS2* gene was the gift of M.C. Gustin (Rice University, Houston, TX). A 3.7-kb *SacI*–*SpeI* fragment was used for yeast transformation. Yeast transformation was by the lithium ace-

Table 2. *S. cerevisiae* strains

Strain	Genotype	Source
EG123	<i>MATα leu2-3,112 ura3-52 trp1-1 his4 can1^r</i>	I. Herskowitz (University of California)
1783	<i>MATα EG123</i>	I. Herskowitz (University of California)
1788	<i>MATα/MATα isogenic diploid of EG123</i>	I. Herskowitz (University of California)
DL251	<i>MATα/MATα 1788 <i>bck1Δ::URA3/bck1Δ::URA3</i></i>	Lee and Levin (1992)
DL456	<i>MATα/MATα 1788 <i>mpk1Δ::TRP1/mpk1Δ::TRP1</i></i>	Lee et al. (1993)
DL744	<i>MATα/MATα DL456 (YE351[<i>MPK1</i>])</i>	this study
DL807	<i>MATα/MATα DL456 (YE352[<i>MPK1::HA</i>])</i>	this study
DL922	<i>MATα/MATα DL456 (YE351[<i>MPK1::HA</i>])</i>	this study
DL923	<i>MATα/MATα DL922 <i>bck1Δ::URA3/bck1Δ::URA3</i></i>	this study
DL1049	<i>MATα/MATα DL807 <i>pbs2Δ::LEU2/pbs2Δ::LEU2</i></i>	this study
DL1101	<i>MATα 1783 (YE352[<i>MPK1::HA</i>])</i>	this study
DL1102	<i>MATα DL1101 <i>pkc1Δ::LEU2</i> (YCp50ΔURA3[<i>pkc1-2</i>])</i>	this study
DL1157	<i>MATα/MATα DL456 (pB11)</i>	this study
DL1158	<i>MATα/MATα DL456 (CS11)</i>	this study
DL1161	<i>MATα/MATα 1788 (pB11)</i>	this study
DL1162	<i>MATα/MATα 1788 (CS11)</i>	this study
DL1182	<i>MATα/MATα 1788 (YE352[<i>mpk1-T190A</i>, <i>Y192F::HA</i>])</i>	this study
DL1183	<i>MATα/MATα 1788 <i>BCK1-20/BCK1-20</i> (YE352[<i>MPK1::HA</i>])</i>	this study
DL1186	<i>MATα 1783 <i>pkc1Δ::LEU2</i> (pB11)</i>	this study
DL1210	<i>MATα ade2-1 his3-11,15 leu2-3,112 trp1-1 ura3-1 can1-100</i> (YE351[<i>MPK1::HA</i>])	T. Davis (University of Washington)
DL1211	<i>MATα DL1210 <i>cmd1-6</i></i>	T. Davis (University of Washington)
DL1232	<i>MATα/MATα DL456 (YCp50LEU2[<i>MPK1::HA</i>])</i>	this study

tate method (Ito et al. 1983). General genetic manipulation of yeast cells was carried out as described previously (Rose et al. 1990). Induced thermotolerance of yeast strains was tested as described by Sanchez et al. (1992). Oligonucleotide labeling, hybridization, and DNA sequence analysis were also carried out as described previously (Levin et al. 1987).

E. coli DH5 α (Hanahan 1983), HB101 (Boyer and Roilland-Dussoix 1969), and TG1 (Sambrook et al. 1989) were used for the propagation of all plasmids and phage. Phage M13mp18 (Norlander et al. 1983) was used to generate single-stranded template DNA for site-directed mutagenesis and DNA sequence analysis. *E. coli* cells were cultured in Luria broth or YT medium (0.75% Bacto-tryptone, 0.5% Bacto-yeast extract, 0.5% NaCl) and transformed or infected with M13 by standard methods (Maniatis et al. 1982).

Epitope tagging of MPK1

MPK1 was tagged at its carboxyl terminus with an epitope from the influenza virus HA protein (Wilson et al. 1984). In the first of two steps, a 2.7-kb *EcoRI*–*SphI* fragment bearing various alleles of the entire *MPK1* gene in M13mp18 was subjected to mutagenesis with the primer 5'-GATAGAAAATATTTTGCGGCCGCTAGGACAAAACTA-3'. This resulted in the placement of a *NotI* site immediately 5' to the translational terminator. Next, the *NotI* fragment from pSM491 (gift of B. Futcher, Cold Spring Harbor Laboratory), which carries the 9-amino-acid HA epitope in tandem triplication, was subcloned into the newly created *NotI* site of *MPK1*.

Indirect immunofluorescence

Indirect immunofluorescence was done by modification of the procedure described in Berkower et al. (1994). Cultures of log phase cells ($\sim 10^8$ cells) growing in YEPD were fixed by direct addition of formaldehyde to the medium to a final concentration of 4%. Cells were fixed for 30 min at room temperature before harvest by centrifugation. Fixed cells were washed twice with 5 ml of KP buffer (0.1 M potassium phosphate, pH 6.5) and resuspended in 2 ml of KPS buffer (KP with 250 mM sorbitol). To remove the cell wall, 0.5 ml of cell suspension was incubated with 20 μ g/ml of Zymolyase 100T (ICN) and 0.2% β -mecaptoethanol at 30°C for 30 min. Spheroplasts were washed with 5 ml of KPS and harvested by gentle centrifugation (3000g for 2 min) and resuspended in 0.5 ml of KPST (KPS with 0.05% Tween 20). Fixed spheroplasts were adsorbed to polylysine-coated multiwell slides for 15 min, and blocked for 15 min with PBST–BSA [0.18% K_2HPO_4 , 0.03% KH_2PO_4 (pH 7.3), 0.8% NaCl, 0.02% KCl, 0.05% Tween 20, and 0.1% BSA]. The wells were incubated with 15 μ l of primary antibody (12CA5) diluted 1:1000 in PBST–BSA for 2 hr at room temperature. After four washes with PBST–BSA, wells were incubated for 2 hr with 15 μ l of secondary antibody (FITC-conjugated goat-anti-mouse; Boehringer Mannheim) diluted 1:500. After four washes with PBST–BSA, cells were stained with 1 μ g/ml of DAPI in PBS (PBST–BSA without Tween 20 or BSA) for 5 min. The wells were washed once with PBS before mounting medium (Rose et al. 1990) was added.

Treatment of cells and preparation of extracts

Cells grown to $A_{600}=0.5$ – 1.0 in YEPD medium were treated with various stresses before harvesting. For heat shock, cells growing at room temperature (23°C–25°C) were diluted 1:1 with fresh medium prewarmed to 55°C and then maintained at 39°C for the indicated times. Control cells were diluted with medium

at room temperature. The cell response was terminated by further diluting the culture (1:1) with ice-cold stop mix (0.9% NaCl, 1 mM NaN_3 , 10 mM EDTA, and 50 mM NaF) (Surana et al. 1991).

Cells were harvested from 50 ml of medium by centrifugation at 3000g for 5 min and washed once with cold stop mix. The cell pellet was suspended in 0.4 ml of ice-cold lysis buffer [50 mM Tris-HCl (pH 7.5), 150 mM NaCl, 5 mM EDTA, 5 mM EGTA, 0.2 mM Na_3VO_4 , 50 mM KF, 30 mM sodium pyrophosphate, 15 mM *p*-nitrophenylphosphate, 20 μ g/ml of leupeptin, 20 μ g/ml of benzamidine, 10 μ g/ml of pepstatin A, 40 μ g/ml of aprotinin, and 1 mM PMSF]. An equal volume of glass beads (0.3 mm diam.) and 2 μ l of Antifoam A (Sigma) were added to this suspension, and cells were broken by vigorous vortexing for 4 min at 4°C. The beads and cell debris were removed by centrifugation at 13,000g at 4°C, and the supernatant was further clarified by two additional 10-min centrifugations. Glycerol was added to the resulting lysate to a final concentration of 30% before storage at -20°C . Protein concentrations of cell extracts were measured as described (Bradford 1976) with Bio-Rad protein determination reagent.

Immunoprecipitations

MPK1^{HA} was immunoprecipitated from cell extracts using the 12CA5 monoclonal antibody (BabCo), which recognizes the HA epitope. Cell extracts (50 μ g of protein) were added to immunoprecipitation buffer (lysis buffer without protease inhibitors, with 1% NP-40 and sodium pyrophosphate at pH 7.5) to a final volume of 0.5 ml, to which was added 2 μ l of mAb 12CA5 raw ascites diluted 1:10 in Tris-buffered saline (TBS; 50 mM Tris-HCl (pH 7.4), 8% NaCl, 0.2% KCl). This mixture was incubated for 1 hr at 4°C with gentle agitation. To the extract/antibody mixture was added 20 μ l of a 50% suspension of protein A–Sepharose beads (Sigma) equilibrated in immunoprecipitation buffer, followed by a further incubation for 1 hr at 4°C with gentle agitation. Immune complexes were washed four times with immunoprecipitation buffer (without sodium pyrophosphate or KF), twice with buffer M [25 mM MOPS (pH 7.5), 1 mM EGTA, 0.1 mM Na_3VO_4 , 15 mM *p*-nitrophenylphosphate] and once with kinase assay buffer (buffer M with 15 mM $MgCl_2$).

In vitro phosphorylation assay

Phosphorylation assays were conducted using immunoprecipitated MPK1^{HA}. Immunoprecipitates were suspended in 18 μ l of kinase assay buffer with 10 μ g of MBP (from bovine brain; Sigma). This mixture was preincubated for 3 min at 30°C before the reaction was initiated by adding 2 μ l of 1 mM [γ -³²P]ATP (16 μ Ci/nmole; ICN). After incubation for 20 min at 30°C, the reaction was terminated by addition of 30 μ l of 2 $\times$ Laemmli's sample buffer (Laemmli 1970), and samples were boiled and subjected to SDS-PAGE on 12.5% acrylamide gels. After electrophoresis, gels were immersed in 12.5% trichloroacetic acid for 1 hr, followed by three washes for 30 min in 10% methanol, 10% acetic acid, to reduce background. Dried gels were subjected to autoradiography or phosphorimaging (BAS 1000, Fujix).

Immunoblots

Immunoprecipitated complexes were subjected to SDS-PAGE on 7.5% acrylamide gels followed by electroblotting onto nitrocellulose using a Hoefer semidry blotting system. Blots were blocked by incubation for 30 min at room temperature in TBS with 4% nonfat dry milk. Blots were then washed twice for 10

min with TBST (TBS with 0.05% Tween 20) and incubated with the mAb 12CA5 diluted 1:10,000 in TBST for 1.5 hr. After two washes with TBST, blots were incubated for 1 hr with peroxide-linked secondary antibody (Amersham) diluted 1:10,000 in TBST. After two final washes with TBST, blots were developed using the Enhanced chemiluminescence detection kit (Amersham).

β -Galactosidase and catalase assays

HSE-bearing reporter plasmids were pB11 (carries *HSP26/SSA1/CYC1* chimeric promoter, Susek and Lindquist 1990) and CS11 (carries four synthetic HSEs fused to *CYC1* promoter), which express *E. coli lacZ* in response to heat shock (gift of S. Lindquist, University of Chicago, IL). Cells bearing either of these plasmids were grown at 23°C to $A_{600} = 1.0$ in the presence or absence of 0.5 M potassium acetate, and heat-shocked at 39°C by diluting 1:1 in medium prewarmed to 55°C. After 30 min at 39°C, the temperature of the culture was reduced to 23°C for 1.5 hr to allow expression of β -galactosidase (and catalase) without further stress. The temperature was reduced to protect *PKC1* pathway mutants from thermal killing. Cell extracts were made and β -galactosidase assays were done with 15 μ g of protein, as described (Rose et al. 1990). The same extracts were used for catalase T assays, in which 30 μ g of protein was added to 1 ml of catalase buffer [100 mM Na_2HPO_4 (pH 7.0), 0.01% Triton X-100, 0.05% H_2O_2]. The decrease in A_{240} was followed for 2–3 min.

Acknowledgments

We thank K. Lee, S. Lindquist, T. Davis, and M. Gustin for strains and plasmids; H. Iida and M. Gustin for communication of results before publication; and D. Loayza and K. Fujimura-Kamada for helpful discussions. This work was supported by grants from the National Institutes of Health (GM48533) and the American Cancer Society (FRA-446) to D.E.L., and by Center Grant ES-03819 from the National Institutes of Environmental Health Sciences.

The publication costs of this article were defrayed in part by payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 USC section 1734 solely to indicate this fact.

References

- Ahn, N.G., R. Seger, and E.G. Krebs. 1992. The MAP kinase activator. *Curr. Opin. Cell Biol.* **4**: 992–999.
- Angel, P., M. Imagawa, R. Chiu, B. Stein, R.J. Imbra, H.J. Rahmsdorf, C. Jonat, P. Herrlich, and M. Karin. 1987. Phorbol ester-inducible genes contain a common cis element recognized by a TPA-modulated trans-acting factor. *Cell* **49**: 729–739.
- Angel, P., K. Hattori, T. Smeal, and M. Karin. 1988. The *jun* proto-oncogene is positively autoregulated by its product, Jun/AP-1. *Cell* **55**: 875–885.
- Babu, Y.S., C.E. Bugg, and W.J. Cok. 1988. Structure of calmodulin refined at 2.2 Å resolution. *J. Mol. Biol.* **204**: 191–204.
- Baeuerle, P.A. and D. Baltimore. 1988. Activation of DNA-binding activity in an apparently cytoplasmic precursor of the NF- κ B transcription factor. *Cell* **53**: 211–217.
- Berkower, C., D. Loayza, and S. Michaelis. 1994. Metabolic instability and constitutive endocytosis of STE6, the α -factor transporter of *Saccharomyces cerevisiae*. *Mol. Biol. Cell.* **5**: 1185–1198.
- Berridge, M.J. and R.F. Irvine. 1984. Inositol trisphosphate, a novel second messenger in cellular signal transduction. *Nature* **312**: 315–321.
- Boyer, H. and D. Roiland-Dussoix. 1969. A complementation analysis of the restriction and modification of DNA in *Escherichia coli*. *J. Mol. Biol.* **41**: 459–472.
- Bradford, M.M. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* **76**: 248–254.
- Brewster, J.L., T. de Valoir, N.D. Dwyer, E. Winter, and M.C. Gustin. 1993. An osmosensing signal transduction pathway in yeast. *Science* **259**: 1760–1763.
- Chen, R.-H., C. Sarnecki, and J. Blenis. 1992. Nuclear localization and regulation of *erk*- and *rsk*-encoded protein kinases. *Mol. Cell. Biol.* **12**: 915–927.
- Chiu, R., M. Imagawa, R.J. Imbra, J.R. Bockover, and M. Karin. 1987. Multiple cis- and trans-acting elements mediate the transcriptional response to phorbol esters. *Nature* **329**: 648–651.
- Cobb, M.H., T.G. Boulton, and D.J. Robbins. 1991. Extracellular signal-regulated kinases: ERKs in progress. *Cell Regul.* **2**: 965–978.
- Colamonici, O.R., J.B. Trepel, C.A. Vidal, and L.M. Neckers. 1986. Phorbol ester induces c-myc gene transcription in stem cell line K-562. *Mol. Cell. Biol.* **6**: 1847–1850.
- Coughlin, S.R., W.M. Lee, P.W. Williams, G.M. Giels, and L.T. Williams. 1985. c-myc gene expression is stimulated by agents that activate protein kinase C and does not account for the mitogenic effect of PDGF. *Cell* **43**: 243–251.
- Dekker, L.V. and P.J. Parker. 1994. Protein kinase C—A question of specificity. *Trends Biochem. Sci.* **19**: 73–77.
- Derijard, B., M. Hibi, I.-H. Wu, T. Barrett, B. Su., T. Deng, M. Karin, and R.J. Davis. 1994. JNK1: A protein kinase stimulated by UV light and Ha-Ras that binds and phosphorylates the c-Jun activation domain. *Cell* **76**: 1025–1037.
- Drubin, D.G. 1991. Development of cell polarity in budding yeast. *Cell* **65**: 1093–1096.
- Errede, B., R.M. Cade, B.M. Yasar, Y. Kamada, D.E. Levin, K. Irie, and K. Matsumoto. 1995. Dynamics and organization of MAP kinase signal pathways. *Mol. Reprod. Dev.* (in press).
- Farkas, V., J. Kovarik, A. Kosinova, and S. Bauer. 1974. Autoradiographic study of mannan incorporation into the growing cell walls of *Saccharomyces cerevisiae*. *J. Bacteriol.* **117**: 265–269.
- Farley, J. and S. Auerbach. 1986. Protein kinase C activation induces conductance changes in Hermissenda photoreceptors like those seen in associative learning. *Nature* **319**: 220–223.
- Flick, J.S. and J. Thorner. 1993. Genetic and biochemical characterization of a phosphatidylinositol-specific phospholipase C in *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* **13**: 5861–5876.
- Galcheva-Gargova, Z., B. Derijard, I.-H. Wu, and R. Davis. 1994. An osmosensing signal transduction pathway in mammalian cells. *Science* **265**: 806–808.
- Gartner, A., K. Nasmyth, and G. Ammerer. 1992. Signal transduction in *Saccharomyces cerevisiae* requires tyrosine and threonine phosphorylation of FUS3 and KSS1. *Genes & Dev.* **6**: 1280–1292.
- Geiser, J.R., D. van Tuinen, S.E. Brockerhoff, M.M. Neff, and T.N. Davis. 1991. Can calmodulin function without binding calcium? *Cell* **65**: 949–959.
- Greenberg, M.E. and E.B. Ziff. 1984. Stimulation of 3T3 cells induces transcription of the c-fos proto-oncogene. *Nature* **311**: 433–438.

- Gustin, M.C., X.-L. Zhou, B. Martinac, and C. Kung. 1988. A mechanosensitive ion channel in the yeast plasma membrane. *Science* **242**: 762–765.
- Han, J., J.-D. Lee, L. Bibbs, and R.J. Ulevitch. 1994. A MAP kinase targeted by endotoxin and hyperosmolarity in mammalian cells. *Science* **265**: 808–811.
- Hanahan, D. 1983. Studies on transformation of *Escherichia coli* with plasmids. *J. Mol. Biol.* **166**: 557–580.
- Hardy, S.P., H.R. Goodfellow, M.A. Valverde, D.R. Gill, F.V. Sepulveda, and C.F. Higgins. 1995. Protein kinase C-mediated phosphorylation of the human multidrug resistance P-glycoprotein regulates cell volume-activated chloride channels. *EMBO J.* **14**: 68–75.
- Herskowitz, I. 1995. MAP kinase pathways in yeast: for mating and more. *Cell* **80**: 187–197.
- Hibi, M., A. Lin, T. Smeal, A. Minden, and M. Karin. 1993. Identification of an oncoprotein- and UV-responsive protein kinase that binds and potentiates the c-Jun activation domain. *Genes & Dev.* **7**: 2135–2148.
- Hill, C.S. and R. Treisman. 1995. Transcriptional regulation of extracellular signals: Mechanisms and specificity. *Cell* **80**: 199–211.
- Hokin, L.E. 1985. Receptors and phosphoinositol-generated second messengers. *Annu. Rev. Biochem.* **54**: 205–235.
- Imagawa, M., R. Chiu, and M. Karin. 1987. Transcription factor AP-2 mediates induction by two different signal-transduction pathways: Protein kinase C and cAMP. *Cell* **51**: 251–260.
- Imbra, R.J. and M. Karin. 1986. Phorbol ester induces the transcriptional stimulatory activity of the SV40 enhancer. *Nature* **323**: 555–558.
- Irie, K., M. Takase, K.S. Lee, D.E. Levin, H. Araki, K. Matsumoto, and Y. Oshima. 1993. *MKK1* and *MKK2*, which encode *S. cerevisiae* MAP kinase-kinase homologs function in the pathway mediated by protein kinase C. *Mol. Cell. Biol.* **13**: 3076–3083.
- Ito, H., Y. Fukuda, K. Murata, and A. Kimura. 1983. Transformation of intact yeast cells treated with alkali cations. *J. Bacteriol.* **153**: 163–168.
- Kaibuchi, K., Y. Takai, and Y. Nishizuka. 1985. Protein kinase C and calcium ion in mitogenic response of macrophage-depleted human peripheral lymphocytes. *J. Biol. Chem.* **260**: 1366–1369.
- Kelly, K., B.H. Cochran, C.D. Stiles, and P. Leder. 1983. Cell-specific regulation of the c-myc gene by lymphocyte mitogens and platelet-derived growth factor. *Cell* **35**: 603–610.
- Kishimoto, A., Y. Takai, T. Mori, U. Kikkawa, and Y. Nishizuka. 1980. Activation of calcium and phospholipid dependent protein kinase by diacylglycerol: Its possible relation to phosphatidyl inositol turnover. *J. Biol. Chem.* **255**: 2273–2276.
- Kruijer, W., J.A. Cooper, T. Hunter, and I. Verma. 1984. Platelet-derived growth factor induces rapid but transient expression of the c-fos gene and protein. *Nature* **312**: 711–720.
- Kyriakis, J.M., P. Banerjee, E. Nikolakaki, T. Dai, E.I. Rubie, M.F. Ahmad, J. Avruch, and J.R. Woodgett. 1994. The stress-activated protein kinase subfamily of c-jun kinases. *Nature* **369**: 156–160.
- Laemmli, U.K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**: 680–685.
- Lee, K.S. and D.E. Levin. 1992. Dominant mutations in a gene encoding a putative protein kinase (*BCK1*) bypass the requirement for a *Saccharomyces cerevisiae* protein kinase C homolog. *Mol. Cell. Biol.* **12**: 172–182.
- Lee, K.S., K. Irie, Y. Gotoh, Y. Watanabe, H. Araki, E. Nishida, K. Matsumoto, and D.E. Levin. 1993. A yeast mitogen-activated protein kinase homolog (Mpk1p) mediates signalling by protein kinase C. *Mol. Cell. Biol.* **13**: 3067–3075.
- Lee, W., A. Haslinger, M. Karin, and R. Tjian. 1987. Activation of transcription by two factors that bind promoter and enhancer sequences of the human metallothionein gene and SV40. *Nature* **325**: 368–372.
- Lenormand, P., C. Sardet, G. Pages, G. L'Allemain, A. Brunet, and J. Pouyssegur. 1993. Growth factors induce nuclear translocation of MAP kinases (p42^{mapk} and p44^{mapk}) but not their activator MAP kinase kinase (p45^{mapkk}) in fibroblasts. *J. Cell Biol.* **122**: 1079–1088.
- Levin, D.E. and E. Bartlett-Heubusch. 1992. Mutants in the *S. cerevisiae* *PKC1* gene display a cell cycle-specific osmotic stability defect. *J. Cell Biol.* **116**: 1221–1229.
- Levin, D.E. and B. Errede. 1995. The proliferation of MAP kinase signalling pathways in yeast. *Curr. Opin. Cell Biol.* **7**: 197–202.
- Levin, D.E., C.I. Hammond, R.O. Ralston, and J.M. Bishop. 1987. Two yeast genes that encode unusual protein kinases. *Proc. Natl. Acad. Sci.* **84**: 6035–6039.
- Levin, D.E., F.O. Fields, R. Kunisawa, J.M. Bishop, and J. Thorner. 1990. A candidate protein kinase C gene, *PKC1*, is required for the *S. cerevisiae* cell cycle. *Cell* **62**: 213–224.
- Levin, D.E., B. Bowers, C.-Y. Chen, Y. Kamada, and M. Watanabe. 1994. Dissecting the protein kinase C/MAP kinase signalling pathway of *Saccharomyces cerevisiae*. *Cell Mol. Biol. Res.* **40**: 229–239.
- Lew, D.J. and S.I. Reed. 1993. Morphogenesis in the yeast cell cycle: Regulation by Cdc28 and cyclins. *J. Cell Biol.* **120**: 1305–1320.
- Madison, D.V., R.C. Malenka, and R.A. Nicoll. 1986. Phorbol esters block a voltage-sensitive chloride current in hippocampal pyramidal cells. *Nature* **321**: 695–697.
- Marchler, G., C. Schuler, G. Adams, and H. Ruis. 1993. A *Saccharomyces cerevisiae* UAS element controlled by protein kinase A activates transcription in response to a variety of stress conditions. *EMBO J.* **12**: 1997–2003.
- Maniatis, T., E.F. Fritsch, and J. Sambrook. 1982. *Molecular cloning: A laboratory manual*. Cold Spring Harbor Laboratory, Cold Spring Harbor, New York.
- Marshall, C.J. 1995. Specificity of receptor tyrosine kinase signaling: Transient versus sustained extracellular signal-regulated kinase activation. *Cell* **80**: 179–185.
- Martinac, B., J. Adlerm, and C. Kung. 1990. Mechanosensitive ion channels of *E. coli* activated by amphipaths. *Nature* **348**: 261–263.
- Mazzoni, C., P. Zarzov, A. Rambourg, and C. Mann. 1993. The *SLT2 (MPK1)* MAP kinase homolog is involved in polarized cell growth in *Saccharomyces cerevisiae*. *J. Cell Biol.* **123**: 1821–1833.
- Michak, H., J.H. Pierce, J. Goodnight, M.G. Kazanietz, P.M. Blumberg, and J.F. Mushinski. 1993. Phorbol ester-induced myeloid differentiation is mediated by protein kinase C- α and not by protein kinase C- β II, - ϵ , - ζ , and - η . *J. Biol. Chem.* **268**: 20,110–20,115.
- Morimoto, R.I., A. Tissieres, and C. Georgopoulos. 1994. Progress and perspectives on the biology of heat shock proteins and molecular chaperones. In *The biology of heat shock proteins and molecular chaperones* (ed. R.I. Morimoto, A. Tissieres, and C. Georgopoulos), pp. 1–30. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York.
- Munro, S. and H. Pelham. 1985. What turns on heat shock genes? *Nature* **317**: 476–478.

- Negro-Vilar, A. and E.G. Lapetina. 1985. 1,2 didecanoylglycerol and phorbol 12, 13 dibutyrate enhance anterior pituitary hormone secretion in vitro. *Endocrinology* **117**: 1559–1564.
- Nishizuka, Y. 1986. Studies and perspectives of protein kinase C. *Science* **233**: 305–312.
- . 1988. The molecular heterogeneity of protein kinase C and its implications for cellular regulation. *Nature* **334**: 661–665.
- . 1992. Intracellular signaling by hydrolysis of phospholipids and activation of protein kinase C. *Science* **258**: 607–614.
- Norrander, J., T. Kempe, and J. Messing. 1983. Construction of improved M13 vectors using oligodeoxynucleotide-directed mutagenesis. *Gene* **26**: 101–106.
- Ohmura, E. and H.G. Friesen. 1985. 12-O-tetradecanoyl phorbol-13-acetate stimulates rat growth hormone (GH) release from different pathways from that of human pancreatic GH-releasing factor. *Endocrinology* **116**: 728–733.
- Paravicini, G., M. Cooper, L. Friedli, D.J. Smith, J.-L. Carpentier, L.S. Klig, and M.A. Payton. 1992. The osmotic integrity of the yeast cell requires a functional *PKC1* gene product. *Mol. Cell. Biol.* **12**: 4896–4905.
- Parsell, D.A. and S. Lindquist. 1994. Heat shock proteins and stress tolerance. In *The biology of heat shock proteins and molecular chaperones* (ed. R.I. Morimoto, A. Tissieres, and C. Georgopoulos), pp. 457–495. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York.
- Payne, W.E. and M. Fitzgerald-Hayes. 1993. A mutation in *PLC1*, a candidate phosphoinositide-specific phospholipase C gene from *Saccharomyces cerevisiae*, causes aberrant mitotic chromosome segregation. *Mol. Cell. Biol.* **13**: 4351–4364.
- Pelech, S.L. and J.S. Sanghera. 1992. Mitogen-activated protein kinases: Versatile transducers for cell signalling. *Trends Biochem. Sci.* **17**: 233–238.
- Persons, D.A., W.O. Wilkison, R.M. Bell, and O.J. Finn. 1988. Altered growth regulation and enhanced tumorigenicity of NIH 3T3 fibroblasts transfected with protein kinase C-1 cDNA. *Cell* **52**: 447–458.
- Rose, M.D., F. Winston, and P. Hieter. 1990. *Methods in yeast genetics*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York.
- Rosengurt, E., A. Rodriguez-Pena, M. Coombs, and J. Sinnett-Smith. 1984. Diacylglycerol stimulates DNA synthesis and cell division in mouse 3T3 cells: Role of Ca^{+2} -sensitive phospholipid-dependent protein kinase. *Proc. Natl. Acad. Sci.* **81**: 5748–5752.
- Rouse, J., P. Cohen, S. Trigon, M. Morange, A. Alonso-Llamazares, D. Zamanilo, T. Hunt, and A.R. Nebreda. 1994. A novel kinase cascade triggered by stress and heat shock that stimulates MAPKAP kinase-2 and phosphorylation of the small heat shock proteins. *Cell* **78**: 1027–1037.
- Sambrook, J., E.F. Fritsch, and T. Maniatis. 1989. *Molecular cloning: A laboratory manual*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York.
- Sanchez, Y., J. Taulien, K.A. Borkovich, and S. Lindquist. 1992. Hsp104 is required for tolerance to many forms of stress. *EMBO J.* **11**: 2357–2364.
- Sanghera, J.S., M. Peter, E.A. Nigg, and S.L. Pelech. 1992. Immunological characterization of avian MAP kinases: Evidence for nuclear localization. *Mol. Biol. Cell* **3**: 775–787.
- Schuller, C., J.L. Brewster, M.R. Alexander, M.C. Gustin, and H. Ruis. 1994. The HOG pathway controls regulation of transcription via the stress response element (STRE) of the *Saccharomyces cerevisiae* *CTT1* gene. *EMBO J.* **13**: 4382–4389.
- Sheetz, M.P. and S.J. Singer. 1974. Biological membranes as bilayer couples. A molecular mechanism of drug-erythrocyte interactions. *Proc. Natl. Acad. Sci.* **71**: 4457–4461.
- Siliciano, P.G. and K. Tatchell. 1984. Transcription and regulatory signals at the mating type locus in yeast. *Cell* **37**: 969–978.
- Sorger, P.K. and H.R.B. Pelham. 1988. Yeast heat shock factor is an essential DNA-binding protein that exhibits temperature-dependent phosphorylation. *Cell* **54**: 855–864.
- Su, B., E. Jacinto, M. Hibi, T. Kallunki, M. Karin, and Y. Ben-Neriah. 1994. JNK is involved in signal integration during costimulation of T lymphocytes. *Cell* **77**: 727–736.
- Surana, U., H. Robitsch, C. Price, T. Schuster, I. Fitch, A.B. Fletcher, and K. Nasmyth. 1991. The role of CDC28 and cyclins during mitosis in the budding yeast *S. cerevisiae*. *Cell* **65**: 145–161.
- Susek, R.E. and S. Lindquist. 1990. Transcriptional derepression of the *Saccharomyces cerevisiae* *HSP26* gene during heat shock. *Mol. Cell. Biol.* **10**: 6362–6273.
- Takai, Y., A. Kishimoto, U. Kikkawa, T. Mori, and Y. Nishizuka. 1979. Unsaturated diacylglycerol as a possible messenger for the activation of calcium-activated, phospholipid-dependent protein kinase system. *Biochem. Biophys. Res. Commun.* **91**: 1218–1224.
- Thomas, G. 1992. MAP kinase by any other name smells just as sweet. *Cell* **68**: 3–6.
- Torres, L., H. Martin, M.R. Garcia-Saez, J. Arroyo, M. Molina, M. Sanchez, and C. Nombela. 1991. A protein kinase gene complements the lytic phenotype of *Saccharomyces cerevisiae* *lyt2* mutants. *Mol. Microbiol.* **5**: 2845–2854.
- Wiederrecht, G., D. Seto, and C.S. Parker. 1988. Isolation of the gene encoding the *S. cerevisiae* heat shock transcription factor. *Cell* **54**: 841–853.
- Wilson, I.A., H.L. Niman, R.A. Houghten, A.R. Cherenon, M.L. Connolly, and R.A. Lerner. 1984. The structure of an antigenic determinant in a protein. *Cell* **37**: 767–778.
- Yoko-o, T., Y. Matsui, H. Yagisawa, H. Nojima, I. Uno, and A. Toh-e. 1993. The putative phosphoinositide-specific phospholipase C gene, *PLC1*, of the yeast *Saccharomyces cerevisiae* is important for cell growth. *Proc. Natl. Acad. Sci.* **90**: 1804–1808.

The protein kinase C-activated MAP kinase pathway of *Saccharomyces cerevisiae* mediates a novel aspect of the heat shock response.

Y Kamada, U S Jung, J Piotrowski, et al.

Genes Dev. 1995, **9**:

Access the most recent version at doi:[10.1101/gad.9.13.1559](https://doi.org/10.1101/gad.9.13.1559)

References

This article cites 86 articles, 31 of which can be accessed free at:
<http://genesdev.cshlp.org/content/9/13/1559.full.html#ref-list-1>

License

Email Alerting Service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or [click here](#).